INFORMATION TO USERS

This reproduction was made from a copy of a document sent to us for microfilming. While the most advanced technology has been used to photograph and reproduce this document, the quality of the reproduction is heavily dependent upon the quality of the material submitted.

The following explanation of techniques is provided to help clarify markings or notations which may appear on this reproduction.

1. The sign or "target" for pages apparently lacking from the document photographed is "Missing Page(s)". If it was possible to obtain the missing page(s) or section, they are spliced into the film along with adjacent pages. This may have necessitated cutting through an image and duplicating adjacent pages to assure complete continuity.
2. When an image on the film is obliterated with a round black mark, it is an indication of either blurred copy because of movement during exposure, duplicate copy, or copyrighted materials that should not have been filmed. For blurred pages, a good image of the page can be found in the adjacent frame. If copyrighted materials were deleted, a target note will appear listing the pages in the adjacent frame.
3. When a map, drawing or chart, etc., is part of the material being photographed, a definite method of "sectioning" the material has been followed. It is customary to begin filming at the upper left hand corner of a large sheet and to continue from left to right in equal sections with small overlaps. If necessary, sectioning is continued again—beginning below the first row and continuing on until complete.
4. For illustrations that cannot be satisfactorily reproduced by xerographic means, photographic prints can be purchased at additional cost and inserted into your xerographic copy. These prints are available upon request from the Dissertations Customer Services Department.
5. Some pages in any document may have indistinct print. In all cases the best available copy has been filmed.

**University
Microfilms
International**

300 N. Zeeb Road
Ann Arbor, MI 48106

Elmigri, Mohamed Rhuma

**RESPONSE OF BARLEY GENOTYPES TO NON-SALINE AND SALINE
ENVIRONMENTS**

The University of Arizona

PH.D. 1982

**University
Microfilms
International** 300 N. Zeeb Road, Ann Arbor, MI 48106

PLEASE NOTE:

In all cases this material has been filmed in the best possible way from the available copy.
Problems encountered with this document have been identified here with a check mark ✓.

1. Glossy photographs or pages _____
2. Colored illustrations, paper or print _____
3. Photographs with dark background _____
4. Illustrations are poor copy _____
5. Pages with black marks, not original copy _____
6. Print shows through as there is text on both sides of page _____
7. Indistinct, broken or small print on several pages ✓ _____
8. Print exceeds margin requirements _____
9. Tightly bound copy with print lost in spine _____
10. Computer printout pages with indistinct print _____
11. Page(s) _____ lacking when material received, and not available from school or author.
12. Page(s) _____ seem to be missing in numbering only as text follows.
13. Two pages numbered _____. Text follows.
14. Curling and wrinkled pages _____
15. Other _____

University
Microfilms
International

RESPONSE OF BARLEY GENOTYPES TO NON-SALINE
AND SALINE ENVIRONMENTS

by

Mohamed Rhuma Elmigri

A Dissertation Submitted to the Faculty of the

DEPARTMENT OF PLANT SCIENCES

In Partial Fulfillment of the Requirements
For the Degree of

DOCTOR OF PHILOSOPHY
WITH A MAJOR IN AGRONOMY AND PLANT GENETICS

In the Graduate College

THE UNIVERSITY OF ARIZONA

1 9 8 2

THE UNIVERSITY OF ARIZONA
GRADUATE COLLEGE

As members of the Final Examination Committee, we certify that we have read
the dissertation prepared by Mohamed Rhuma Elmigri
entitled Response of Barley Genotypes to Non-Saline and Saline Environments

and recommend that it be accepted as fulfilling the dissertation requirement
for the Degree of Doctor of Philosophy.

A. D. Day,
Robert Dennis

1-5-82
Date

1-5-82
Date

E. B. Jackson

1-7-82
Date

Date

Date

Final approval and acceptance of this dissertation is contingent upon the
candidate's submission of the final copy of the dissertation to the Graduate
College.

I hereby certify that I have read this dissertation prepared under my
direction and recommend that it be accepted as fulfilling the dissertation
requirement.

A. D. Day
Dissertation Director

1-5-82
Date

STATEMENT BY AUTHOR

This dissertation has been submitted in partial fulfillment of requirements for an advanced degree at The University of Arizona and is deposited in The University Library to be made available to borrowers under the rules of the Library.

Brief quotations from this dissertation are allowable without special permission, provided that accurate acknowledgement of source is made. Requests for permission for extended quotation from or reproduction of this manuscript in whole or in part may be granted by the head of the major department or the Dean of the Graduate College when in his judgement the proposed use of the material is in the interests of scholarship. In all other instances, however, permission must be obtained from the author.

SIGNED: _____

M. R. Slinger

ACKNOWLEDGEMENTS

The author expresses his sincere appreciation to Dr. A. D. Day, his Major Advisor, for his stimulating suggestions, constructive criticism, continuous encouragement, and overall supervision of the field and laboratory research.

Gratitude is extended to Dr. Fred Turner, Superintendent of the Safford Experiment Station, Safford, Arizona for his suggestions, advice, and for the privilege of using the field research facilities on the Safford Experiment Station.

Sincere thanks are expressed to Dr. R. E. Dennis and Dr. E. B. Jackson for their detailed reviews of the manuscripts for the Ph.D. Dissertation and for serving on the Ph.D. Examination Committee.

Appreciation is also extended to Dr. P. G. Bartels and Dr. K. Matsuda for representing Botany, which was the Minor Area of Specialization in this Ph.D. Program.

I owe much to my family and friends for their support of my Ph.D. Program and to The University of El-Fateh, Tripoli, Libya for sponsoring my graduate studies at The University of Arizona.

To all others who contributed in any way and are not mentioned here, I am deeply grateful.

TABLE OF CONTENTS

	Page
LIST OF TABLES	v
ABSTRACT	vii
INTRODUCTION	1
LITERATURE REVIEW	3
Temperature as a Factor in Salt Tolerance	3
Specific Salinity Effects	4
Photosynthesis and Growth	4
Water Availability	5
Root Growth	6
Nutrient Uptake	7
General Effects of Salinity on Growth and Development of	
Cereal Grains	8
Barley	8
Rice	10
Corn and Sorghum	11
Cultivar Differences in Response to Salinity	11
Yield and Yield Components	13
MATERIALS AND METHODS	15
RESULTS AND DISCUSSION	19
Plant Height	19
Lodging	22
Heads per Hill	32
Seeds per Head	33
Seed Weight	34
Grain Yield	34
Straw Yield	37
Grain to Straw Ratio	38
Planting to Flowering	38
Flowering to Maturity	39
CONCLUSIONS	41
LITERATURE CITED	43

LIST OF TABLES

Table	Page
1. Typical chemical characteristics of Grabe Clay Loam Soil irrigated with river water at the Safford Experiment Station, Safford, Arizona in 1976	16
2. Typical chemical characteristics of Grabe Clay Loam Soil irrigated with well water at the Safford Experiment Station, Safford, Arizona in 1976	17
3. Typical chemical properties of river water at the Safford Experiment Station, Safford, Arizona in 1976	20
4. Typical chemical properties of well water at the Safford Experiment Station, Safford, Arizona in 1976	21
5. Analysis of variance for plant height, lodging, heads per hill, seeds per head, and seed weight for 15 barley genotypes and varieties grown in two different environments at Safford, Arizona in 1976 and 1977	23
6. Combined analysis of variance for plant height, lodging, heads per hill, seeds per head, and seed weight for 15 barley genotypes and varieties grown in two different environments at Safford, Arizona in 1976 and 1977	24
7. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with river water in 1976	25
8. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with river water in 1977	26
9. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with river water in 1976 and 1977	27
10. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with well water in 1976	28

LIST OF TABLES---Continued

Table	Page
11. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with well water in 1977	29
12. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with well water in 1976 and 1977	30
13. Correlations between grain yield per hill and 9 characteristics for 15 barley genotypes and varieties grown in two environments at Safford, Arizona in 1976 and 1977	31
14. Analysis of variance for grain yield per hill, straw yield per hill, grain to straw ratio, days from planting to flowering, and days from flowering to maturity for 15 barley genotypes and varieties grown in two different environments at Safford, Arizona in 1976 and 1977	35
15. Combined analysis of variance for grain yield per hill, straw yield per hill, grain to straw ratio, days from planting to flowering, and days from flowering to maturity for 15 barley genotypes and varieties grown in two different environments at Safford, Arizona in 1976 and 1977	36

ABSTRACT

A 2-year study (1976-1977) was conducted at the Safford Experiment Station, Safford, Arizona to investigate the response of barley (Hordeum vulgare L.) genotypes to both non-saline and saline environments. The soil type was a Grabe Clay Loam. One environment had received only river irrigation water for the previous ten years and throughout the experiment (non-saline environment) and the other environment had been irrigated with only well water for the previous ten years and throughout the experiment (saline environment). Fifteen barley genotypes were grown in each environment each year. The following data were recorded for each genotype each year: (1) plant height, (2) lodging, (3) number of heads per hill, (4) number of seeds per head, (5) seed weight, (6) grain yield, (7) straw yield, (8) grain-to-straw ratio, (9) days from planting to flowering, and (10) days from flowering to maturity.

The exchange capacity, soluble ions, and ESP of the soil irrigated with well water were all much higher than the exchange capacity, soluble ions, and ESP of the soil irrigated with river water. The soluble salts, calcium, magnesium, sodium, chloride, sulfate, bicarbonate, and sodium adsorption ratio were all much higher in well irrigation water than were the same chemical properties in river irrigation water. The foregoing soil and water chemical properties

indicate that the non-saline environment should be much more conducive to optimum plant growth than the saline environment.

Most of the barley genotypes germinated more uniformly, grew better, and produced more forage and grain in the non-saline environment than they did in the saline environment. It required a longer period for barley to reach maturity in the saline environment than it did in the non-saline environment. When the data from the two years were combined, there were positive correlations between grain yield and plant height, number of heads per unit area, and straw yield in both non-saline and saline environments. Since there were significant differences between barley genotypes in a number of growth and yield characteristics in both environments, it should be possible to develop improved barley cultivars for both non-saline and saline environments using plant breeding techniques.

INTRODUCTION

The loose material covering the continents of the earth is called soil. Soil, which was formed through the ages by meteorological, geological, and biological actions on rock, is a living entity that is continually changing as a result of the forces of climate. The soil not only lives but it continually renews life. Animal and plant residues decay into simpler constituents and nutrient elements are made available for new life in a perpetual cycle.

Soils differ, just as plants and animals differ. They have easily recognizable characteristics which classify them into distinct bodies of nature. Soils acquire their individual properties from the forces which act upon them. For example, desert soils in the southwestern United States are usually alkaline in nature and contain large quantities of soluble salts, while the soils in the humid Northeast are normally acidic and have had many of the soluble salts leached down below the plant root zone. The accumulation of soluble salts in desert soils may become so intense that it actually interferes with normal plant growth. The danger from soluble salt accumulation in the root zone of plants in semiarid regions requires that a certain amount of irrigation water be used for leaching the salts down below the root zone. In humid climates, leaching is accomplished naturally by the high rainfall.

The accumulation of soluble salts in soils in semiarid regions is frequently further complicated by the presence of high concentrations of soluble salts in the normal irrigation water, especially if it is pumped from wells in the local areas. Well irrigation water usually contains higher concentrations of soluble salts than does river irrigation water from the accumulation of surface rainfall.

In the southwestern United States and in similar semiarid regions throughout the world the concentrations of soluble salts in the soil and in the irrigation water are usually so high that they interfere with the normal growth of most field crop plants.

LITERATURE REVIEW

Agronomic and horticultural crops vary in their tolerance to salts. However, yields of the most salt tolerant species and cultivars are reduced when the exchangeable sodium percentage (ESP) exceeds 40-60 (Richards, 1954). Soils of irrigated deserts frequently contain salts in excess of 3000 ppm and with ESP values sufficiently high to reduce yields (Fuller, 1965). Dennis et al. (1978) suggest the use of the most salt-tolerant cultivars of barley (Hordeum vulgare L.), where salt is expected to be a problem. Although salinity can usually be controlled, there is an urgent need to develop more salt-tolerant cultivars for use in the irrigated deserts of the world.

Temperature as a Factor in Salt Tolerance

Magistard et al. (1943) found that salt-tolerance varied with temperature and that with increasing temperature and salinity, decreases in crop yields occurred. Seed germination of soybean (Glycine max L. Merrill) was inhibited by 0.2% NaCl solution at 30°C but not by 0.3% NaCl solution at 15°C. The interaction of temperature and salinity on germination of sugarbeets (Beta vulgaris L.) was highly significant. Salinity had little effect on germination of sugarbeet seed at 10°C and 15°C but the inhibition on germination increased as temperature increased to 40°C. Francois and Goodin (1972) found that salinity was increasingly damaging to sugarbeets as temperature increased from 25°C to 40°C. In their experiments using several

different varieties, they determined that the germination of the sugar-beet cultivar 'U.S.H2' was most adversely affected by salinity.

Ahi and Powers (1938) conducted experiments using alfalfa (Medicago sativa L.) in which salinity and temperature were variables. They found that alfalfa germination and seedling emergence were greatly reduced by the combination of high salinity and high temperature. Brown and Hayward (1956) continued these studies with alfalfa and noted that the average yield for all cultivars was reduced to 79% of the control by their low salt treatment and 60% by the intermediate salt treatment. High relative humidity increased the salt-tolerance of barley and corn (Zea mays L.) but did not affect the tolerance of wheat (Triticum vulgare Vill aestivum L.) to salt (Hoffman and Jobes, 1978).

Specific Salinity Effects

Photosynthesis and Growth

Boyers (1965) reported that high levels of NaCl reduced the rate of photosynthesis. He theorized that this was the result of solute effects of enzymes and that it did not result from stomatal closure. Nieman (1962) in similar studies, however, found that photosynthesis rates were not affected when he increased osmotic stress with NaCl. This was true for all of the 12 different crop plants studied. Results obtained by Ackerson and Youngner (1975) agreed with those of Nieman.

Abel and Mackenzie (1964) stated that high salinity in the soil resulted in increased plant mortality, increased leaf necrosis, reduced

green leaf color, decreased stem production, decreased seed yield, and decreased seed quality. The negative effects of salinity on the growth of cotton (Gossypium hirsutum L.) plants were studied by Thomas (1980). Regression coefficients developed from his data suggested that an increase of 1 bar in total soil water suction decreased growth of cotton stems by 3.5cm during the 70 to 85-day growth period and 0.9cm during the 85 to 115-day growth period.

Sepaskhah and Boersma (1979) found that the rates of elongation of wheat leaves decreased at an accelerated rate as the osmotic potential of the water used for irrigation increased. Bernstein (1961) determined that the fresh weights of tops and roots of cotton were decreased by increasing the percentage of NaCl in the solution culture. Hassan et al. (1970) obtained results that differ from those of most researchers. Their experiments indicated that moderate soil salinity may enhance metabolism and increase dry matter production.

Water Availability

Langdale and Thomas (1971) related the growth responses of several plants to salt concentrations in the soil. They found a strong negative correlation between growth and salinity, and that the decrease in growth rates resulted largely from the physiological unavailability of water and/or from specific ion toxicities. In similar studies, Donovan and Day (1969) found that restricted crop growth in saline soils may be due to either reduction in moisture availability, toxic effects of specific ions, or salt-induced nutritional deficiencies. The first reaction of plants to high salt was reduced water entry which

caused decreased growth because of the inability of subcellular organelles, such as mitochondria, to withstand the internal drought conditions (Bernstein, 1961).

Bower and Tamimi (1979) studied the effects of salinity on the moisture content of plants. They found that shoots of cultivars grown in saline solutions had a lower water content than shoots grown in non-saline solutions, indicating that salt injury involved water stress. Narale, Subramanyan, and Mukherjee (1969) found that growth depression attributed to salinity was caused by the kind as well as the quantity of the salt present in the root environment. Negative effects of specific ions resulted primarily from imbalances in the concentration of various ion species. The nonspecific deleterious effects resulting from high salinity were caused by the total concentration of salt, regardless of the species present.

Hopkins (1939) concluded that the effects of salinity on plant growth were similar to those resulting from drought stress. Plants grown in a saline environment were shorter as a result of a shortening of the internodes. Examination of the shortened internodes suggested that this reduced growth resulted from smaller cells.

Root Growth

Gerard (1978) measured root growth of sugarcane (Saccharum officinarum L.) in mm/cm^2 of soil. He found that root growth increased with increases in salinity of irrigation water from 1.1 to 8 mmhos/cm. In spite of the fact that there were increases in root growth, top

growth of plants was reduced by 50% at the highest salt concentration used in these studies.

In bermudagrass (Cynodon dactylon (L.) Pers), Ackerson and Youngner (1975) found that dry weight of tops decreased, while dry weight of roots and total non-structural carbohydrate concentration of crowns increased, with increased salinity of the culture solution. These researchers also noted that the root/shoot ratio of several halophytes was relatively high, compared with that of glucophytes grown under saline conditions. This suggested that root growth of halophytes may be less affected by salinity than root growth of glucophytes. The response of plants to different levels of salinity may be conditioned by a small area in the root that contains more or less salt than the soil (Black, 1957).

Nutrient Uptake

There is a significant negative correlation between soil salinity, and the uptake of P, K, Ca, Fe, and Cu by all parts of barley plants but a significant positive correlation between soil salinity and the uptake of Mn, and Na (Hassan et al., 1970). For most irrigated desert soils, nitrogen is the principal fertilizer element applied (Dennis et al., 1978). Water-stress induced by a high salt content of the soil resulted in the inhibition of nitrogen metabolism (Wadleigh and Ayers, 1945).

General Effects of Salinity on
Growth and Development of Cereal Grains

All agronomic and horticultural crops are affected by salinity levels in the soil. However, they differ greatly in their tolerance to salt (Richards, 1954). In general, plants are more tolerant of salt as they approach maturity (Lunin, Gallatin, and Batchelder, 1963). Plants are most susceptible to salt during the germination and seedling stages although other stages are affected (Maliwal and Paliwal, 1967). This effect of salinity on plant growth was studied by Pearson and Bernstein (1959) and Ehler (1958). These researchers found that salinity inhibited growth more severely at early stages of plant development than at later stages. They also found that vegetative stages of growth were less affected by salinity than were reproductive stages. Harlan and Martini (1936) found that barley was particularly susceptible to salt in the soil and water during germination. Ayers, Brown, and Wadleigh (1952) concluded that high levels of salinity in the soil and irrigation water had a greater negative effect on the reproductive stages of growth of wheat plants than it did on the vegetative stages.

Barley

Barley is one of the most salt-tolerant agronomic crops (Ayers, Brown, and Wadleigh, 1952). Because of this characteristic, it is adapted to wide areas throughout the world (Donovan and Day, 1969).

Although barley is salt-tolerant, it is affected by the level of dissolved salts in the environment at germination and at other stages of growth (Ayers and Hayward, 1948).

In Arizona, most barley is of the spring type but it is grown as a winter-annual (Dennis et al., 1978). Barley at the lower elevations of Arizona is usually planted in December or January and it matures in May or June of the following year. Since most of the growth of barley plants occurs during the cooler time of the year, they are able to withstand relatively high levels of salt in the soil and irrigation water (Dennis et al., 1978).

The date of planting for barley, as for other crops, greatly affects yield potential. Beard (1961) found that highest yields were obtained from early planting. Woodward (1956) found that early planted barley plants had stiffer straw, larger heads, and a higher test weight, since the plants had a longer period for each of the several stages of growth. In similar experiments, Day and Thompson (1970) obtained the highest yields for barley when plantings in central Arizona were made in mid-November and with a seeding rate of 25 kg/ha. The lower seeding rate they used for barley decreased the number of fertile heads per unit of area but increased the number of seeds per head and the weight of individual seeds.

Wagenet et al. (1980) studied the effect of irrigation frequency and its interaction with salinity on the yield of barley. They found that grain yields were reduced with decreasing irrigation frequency (2 to 8 days) and that increasing the salinity of irrigation

water from 0.5 to 13.5 mmhos/cm also decreased grain yield. Increasing the seeding rate helped to compensate for the decreased germination and the increased seedling mortality that occurred in saline-sodic soils (Dargon, Abrol and Bhumla, 1974).

The method of planting also has an impact on the growth and development of barley plants. Day, Turner, and Kirkpatrick (1971) found that drilling on beds resulted in higher grain yields than planting with two rows per bed. They also determined that planting 54 kg/ha of seed resulted in higher grain yields than planting 28 kg/ha. In related experiments, they also determined that when barley was grown in saline soil and irrigated with high-salt water, highest yields were obtained when stands were obtained by planting on beds at a low rate of seeding.

Lodging of barley plants is also an important factor related to yield loss when plantings are made in the irrigated deserts of the Southwest. When plants were artificially lodged to 90° (Day, 1957) the yield of grain, test weight, and weight per 1000 seeds were decreased.

Rice

Salinity of soil and water is most damaging to rice (Oryza sativa L.) plants when it is imposed early in the life cycle (Pearson and Bernstein, 1959). In addition, high salts appear to affect pollen germination, and therefore, fertilization of rice florets. Kapp (1947) also studied the interaction of stage of growth of rice plants and salinity levels. He found that high soil salinity at planting time was a more severe treatment for rice than high salinity when seedlings

were six weeks old. Kaddah, Lehman, and Robinson (1973) obtained similar results. They also found that relatively high salinity after the boot stage did not significantly affect the yield of rice plants.

Narale, Subramanyan, and Mukherjee (1969) concluded that grain yield, and vegetative growth were markedly hampered at an EC_e of 7.2 and 9.8 mmhos/cm, and that rice could germinate normally in a medium having an EC_e up to 4.5 mmhos/cm.

Corn and Sorghum

Kaddah and Ghowail (1964) conducted experiments using corn. They found that the tolerance of corn to salt was moderate and that it increased as plants grew older. Lyles and Fanning (1964) used sorghum (Sorghum vulgare L.) plants in similar research. They found that increases in soil salinity delayed and decreased seedling emergence.

Cultivar Differences in Response to Salinity

Ayers (1953) determined that barley cultivars differ in their tolerance to salinity. These early studies have been repeated in greater detail by other researchers. Shourbagy and Wallace (1965) found that high levels of sodium in the soil and water decreased yields for four of five cultivars of barley. In their studies, 'Arivat' was most severely affected by high salinity while 'Firlbecks III' was least affected. Donovan and Day (1969) found that the cultivar 'California Mariout' was exceptionally tolerant of high levels of salinity, while Arivat had a very low level of salt-tolerance. These researchers also concluded that at moderate salinity levels most barley varieties had

uniform seedling emergence but that at high levels of salinity only the most salt-tolerant cultivars produced satisfactory seedling growth.

Taylor, Young, and Rivera (1975) found that cultivars of sorghum also differ significantly in their tolerance to salt. They concluded that plant breeders have an opportunity to develop new cultivars with extraordinary salt-tolerance. Abel and Mackenzie (1964) found that salinity decreased the percent and rate of seedling emergence and that there was a difference between varieties of soybeans in the decrease in the rate of emergence.

Dewey (1960) studied the relative salt tolerance of twenty-five different strains of Agropyron (sp.). He reported that differences in salt tolerance among strains indicated that selection would prove successful in developing more salt-tolerant strains. Dewey (1960) also stated that salt-tolerance is not absolute and depends on environmental factors, such as relative humidity and management of irrigation and fertility.

Hunt (1965) noted that success in increasing salt-tolerance in a given species depends on heritable variation in the species and the effectiveness of selection techniques. Little or no relationship was found between salt-tolerance of seedlings and salt-tolerance of mature plants.

Dobrenz et al. (1980) were very successful in increasing salt-tolerance in alfalfa. They selected plants that appeared to be unusually salt-tolerant. Three years of yield data with these

salt-tolerant alfalfas have demonstrated that their yields are higher than 'Mesa-Sirsa', the check variety.

Mineral analysis may be a way to evaluate tolerance to salinity. Shannon (1978) found that tolerance to salinity was associated with restricted accumulation of Na, Ca, and Cl. Francois (1976) determined that Na uptake increased proportionally in the shoots of tall wheatgrass (Agropyron elongatum Beauv.) with increases in salinity, whereas Cl uptake seemed to remain relatively constant and was not dependent on salt concentration in the soil and water.

Yield and Yield Components

Stickler and Pauli (1961) stated that the yield components for barley and wheat were: (1) number of heads per unit area, (2) number of seeds per head, and (3) weight of individual seeds. Knott and Talukdar (1971) found that individual seed weight of wheat was positively correlated with yield and negatively correlated with the number of kernels per plot. On the average, wheat lines with high seed weight produced the higher yields of grain.

Results obtained by McNeal (1960) differed from those of Knott and Talukdar (1971). McNeal stated that of the three yield components, kernel weight was the least important in determining yield. McNeal, Smith, and Berg (1974) found that yield increases for wheat could be obtained by increasing one or more of the three components of yield.

Grafius (1956) stated that for oats yield components were: (1) panicles per unit area, (2) number of kernels per panicle, and

(3) kernel weight. Fonseca and Patterson (1965) also conducted experiments with oats and found that the three yield components above were highly correlated with grain yield. Frey (1959) working with spring oats, found that the yield variables enumerated by Grafius were influenced by cultivar, location, date of planting, and fertility.

The objectives of this research experiment were as follows:

(1) to compare the chemical characteristics of the soil and irrigation water in both non-saline and saline environments, (2) to study the general growth and response of barley genotypes to both non-saline and saline environments, (3) to determine the correlations between grain yield and plant growth characteristics of barley genotypes grown in non-saline and saline environments, and (4) to evaluate the feasibility of using plant breeding techniques to develop improved barley cultivars for both non-saline and saline environments.

MATERIALS AND METHODS

A two-year experiment (1976 thru 1977) was conducted at the Safford Experiment Station, Safford, Arizona to study the response of barley genotypes to both non-saline and saline environments. The research site was located at about 1,000 m above sea level. The soil type was a Grabe Clay Loam, which is a member of the coarse-loamy, mixed thermic family of Cumulic Haplustolls. The soil was plowed, floated, disked, dragged, and bedded to produce a good seedbed. Nitrogen fertilizer, in the form of ammonium nitrate, was applied prior to planting each year at the rate of 168 kg/ha of elemental N. The experiment was conducted in two research environments each year. One area had received only river irrigation water for the previous ten years (non-saline environment) and the other area had been irrigated with only well water for the previous ten years (saline environment). The typical chemical characteristics of Grabe Clay Loam Soil irrigated with river water compared with that irrigated with well water for ten years are presented in Tables 1 and 2, respectively. The following 15 barley genotypes were grown in each environment each year: (1) 'Mesa 273', (2) 'Mesa 269', (3) 'Yuma 32', (4) 'Yuma 6', (5) 'Mesa 322', (6) 'Yuma 52', (7) 'Yuma 166', (8) 'Mesa 300', (9) 'Mesa 383', (10) 'Yuma 122', (11) 'Arimar', (12) 'Arivat', (13) 'Mesa 497', (14) 'Arizona 1970-1', and (15) 'California Mariout'. Arimar, Arivat, and California Mariout were commercial cultivars. Arizona 1970-1 was an early, short-strawed

Table 1. Typical chemical characteristics of Grabe Clay Loam Soil irrigated with river water at the Safford Experiment Station, Safford, Arizona in 1976.

Soil depth	EC x 10 ³	Soluble ions ¹				ESP	pH
		Ca	Mg	Na	Cl		
(m)	(mmhos/cm)	(meq/l)	(meq/l)	(meq/l)	(meq/l)	(%)	(pH)
0.3	2.2	0.8	0.4	17.4	17.3	15.6	8.6
0.6	3.5	3.1	0.6	20.3	20.0	15.8	8.6
1.0	5.0	2.7	1.1	24.5	24.2	15.9	8.3
1.3	4.5	2.1	1.1	25.0	24.9	16.9	8.3
1.6	3.9	1.3	1.0	25.8	24.2	20.0	8.4
2.0	2.9	0.8	0.5	20.9	19.2	19.7	8.3

¹Determined in the saturation extract.

Table 2. Typical chemical characteristics of Grabe Clay Loam Soil irrigated with well water at the Safford Experiment Station, Safford, Arizona in 1976.

Soil depth	EC x 10 ³	Soluble ions ¹				ESP	pH
		Ca	Mg	Na	Cl		
(m)	(mmhos/cm)	(meq/l)	(meq/l)	(meq/l)	(meq/l)	(%)	(pH)
0.3	3.5	1.6	0.8	34.1	30.2	18.4	8.6
0.6	3.7	3.7	1.7	52.2	51.2	31.4	8.5
1.0	7.7	4.7	2.9	68.6	67.9	32.7	8.4
1.3	7.9	5.8	2.4	70.4	70.3	31.0	8.4
1.6	4.8	4.6	1.3	47.5	53.0	29.9	8.3
2.0	4.2	4.0	1.1	44.2	41.2	27.5	8.3

¹Determined in the saturation extract.

natural mutation from Arimar. All Mesa genotypes were thermal neutron induced mutations from Arivat and all Yuma genotypes were thermal neutron induced mutations from Arimar. The experimental design was a Split Plot Experiment with environments as main plots and genotypes as sub-plots with four replications. Each individual plot consisted of one hill which covered an area of 0.25m^2 on a standard vegetable bed. Torrie (1962) reported that the hill method of planting was used successfully in evaluating strains (genotypes) of soybeans when land was limited, the number of entries was large, or the seed supply was limited. The barley was planted in November at the rate of 12 seeds per hill and the grain and straw were harvested with a hand sickle the following May each year. After planting, the barley was irrigated-up to saturate the soil to a depth of 1 m and irrigated as needed throughout the growing season to insure that the plants were never stressed for moisture. A total of 1 m of irrigation water was required to grow the barley genotypes to maturity. The following data were recorded for each genotype each year: (1) plant height, (2) lodging, (3) number of heads per hill, (4) number of seeds per head, (5) seed weight, (6) grain yield, (7) straw yield, (8) grain-to-straw ratio, (9) days from planting to flowering, and (10) days from flowering to maturity. All data were analyzed using the standard analysis of variance. Means were compared using the Least Significant Differences (LSD) Test. Simple correlations were calculated between grain yield and the remaining nine characteristics observed for each genotype.

RESULTS AND DISCUSSION

It is important to point out at the beginning of the results and discussion that the growth of each of the 15 barley genotypes was evaluated in both a non-saline environment and in a saline-environment on Grabe Clay Loam Soil at Safford, Arizona. The non-saline environment was created by irrigating the soil with river water for 10 years prior to the experiment and throughout the experiment. The saline environment was created by irrigating the soil with well water for 10 years prior to the experiment and throughout the experiment. The exchange capacity, soluble ions, and ESP of the soil irrigated with well water were all much higher than the exchange capacity, soluble ions, and ESP of the soil irrigated with river water (Tables 1 and 2). The pH values for the soils in the two environments were similar. The soluble salts, calcium, magnesium, sodium, chloride, sulfate, bicarbonate, and sodium adsorption ratio were all much higher in well irrigation water than were the same chemical properties in river irrigation water (Tables 3 and 4). The foregoing soil and water chemical properties indicate that the non-saline environment should be much more conducive to optimum plant growth than the saline environment.

Plant Height

The analysis of variance for plant height for 1976 and 1977 combined showed a significant difference between environments and also

Table 3. Typical chemical properties of river water at the Safford Experiment Station, Safford, Arizona in 1976.

Chemical properties	1976
	(ppm)
Soluble salts	518.0
Calcium	29.0
Magnesium	8.7
Sodium	88.0
Chloride	107.0
Sulfate	56.0
Carbonate	0.0
Bicarbonate	222.0
Fluoride	2.2
Nitrate	5.0
Sodium Adsorption Ratio (SAR)	3.7

Table 4. Typical chemical properties of well water at the Safford Experiment Station, Safford, Arizona in 1976.

Chemical properties	1976
	(ppm)
Soluble salts	2044.0
Calcium	98.0
Magnesium	19.0
Sodium	500.0
Chloride	630.0
Sulfate	260.0
Carbonate	0.0
Bicarbonate	532.0
Fluoride	2.4
Nitrate	3.2
Sodium Adsorption Ratio (SAR)	12.1

between genotypes (Tables 5 and 6). In general, barley genotypes grown in a non-saline environment produced taller plants than did the same genotypes grown in a saline environment. This indicates that barley was unable to reach its full height potential in the saline environment. Since there were significant differences between genotypes in both the non-saline environment (Tables 7, 8, and 9) and in the saline environment (Tables 10, 11, and 12), it should be possible to select and/or breed for either taller or shorter barley in each environment. When barley is grown for forage or mulching material, tall plants are desired; however, when grain production is the primary objective, short plants are preferred. In most instances, when barley was grown in both the non-saline environment and also in the saline environment, there was a positive correlation between grain yield and plant height (Table 13). This suggests that when a plant breeder selects tall barley plants he is also selecting for high grain yield at the same time.

Lodging

The 1976 and 1977 combined analysis of variance for lodging did not show a significant difference between environments; however, it did indicate a significant difference between genotypes (Tables 5 and 6). This indicated that barley genotypes were as susceptible to lodging in a non-saline environment as they were in a saline environment. In general, it is desirable to grow barley with as low a lodging percentage as possible in all environments because as the lodging percentage increases, the general quality of the plant

Table 5. Analysis of variance for plant height, lodging heads per hill, seeds per head, and seed weight for 15 barley genotypes and varieties grown in two different environments at Safford, Arizona in 1976 and 1977.

Source of variation	df	Plant height		Lodging		Heads per hill		Seeds per head		Seed weight	
		MS	F-ratio	MS	F-ratio	MS	F-ratio	MS	F-ratio	MS	F-ratio
Replications (R)	3	247.58		567.97		56.18		32.84		7.49	
Environments (E)	1	8213.40	30.12*	0.07	0.00	3673.84	8.16	2214.34	50.93**	380.02	110.15**
Error a	3	272.70		248.88		450.12		43.48		3.45	
Genotypes (G)	14	878.66	15.22**	594.86	4.67**	739.26	2.77**	67.59	1.35	207.41	32.77**
GE	14	42.48	0.74	222.37	1.75*	342.61	1.28	63.78	1.27	22.53	3.56**
GER	42	29.40	0.51	48.03	0.38	188.39	0.71	28.39	0.57	6.60	1.04
Error b	162	57.74		127.38		267.07		50.25		6.33	

* Significant at the 5% level of probability.

** Significant at the 1% level of probability.

Table 6. Combined analysis of variance for plant height, lodging, heads per hill, seeds per head, and seed weight for 15 barley genotypes and varieties grown in two different environments at Safford, Arizona in 1976 and 1977.

Source of variation	df	Plant Height		Lodging		Heads per hill		Seeds per head		Seed weight	
		MS	F-ratio	MS	F-ratio	MS	F-ratio	MS	F-ratio	MS	F-ratio
Replications	3	247.58		587.97		56.18		32.84		7.49	
Environments (E)	1	8213.40	30.12*	0.07	0.00	3673.84	8.16	2214.34	50.93**	380.02	110.15**
Error a	3	272.70		248.88		450.12		43.48		3.45	
Years (Y)	1	912.60	8.24*	3936.60	13.25*	1111.20	50.67**	1219.50	10.48*	56.42	4.67
YE	1	1717.35	15.50**	2561.07	8.62*	6273.04	28.61**	885.50	7.61*	120.42	11.17*
Error b	6	110.81		297.20		219.30		116.42		10.78	
Genotypes (G)	14	878.66	26.67**	594.86	8.41**	739.26	4.45**	67.59	2.27**	207.41	39.76**
GE	14	42.48	1.29	222.37	3.15**	342.61	2.06*	63.78	2.14*	22.53	4.32**
GY	14	88.75	2.69**	52.67	0.75	153.89	0.93	83.45	2.80**	8.82	1.69
GYE	14	36.95	1.12	125.55	1.78*	172.74	1.04	24.99	0.84	4.80	0.92
Error c	168	32.95		70.69		166.10		29.83		5.22	

* Significant at the 5% level of probability.

** Significant at the 1% level of probability.

Table 7. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with river water in 1976.

Treatments (Genotypes)	Plant height	Lodging	Heads per hill	Seeds per head	Seed weight	Grain yield per hill	Straw yield per hill	Grain to straw ratio	Planting to flowering	Flowering to maturity
	(cm)	(%)	(no.)	(no.)	(g/1000)	(g)	(g)	(ratio)	(day)	(day)
Mesa 273	69 ab*	8 bc	50 ab	29 bc	45 bc	72 abc	55 bc	1.2 a	106 b	63 c
Mesa 269	73 ab	7 bc	77 ab	32 ab	46 ab	116 ab	101 ab	1.3 a	106 b	62 d
Yuma 32	67 ab	9 bc	81 a	31 abc	42 bc	106 ab	112 ab	1.1 a	105 c	63 c
Yuma 6	68 ab	6 bc	65 ab	38 ab	38 cd	96 ab	75 bc	1.3 a	106 b	62 d
Mesa 322	75 a	3 bc	61 ab	39 a	43 bc	102 ab	89 ab	1.2 a	106 b	62 d
Yuma 52	43 c	0 bc	71 ab	25 bc	33 e	56 bc	56 bc	1.1 a	107 a	62 d
Yuma 166	74 a	3 bc	52 ab	36 ab	44 bc	82 abc	64 bc	1.3 a	107 a	61 e
Mesa 300	65 b	2 bc	64 ab	37 ab	43 bc	102 ab	79 abc	1.4 a	105 c	63 c
Mesa 383	75 a	2 bc	66 ab	36 ab	47 ab	107 ab	72 bc	1.5 a	107 a	61 e
Yuma 122	76 a	13 bc	75 ab	36 ab	41 cd	109 ab	106 ab	1.0 a	105 c	63 c
Arimar	78 a	17 b	71 ab	36 ab	50 a	127 ab	107 ab	1.2 a	106 b	62 d
Arivat	78 a	13 bc	82 a	35 ab	46 ab	130 a	127 a	1.1 a	106 b	62 d
Mesa 497	79 a	6 bc	53 ab	36 ab	48 ab	90 ab	78 abc	1.2 a	107 a	61 e
Arizona 1970-1	62 b	6 bc	55 ab	28 bc	48 ab	75 abc	53 bc	1.5 a	102 e	67 a
Calif. Mariout	73 ab	54 a	57 ab	31 abc	49 ab	87 abc	94 ab	1.0 a	103 d	65 b
LSD	8	14	21	7	4	32	35	0.6	1	1

* Means followed by the same letter are not significantly different at 5% level of probability.

Table 8. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with river water in 1977.

Treatments (Genotypes)	Plant height	Lodging	Heads per hill	Seeds per head	Seed weight	Grain yield per hill	Straw yield per hill	Grain to straw ratio	Planting to flowering	Flowering to maturity
	(cm)	(%)	(no.)	(no.)	(g/1000)	(g)	(g)	(ratio)	(day)	(day)
Mesa 273	62 abc*	12 b	42 ab	38 b	47 ab	73 bc	39 abc	1.9 ab	87 ab	31 bc
Mesa 269	64 ab	12 b	49 ab	39 b	49 ab	92 ab	50 ab	1.9 ab	85 bc	32 b
Yuma 32	63 ab	14 ab	57 a	42 ab	43 bc	98 ab	57 ab	1.8 ab	86 bc	31 bc
Yuma 6	62 abc	7 bc	33 bc	44 ab	40 cd	51 bc	35 bc	1.6 ab	88 ab	30 bc
Mesa 322	57 bc	7 bc	35 abc	41 ab	45 bc	64 bc	35 bc	1.9 ab	86 bc	31 bc
Yuma 52	44 cd	1 bc	40 ab	40 b	34 e	56 c	39 abc	1.5 ab	88 ab	29 c
Yuma 166	70 ab	11 b	36 abc	41 ab	43 bc	62 bc	41 abc	1.5 ab	89 a	28 d
Mesa 300	51 cd	2 bc	37 abc	39 b	41 cd	59 bc	29 bc	2.0 ab	86 bc	31 bc
Mesa 383	61 bc	13 ab	46 ab	38 b	46 b	80 abc	39 abc	2.1 ab	86 bc	32 b
Yuma 122	69 ab	11 b	41 ab	52 a	41 cd	86 ab	46 ab	1.9 ab	86 bc	31 bc
Arimar	68 ab	21 ab	47 ab	42 ab	51 a	95 ab	52 ab	1.9 ab	85 bc	32 b
Arivat	74 a	15 ab	57 a	42 ab	47 ab	110 a	62 a	1.8 ab	85 bc	32 b
Mesa 497	60 bc	10 b	35 abc	44 ab	47 ab	69 bc	40 abc	1.8 ab	86 bc	31 bc
Arizona 1970-1	57 bc	7 bc	24 bc	48 ab	45 bc	50 c	27 bc	1.9 ab	82 d	35 a
Calif. Mariout	55 bc	29 a	44 ab	40 b	51 a	91 ab	41 abc	2.2 a	83 d	34 a
LSD	8	9	15	8	3	24	16	0.7	2	2

* Means followed by the same letter are not significantly different at 5% level of probability.

Table 9. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with river water in 1976 and 1977.

Treatments (Genotypes)	Plant height	Lodging	Heads per hill	Seeds per head	Seed weight	Grain yield per hill	Straw yield per hill	Grain to straw ratio	Planting to flowering	Flowering to maturity
	(cm)	(%)	(no.)	(no.)	(g/1000)	(g)	(g)	(ratio)	(day)	(day)
Mesa 273	66 ab*	10 bc	46 ab	33 ab	46 bc	73 bc	47 ab	1.6 a	96 a	47 a
Mesa 269	68 ab	9 bc	63 ab	36 ab	47 b	104 ab	75 ab	1.6 a	96 a	47 a
Yuma 32	65 ab	11 bc	69 a	36 ab	43 cd	102 ab	84 ab	1.4 a	96 a	47 a
Yuma 6	65 ab	6 bc	49 ab	41 ab	39 d	77 bc	55 ab	1.5 a	97 a	46 a
Mesa 322	66 ab	5 bc	48 ab	40 ab	44 bc	83 abc	62 ab	1.5 a	96 a	47 a
Yuma 52	43 c	1 bc	56 ab	33 ab	33 e	56 bc	47 ab	1.3 a	97 a	45 a
Yuma 166	72 ab	7 bc	44 ab	39 ab	44 bc	72 bc	53 ab	1.4 a	98 a	45 a
Mesa 300	58 ab	2 bc	50 ab	38 ab	42 cd	80 bc	54 ab	1.7 a	96 a	47 a
Mesa 383	68 ab	7 bc	56 ab	37 ab	47 b	93 ab	55 ab	1.8 a	96 a	46 a
Yuma 122	72 ab	12 bc	58 ab	44 a	41 cd	97 ab	76 ab	1.4 a	95 a	47 a
Arimar	73 ab	19 b	59 ab	39 ab	50 a	111 ab	79 ab	1.6 a	95 a	47 a
Arivat	76 a	14 bc	70 a	38 ab	46 bc	120 a	94 a	1.4 a	96 a	47 a
Mesa 497	70 ab	8 bc	44 ab	40 ab	47 b	80 bc	59 ab	1.5 a	97 a	46 a
Arizona 1970-1	59 ab	7 bc	40 ab	38 ab	47 b	63 bc	40 ab	1.7 a	92 a	51 a
Calif. Mariout	64 ab	41 a	51 ab	36 ab	50 a	89 ab	67 ab	1.6 a	93 a	49 a
LSD	11	16	23	10	3	30	36	0.7	17	25

* Means followed by the same letter are not significantly different at 5% level of probability.

Table 10. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with well water in 1976.

Treatments (Genotypes)	Plant height	Lodging	Heads per hill	Seeds per head	Seed weight	Grain yield per hill	Straw yield per hill	Grain to straw ratio	Planting to flowering	Flowering to maturity
	(cm)	(%)	(no.)	(no.)	(g/1000)	(g)	(g)	(ratio)	(day)	(day)
Mesa 273	53 bc*	0 a	48 ab	30 ab	44 ab	63 ab	39 ab	1.7 bc	107 a	61 b
Mesa 269	58 ab	1 a	45 ab	29 abc	47 a	60 ab	44 ab	1.3 bc	107 a	61 b
Yuma 32	56 abc	5 a	60 ab	29 bc	43 ab	68 ab	56 a	1.4 bc	107 a	62 a
Yuma 6	55 abc	5 a	45 ab	33 ab	39 bc	57 ab	32 ab	1.9 ab	107 a	62 a
Mesa 322	53 bc	2 a	47 ab	38 ab	41 bc	74 ab	47 ab	1.7 bc	107 a	62 a
Yuma 52	33 d	0 a	43 ab	29 bc	36 c	44 ab	51 ab	1.6 bc	107 a	61 b
Yuma 166	57 abc	7 a	58 ab	31 ab	45 ab	82 ab	61 a	1.4 bc	106 b	62 a
Mesa 300	50 bc	2 a	37 ab	33 ab	42 bc	51 ab	29 ab	1.7 bc	107 a	62 a
Mesa 383	52 bc	6 a	45 ab	34 ab	44 ab	70 ab	37 ab	1.9 ab	106 b	62 a
Yuma 122	61 ab	7 a	45 ab	23 bc	40 bc	48 ab	44 ab	1.1 bc	107 a	61 b
Arimar	54 abc	1 a	45 ab	40 a	43 ab	75 ab	35 ab	2.5 a	107 a	61 b
Arivat	66 a	9 a	69 a	32 ab	46 ab	99 a	62 a	1.6 bc	106 b	62 a
Mesa 497	59 ab	1 a	49 ab	31 ab	47 a	72 ab	45 ab	1.6 bc	106 b	62 a
Arizona 1970-1	37 d	0 a	29 ab	27 bc	46 ab	35 ab	17 ab	2.4 ab	102 c	66 a
Calif. Mariout	53 bc	5 a	46 ab	31 ab	42 bc	58 ab	34 ab	1.7 bc	106 b	62 a
LSD	8	14	21	7	4	32	35	0.6	1	1

* Means followed by the same letter are not significantly different at 5% level of probability.

Table 11. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with well water in 1977.

Treatments (Genotypes)	Plant height	Lodging	Heads per hill	Seeds per head	Seed weight	Grain yield per hill	Straw yield per hill	Grain to straw ratio	Planting to flowering	Flowering to maturity
	(cm)	(%)	(no.)	(no.)	(g/1000)	(g)	(g)	(ratio)	(day)	(day)
Mesa 273	52 bc*	19 ab	44 ab	35 ab	42 ab	57 ab	29 ab	2.0 ab	87 ab	35 ab
Mesa 269	51 bc	19 ab	30 bc	35 ab	44 a	46 ab	26 ab	1.8 ab	88 ab	33 bc
Yuma 32	61 ab	15 abc	41 abc	29 ab	42 ab	48 ab	30 ab	1.6 ab	87 ab	34 bc
Yuma 6	53 bc	14 bc	47 ab	28 ab	34 d	47 ab	30 ab	1.6 ab	89 a	32 c
Mesa 322	55 ab	15 abc	44 ab	37 ab	41 ab	67 ab	34 ab	2.1 ab	84 bc	37 a
Yuma 52	40 cd	10 bc	55 ab	28 ab	33 d	50 ab	39 ab	1.3 ab	88 ab	33 bc
Yuma 166	62 ab	20 ab	45 ab	32 ab	43 ab	64 ab	37 ab	1.7 ab	86 bc	35 ab
Mesa 300	47 cd	12 bc	36 bc	32 ab	38 c	44 ab	21 ab	2.2 ab	86 bc	35 ab
Mesa 383	55 ab	19 ab	59 a	31 ab	43 ab	77 ab	41 a	1.9 ab	84 bc	37 a
Yuma 122	62 ab	17 abc	42 abc	29 ab	38 c	45 ab	33 ab	1.4 ab	87 ab	34 bc
Arimar	55 ab	23 ab	44 ab	29 ab	43 ab	55 ab	30 ab	2.1 ab	86 bc	36 ab
Arivat	67 a	21 ab	47 ab	31 ab	43 ab	60 ab	37 ab	1.7 ab	86 bc	35 ab
Mesa 497	58 ab	18 abc	55 ab	39 a	42 ab	83 a	44 a	2.1 ab	84 bc	37 a
Arizona 1970-1	50 c	18 abc	28 bc	37 ab	43 ab	42 ab	23 ab	2.4 a	86 bc	35 ab
Calif. Mariout	52 bc	30 a	43 ab	30 ab	43 ab	53 ab	29 ab	2.0 ab	85 bc	36 ab
LSD	8	9	5	8	3	24	16	0.7	2	

* Means followed by the same letter are not significantly different at 5% level of probability.

Table 12. Average of 10 characteristics for 15 barley genotypes and varieties grown at Safford, Arizona with well water in 1976 and 1977.

Treatments (Genotypes)	Plant height	Lodging	Heads per hill	Seeds per head	Seed weight	Grain yield per hill	Straw yield per hill	Grain to straw ratio	Planting to flowering	Flowering to maturity
	(cm)	(%)	(no.)	(no.)	(g/1000)	(g)	(g)	(ratio)	(day)	(day)
Mesa 273	53 abc*	9 a	46 ab	33 ab	43 ab	60 ab	34 a	1.9 ab	97 a	48 a
Mesa 269	54 ab	10 a	38 ab	32 ab	46 a	53 ab	35 a	1.5 ab	97 a	47 a
Yuma 32	59 ab	10 a	50 ab	29 ab	43 ab	58 ab	43 a	1.5 ab	97 a	48 a
Yuma 6	54 ab	9 a	46 ab	30 ab	36 d	52 ab	31 a	1.7 ab	98 a	47 a
Mesa 322	54 ab	8 a	46 ab	38 a	41 bc	71 ab	40 a	1.9 ab	95 a	49 a
Yuma 52	37 c	5 a	49 ab	28 ab	35 d	47 ab	45 a	1.4 ab	97 a	47 a
Yuma 166	60 ab	14 a	52 a	32 ab	44 ab	73 ab	49 a	1.5 ab	96 a	49 a
Mesa 300	49 bc	7 a	36 ab	33 ab	40 bc	47 ab	25 a	1.9 ab	96 a	48 a
Mesa 383	53 abc	13 a	52 a	32 ab	44 ab	73 ab	39 a	1.9 ab	95 a	49 a
Yuma 122	62 ab	12 a	44 ab	25 ab	39 c	47 ab	38 a	1.3 ab	97 a	48 a
Arimar	55 ab	12 a	44 ab	34 ab	43 ab	65 ab	32 a	2.3 ab	96 a	48 a
Arivat	67 a	15 a	58 a	31 ab	44 ab	80 a	50 a	1.6 ab	96 a	48 a
Mesa 497	58 ab	9 a	52 a	35 ab	44 ab	78 a	44 a	1.8 ab	95 a	50 a
Arizona 1970-1	43 bc	9 a	29 ab	32 ab	44 ab	38 ab	20 a	2.4 a	94 a	50 a
Calif. Marlout	52 abc	17 a	44 ab	31 ab	42 ab	55 ab	31 a	2.0 ab	96 a	49 a
LSD	11	16	23	10	3	30	36	0.7	17	25

* Means followed by the same letter are not significantly different at 5% level of probability.

Table 13. Correlations between grain yield per hill and 9 characteristics for 15 barley genotypes and varieties grown in two environments at Safford, Arizona in 1976 and 1977.

Characteristics	Safford barley grown with well water			Safford barley grown with river water		
	1976	1977	1976-77	1976	1977	1976-77
Plant height	0.71**	0.32	0.51*	0.69**	0.59*	0.73**
Lodging	0.47	0.18	-0.11	0.08	0.71**	0.20
Heads per hill	0.85**	0.72**	0.80**	0.63*	0.90**	0.78**
Seeds per head	0.50	0.40	0.41	0.62*	-0.08	-0.19
Seed weight	0.38	0.33	0.41	0.40	0.55*	0.38
Straw yield per hill	0.68**	0.81**	0.73**	0.86**	0.91**	0.83**
Grain to straw ratio	-0.12	0.13	-0.09	-0.04	0.36	-0.35
Planting to flowering	0.31	-0.74**	0.25	0.22	-0.30	0.47
Flowering to maturity	-0.30	0.74**	0.29	-0.22	0.30	0.49

* Significant at the 5% level of probability.

** Significant at the 1% level of probability.

products decreases. Since there were significant differences in lodging between genotypes in most instances (Tables 7, 8, 9, and 11), it is reasonable to expect that new genotypes could be developed for each environment that would be more resistant to lodging than the presently available cultivars. Only in 1977 in the non-saline environment was there a significant positive correlation between grain yield and lodging percentage (Table 13). In all other instances the correlations between grain yield and lodging were not significant. These data suggest that it would be difficult for a plant breeder to select barley genotypes for high grain yield in environments similar to those present in this experiment and have any reasonable assurance that he was also selecting for increased tolerance to lodging.

Heads Per Hill

The 1976 and 1977 combined analysis of variance for number of heads per hill did not show a significant difference between environments; however, it did indicate a significant difference between genotypes (Tables 5 and 6). This suggested that barley genotypes produced as many heads per hill in a saline environment as they did in a non-saline environment. The barley grower desires a large number of heads per unit area because number of heads per unit area is one of the principal yield components that determine grain yield in small grains. The significant differences between genotypes in number of heads per hill observed in both environments in 1977 indicates that it should be possible to breed new barley cultivars for the future that will produce more heads per unit area in both non-saline and saline

environments than are produced by present commercial cultivars (Tables 8 and 11). There was a positive correlation between grain yield and number of heads per hill in all instances when barley was grown in both non-saline and saline environments at Safford, Arizona in 1976 and 1977 (Table 13). These data suggest that one of the best ways for a barley breeder to be sure that he is selecting for high grain yield is to select plants that produce a large number of heads per unit area.

Seeds Per Head

A second important grain yield component in small grains, such as barley, is number of seeds per head. The analysis of variance for number of seeds per head for 1976 and 1977 combined showed a significant difference between environments and also between genotypes (Table 6). There was also a significant difference between years in number of seeds per head (Table 6). Although the number of seeds per head does vary from year to year as a result of yearly differences in environment, barley usually produces more seeds per head in a non-saline environment than in a saline environment (Tables 9 and 12). The differences between genotypes in number of seeds per head observed in 1976 (Tables 7 and 10) should encourage barley breeders to make an effort to select new genotypes with a higher number of seeds per head because as the number of seeds per head increases, the total grain yield should also increase. Only in 1976 and in the non-saline environment was there a significant positive correlation between grain yield and number of seeds per head (Table 13).

Seed Weight

Seed weight is the third important yield component in small grains that determines the final grain yield. The 1976 and 1977 combined analysis of variance for seed weight showed a significant difference between environments and also between genotypes (Tables 5 and 6). In general, barley genotypes grown in a non-saline environment had heavier seeds than did the same genotypes grown in a saline environment. This indicated that when barley was grown in a saline environment it was unable to produce seeds as large as its genetic potential would permit it to produce in a non-saline environment. It is believed that one reason a number of commercial barley cultivars produce lower grain yields in a saline environment than the same cultivars produce in a non-saline environment is because the cultivars produce lighter seeds when the soil salts are high than when the soil salts are lower. Only in 1977 in the non-saline environment was there a significant positive correlation between grain yield and seed weight (Table 13).

Grain Yield

The principal objective of most small grain farmers is maximum grain yield per unit area of land. The analysis of variance for grain yield for 1976 and 1977 combined showed a significant difference between environments and also between genotypes (Tables 14 and 15). In general, barley genotypes grown in a non-saline environment (Tables 7, 8, and 9) produced higher grain yields than they did in a saline

Table 14. Analysis of variance for grain yield per hill, straw yield per hill, grain to straw ratio, days from planting to flowering, and days from flowering to maturity for 15 barley genotypes and varieties grown in two different environments at Safford, Arizona in 1976 and 1977.

Source of variation	df	Grain yield per hill		Straw yield per hill		Grain to straw ratio		Planting to flowering		Flowering to maturity	
		MS	F-ratio	MS	F-ratio	MS	F-ratio	MS	F-ratio	MS	F-ratio
Replications (R)	3	727.74		343.65		0.33		1.02		0.97	
Environments (E)	1	43362.82	46.37**	40690.10	90.01**	3.98	9.57	14.50	15.66**	136.50	154.78**
Error a	3	935.07		452.05		0.42		0.93		0.88	
Genotypes (G)	14	2695.30	5.84**	1663.18	2.47**	0.71	2.62**	18.59	0.13	18.73	0.06
GE	14	1201.80	2.60**	853.73	1.27	0.24	0.87	7.38	0.05	7.21	0.02
GER	42	528.03	1.14	482.16	0.72	0.27	0.99	2.47	0.02	2.49	0.01
Error b	162	461.42		673.91		0.27		147.99		316.93	

* Significant at the 5% level of probability.

** Significant at the 1% level of probability.

Table 15. Combined analysis of variance for grain yield per hill, straw yield per hill, grain to straw ratio, days from planting to flowering, and days from flowering to maturity for 15 barley genotypes and varieties grown in two different environments at Safford, Arizona in 1976 and 1977.

Source of variation	df	Grain yield per hill		Straw yield per hill		Grain to straw ratio		Planting to flowering		Flowering to maturity	
		MS	F-ratio	MS	F-ratio	MS	F-ratio	MS	F-ratio	MS	F-ratio
Replications	3	727.74		343.65		0.33		1.12		0.97	
Environment (E)	1	43362.82	46.37**	40690.10	90.01**	3.98	9.57	14.50	15.66*	136.50	154.78**
Error a	3	935.07		452.05		0.42		0.93		0.88	
Years (Y)	1	12355.35	42.89**	40794.34	545.07**	9.80	145.71**	23740.70	19992.17**	50837.70	43627.11**
YE	1	2535.00	8.80*	15440.10	206.30**	2.84	42.20**	1.20	1.01	275.20	236.17**
Error b	6	288.06		74.84		0.07		1.19		1.17	
Genotypes (G)	14	2695.30	6.52**	1663.18	4.46**	0.71	3.16**	18.59	15.25**	18.73	15.41**
GE	14	1201.80	2.91**	853.73	2.29**	0.24	1.05	7.38	6.05**	7.21	5.93**
GY	14	417.35	1.01	463.34	1.24	0.23	1.03	6.70	5.50**	6.59	5.42**
GYE	14	360.52	0.87	257.59	0.69	0.08	0.34	2.13	1.75*	2.21	1.82*
Error c	168	413.20		372.90		0.22		1.22		1.22	

* Significant at the 5% level of probability.

** Significant at the 1% level of probability.

environment (Tables 10, 11, and 12). This indicates that the different genotypes were unable to reach their full grain yield potential in a saline environment. Commercial barley growers in the southwestern United States have also observed this phenomenon on a large scale. Since there was greater variation in grain yield between the barley genotypes grown in the non-saline environment (Tables 7, 8, and 9) than there was when the same genotypes were grown in the saline environment (Tables 10, 11, and 12), a plant breeder would have a greater opportunity to select high grain yielding genotypes from these populations in a non-saline environment.

Straw Yield

The analysis of variance for straw yield for 1976 and 1977 combined showed a significant difference between environments and also between genotypes (Tables 14 and 15). Barley genotypes grown in a non-saline environment produced more straw than did the same genotypes grown in a saline environment. This indicates that a saline environment tended to reduce the total vegetative growth of barley. Since there was greater variation in straw yield between the barley genotypes when they were grown in the non-saline environment (Tables 7, 8, and 9) than there was when the same genotypes were grown in the saline environment (Tables 10, 11, and 12), a plant breeder would have a greater opportunity to select high straw yielding genotypes from these populations in a non-saline environment. A similar observation was also made regarding grain yield. High straw yields would be desirable

whenever the barley crop is being grown for hay production or to provide mulching material for the reclamation of disturbed land areas, such as copper mine wastes or coal mine wastes. In both environments, both years, and when the two years were combined there was a significant positive correlation between grain yield and straw yield (Table 13). This indicates that under the conditions of this experiment a barley breeder could select high straw yielding genotypes and be relatively confident that he was also selecting high grain producing genotypes at the same time.

Grain to Straw Ratio

The 1976 and 1977 combined analysis of variance for grain to straw ratio showed a significant difference between genotypes and between years (Tables 14 and 15). There was a tendency for the grain to straw ratios to be larger in the saline environment (Tables 10, 11, and 12) than they were in the non-saline environment (Tables 7, 8, and 9). Similar observations were also made by Day, Turner, and Kirkpatrick (1971). It is believed that the reason grain to straw ratios were larger in saline environments was because straw yields were depressed more by high soil salt than were grain yields. None of the correlations between grain yield and grain to straw ratio were significant in either environment or in either 1976 or 1977 (Table 13).

Planting to Flowering

The analysis of variance for days from planting to flowering for 1976 and 1977 combined showed a significant difference between

environments, between genotypes, and between years (Tables 14 and 15). A number of genotypes required a little longer period from planting to flowering in the saline environment (Tables 10, 11, and 12) than they required in the non-saline environment (Tables 7, 8, and 9). Most barley growers prefer as short a period as possible from planting to flowering because barleys that flower quickly usually mature earlier than barleys that require a longer period from planting to flowering. This relationship was also reported by Dennis et al. (1978). In contrast to the influence of salinity on flowering in barley observed in this study, Pearson and Bernstein (1959) found that flowering in rice was not influenced by salinity. Since the genotypes differed in the length of time required from planting to flowering, within a specific environment (Tables 7, 8, 10, and 11), this is a characteristic that could probably be altered by plant breeding techniques. In 1977 in the saline environment, there was a significant negative correlation between grain yield and the number of days from planting to flowering (Table 13). In all other instances, the correlations between grain yield and number of days from planting to flowering were not significant.

Flowering to Maturity

The 1976 and 1977 combined analysis of variance for days from flowering to maturity showed a significant difference between environments, between genotypes, and between years (Tables 14 and 15). A number of genotypes required a longer period from flowering to maturity in the saline environment (Tables 11 and 12) than they required in

the non-saline environment (Tables 8 and 9). Barley growers usually prefer cultivars that mature quickly after they reach the flowering stage because cultivars that require only a short period from flowering to maturity occupy the land area for a shorter period than do barleys that require a longer period from flowering to maturity. In regions where double-cropping is practiced, the sooner one crop can be matured and harvested the sooner a second crop can be planted on the same area of land. In the southwestern United States, many growers produce barley during the winter months and sorghum during the summer months in a double-cropping program. It should be possible to shorten the period required from flowering to maturity in new barley cultivars because a number of the barley genotypes differed in the length of time required from flowering to maturity in both the non-saline and saline environments in 1976 and 1977 (Tables 7, 8, 10, and 11). In 1977 in the saline environment, there was a significant positive correlation between grain yield and the number of days from flowering to maturity (Table 13). In all other instances, the correlations between grain yield and number of days from flowering to maturity were not significant.

CONCLUSIONS

The exchange capacity, soluble ions, and ESP of the soil irrigated with well water (saline environment) were all much higher than the exchange capacity, soluble ions, and ESP of the soil irrigated with river water (non-saline environment). The soluble salts, calcium, magnesium, sodium, chloride, sulfate, bicarbonate, and sodium adsorption ratio were all much higher in well irrigation water than in river irrigation water. The foregoing soil and water chemical properties indicate that the non-saline environment should be much more conducive to optimum plant growth and high yields of plant products than the saline environment.

Most of the barley genotypes germinated more uniformly, grew more vigorously, and produced more forage and grain in the non-saline environment. It required a longer period for barley to reach maturity in the saline environment. When the data from the two years were combined, there were positive correlations between grain yield and plant height, number of heads per unit area, and straw yield. Since there were significant differences between barley genotypes in a number of growth and yield characteristics in both environments, it should be possible to develop improved barley cultivars for both non-saline and saline environments using plant breeding techniques.

Thermal neutrons were used successfully to induce usable mutations in adapted barley cultivars in this research and may also be

useful in the development of desirable mutations in adapted barley cultivars in other similar environments throughout the semiarid regions of the world.

LITERATURE CITED

- Abel, G. H., and A. J. Mackenzie. 1964. Salt tolerance of soy-bean varieties (Glycine max L. Merrill) during germination and later growth. *Crop Sci.* 4:157-161.
- Ackerson, R. C., and V. B. Youngner. 1975. Response of bermudagrass to salinity. *Agron. J.* 67:678-681.
- Ahi, S. M., and W. L. Powers. 1938. Salt tolerance of plants at various temperatures. *Plant Physiol.* 13:767-789.
- Ayers, A. D. 1953. Germination and emergence of several varieties of barley in salinized soil culture. *Agron. J.* 45:68-71.
- Ayers, A. D., J. W. Brown, and C. H. Wadleigh. 1952. Salt tolerance of barley and wheat in soil plots receiving several salinization regimes. *Agron. J.* 44:307-310.
- Ayers, A. D., and H. E. Hayward. 1948. A method for measuring the effect of soil salinity on seed germination with observation on several crops. *Soil Sci. Amer. Proc.* 13:224-226.
- Beard, B. H. 1961. Effect of date of seeding on agronomic and malting quality characteristics of barley. *Crop Sci.* 1:300-303.
- Bernstein, L. 1961. Osmotic and adjustment of plants to saline media. I. Stead state. *Am. J. Bot.* 48:909-918.
- Black, C. A. 1957. Soil plant relationships. Wiley Book Co., Inc., New York. 332 p.
- Bower, C. A., and Y. N. Tamimi. 1979. Root adjustments associated with salt tolerance in small grains. *Agron. J.* 71:690-693.
- Boyers, J. S. 1965. Effects of osmotic water stress on metabolic rates of cotton plants with open stomata. *Plant Physiol.* 40:229-234.
- Brown, J. W., and H. E. Hayward. 1956. Salt tolerance of alfalfa varieties. *Agron. J.* 48:18-20.
- Day, A. D. 1957. Effect of lodging on yield, test weight and other seed characteristics of spring barley grown under flood irrigation. *Agron. J.* 49:536-539.

- Day, A. D., and R. K. Thompson. 1970. Dates and rates of seeding fall-planted spring barley (Hordeum vulgare L. emend Lam.) in irrigated areas. Agron. J. 62:729-731.
- Day, A. D., Fred Turner, Jr., and R. M. Kirkpatrick. 1971. Growing barley (Hordeum vulgare L.) on beds in saline soil. Agron. J. 63:768-769.
- Dargan, K. S., I. P. Abrol, and D. R. Bhumla. 1974. Performance of rice varieties in a highly saline sodic soil as influenced by plant population. Agron. J. 66:279-280.
- Dennis, R. E., R. K. Thompson, A. D. Day and E. B. Jackson. 1978. Growing barley in Arizona. Univ. of Ariz. Bull. A15, 13 p.
- Dewey, D. R. 1960. Salt tolerance of twenty five strains of Agropyron. Agron. J. 52:631-635.
- Dobrenz, A. K., J. E. Stone, M. H. Schonhorst, and Scott Riggins. 1980. Progress in the development of salt tolerant alfalfa in Arizona. Arizona Coop. Ext. Service and Agric. Exp. Station Report Series P-50:12.
- Donovan, T. J., and A. D. Day. 1969. Some effects of high salinity on germination and emergence of barley (Hordeum vulgare L.). Agron. J. 61:236-238.
- Ehler, W. L. 1958. Some effects of salinity on rice. Bot. Gaz. 122:102-104.
- Fonseca, Santiago, and F. L. Patterson. 1968. Yield component heritabilities and interrelationships in winter wheat (Triticum aestivum L.). Crop Sci. 8:614-617.
- Francois, L. E. 1976. Salt tolerance of prostrate summer cypress (Kochia prostrata). Agron. J. 68:455-456.
- Francois, L. E., and J. R. Goodin. 1972. Interaction of temperature and salinity on sugarbeet germination. Agron. J. 64:272-273.
- Frey, K. J. 1959. Yield components in oats. I. Effect of seeding date. Agron. J. 51:381-383.
- Fuller, W. H. 1965. Water, soil, and crop management principles for the control of salts. Univ. of Ariz. Bull. A-43, 28 p.
- Gerard, C. J. 1978. Root growth along plexiglas surfaces by sugarcane under different salinity conditions. Agron. J. 70:639-643.

- Grafius, J. E. 1956. Component of yield in oats: a geometrical interpretation. *Agron. J.* 48:419-423.
- Harlan, H. V., and M. L. Martini. 1936. Problems, and results in barley breeding. *USDA Yearbook of Agriculture*.
- Hassan, Nouri, A. K. Drew, V. James, Knudsen Delno, and R. A. Olson. 1970. Influence of soil salinity on production of dry matter, and uptake and distribution of nutrients in barley and corn: I. Barley (Hordeum vulgare L.). *Agron. J.* 62:43-45.
- Hoffman, G. J., and J. A. Jobes. 1978. Growth and water relations of cereal crops as influenced by salinity and relative humidity. *Agron. J.* 70:765-769.
- Hopkins, J. W. 1939. Comparative development of two wheat varieties under varying moisture supply. *Can. J. Res.* 17 Sec. C:87-96.
- Hunt, O. J. 1965. Salt tolerance in intermediate wheatgrass. *Crop Sci.* 5:407-409.
- Kaddah, M. T., and S. I. Chowail. 1964. Salinity effects on growth of corn at different stages of development. *Agron. J.* 56:214-217.
- Kaddah, M. T., W. F. Lehman, and F. E. Robinson. 1973. Tolerance of rice (Oryza sativa L.) to salt during boot, flowering, and grain filling stages. *Agron. J.* 65:845-847.
- Kapp, L. C. 1947. The effect of common salt on rice production. *Arkansas Agr. Exp. Sta. Bull.* 465.
- Knott, D. R., and B. Talukdar. 1971. Increasing seed weight in wheat and its effect on yield, yield components, and quality. *Crop Sci.* 11:280-283.
- Langdale, G. W., and J. R. Thomas. 1971. Soil salinity effect on absorption of nitrogen, phosphorus, and protein synthesis by coastal bermudagrass. *Agron. J.* 63:708-711.
- Lunin, J., M. H. Gallatin, and A. R. Batchelder. 1963. Saline irrigation of several vegetable crops at various growth stages. I. Effect on yield. *Agron. J.* 55:107-110.
- Lyles, Leon, and C. D. Fanning. 1964. Effect of presoaking, moisture tension, and soil salinity on the emergence of grain sorghum. *Agron. J.* 56:518-520.

- Magistad, O. C., A. D. Ayers, C. H. Wadleigh, and H. G. Gaugh. 1943. Effect of salt concentration, kind of salt and climate on the plant growth in sand culture. *Plant Physiol.* 18:151-166.
- Maliwal, G. L., and K. V. Paliwal. 1967. Salt tolerance studies of some varieties of wheat and barley at germination stage. *Indian J. Plant Physiol.* 10:26-35.
- McNeal, F. H. 1960. Yield components in a Lemhi x Thatcher wheat cross. *Agron. J.* 52:348-349.
- McNeal, F. H., E. P. Smith, and M. A. Berg. 1974. Plant height, grain yield and yield component relationships in spring wheat. *Agron. J.* 66:575-578.
- Narale, R. P., T. K. Subramanyan, and P. K. Mukherjee. 1969. Influence of salinity on germination, vegetative growth and grain yield of rice (*Oryza sativa* var. Dular). *Agron. J.* 61:341-344.
- Nieman, R. H. 1962. Some effects of sodium chloride on growth photosynthesis, and respiration of 12 crop plants. *Bot. Gaz.* 123:279-385.
- Pearson, G. A., and Leon Bernstein. 1959. Salinity effects at several growth stages of rice. *Agron. J.* 51:654-657.
- Richards, L. A. 1954. Diagnosis and improvement of saline and alkali soils. U.S. Department of Agriculture Handbook No. 60. 248 p.
- Shannon, M. C. 1978. Testing salt tolerance variability among tall wheatgrass lines. *Agron. J.* 70:719-722.
- Sepaskhah, A. R., and L. Boersma. 1979. Elongation of wheat leaves exposed to several levels of matric potential and NaCl-induced osmotic potential of soil water. *Agron. J.* 71:848-852.
- Shourbagy, N. E., and A. Wallace. 1965. Sodium accumulation and sodium responses of five varieties of barley. *Agron. J.* 57:449-450.
- Stickler, Fie, and A. W. Pauli. 1961. Influence of date of planting on yield and yield components in grain sorghum. *Agron. J.* 53:20-22.
- Taylor, R. M., E. F. Young Jr., and R. L. Rivera. 1975. Salt tolerance in cultivars of grain sorghum. *Crop Sci.* 15:734-735.
- Thomas, J. R. 1980. Osmotic and specific salt effects on growth of cotton. *Agron. J.* 72:407-412.

- Torrie, J. H. 1962. Comparison of hills and rows for evaluating soybean strains. Crop Sci. 2:47-49.
- Wadleigh, C. H., and A. D. Ayers. 1945. Growth and biochemical composition of bean plants as conditioned by soil moisture tension and salt concentration. Plant Physiol. 20:106-132.
- Wagenet, R. J., W. F. Campbell, A. M. Bamatraf, and D. L. Turner. 1980. Salinity, irrigation frequency, and fertilization effects on barley growth. Agron. J. 72:969-974.
- Woodward, R. W. 1956. The effect of rate and date of seeding on small grain yield. Agron. J. 48:160-162.