INFORMATION TO USERS

This material was produced from a microfilm copy of the original document. While the most advanced technological means to photograph and reproduce this document have been used, the quality is heavily dependent upon the quality of the original submitted.

The following explanation of techniques is provided to help you understand markings or patterns which may appear on this reproduction.

1. The sign or "target" for pages apparently lacking from the document photographed is "Missing Page(s)". If it was possible to obtain the missing page(s) or section, they are spliced into the film along with adjacent pages. This may have necessitated cutting thru an image and duplicating adjacent pages to insure you complete continuity.
2. When an image on the film is obliterated with a large round black mark, it is an indication that the photographer suspected that the copy may have moved during exposure and thus cause a blurred image. You will find a good image of the page in the adjacent frame.
3. When a map, drawing or chart, etc., was part of the material being photographed the photographer followed a definite method in "sectioning" the material. It is customary to begin photoing at the upper left hand corner of a large sheet and to continue photoing from left to right in equal sections with a small overlap. If necessary, sectioning is continued again -- beginning below the first row and continuing on until complete.
4. The majority of users indicate that the textual content is of greatest value, however, a somewhat higher quality reproduction could be made from "photographs" if essential to the understanding of the dissertation. Silver prints of "photographs" may be ordered at additional charge by writing the Order Department, giving the catalog number, title, author and specific pages you wish reproduced.
5. PLEASE NOTE: Some pages may have indistinct print. Filmed as received.

University Microfilms International

300 North Zeeb Road
Ann Arbor, Michigan 48106 USA
St. John's Road, Tyler's Green
High Wycombe, Bucks, England HP10 8HR

77-7327

PRICE, Mary Vaughan, 1949-
THE ROLE OF MICROHABITAT IN
STRUCTURING DESERT RODENT COMMUNITIES.

The University of Arizona, Ph.D., 1976
Ecology

Xerox University Microfilms, Ann Arbor, Michigan 48106

THE ROLE OF MICROHABITAT IN STRUCTURING
DESERT RODENT COMMUNITIES

By

Mary Vaughan Price

A Dissertation Submitted to the Faculty of the
DEPARTMENT OF ECOLOGY AND EVOLUTIONARY BIOLOGY

In Partial Fulfillment of the Requirements
For the Degree of

DOCTOR OF PHILOSOPHY
WITH A MAJOR IN ZOOLOGY

In the Graduate College
THE UNIVERSITY OF ARIZONA

1 9 7 6

THE UNIVERSITY OF ARIZONA

GRADUATE COLLEGE

I hereby recommend that this dissertation prepared under my
direction by Mary Vaughan Price
entitled The Role of Microhabitat in Structuring Desert
Rodent Communities
be accepted as fulfilling the dissertation requirement for the
degree of Doctor of Philosophy

Edmundell Lockrum
Dissertation Director

4 Oct 1976
Date

As members of the Final Examination Committee, we certify
that we have read this dissertation and agree that it may be
presented for final defense.

Edmundell Lockrum
E A Stahl
Charles Mason
Ron Yellin
Willard Van Arsdale

4 Oct 1976
4 Oct 1976
4 Oct 1976
4 Oct 1976
5 Oct 1976

Final approval and acceptance of this dissertation is contingent
on the candidate's adequate performance and defense thereof at the
final oral examination.

STATEMENT BY AUTHOR

This dissertation has been submitted in partial fulfillment of requirements for an advanced degree at The University of Arizona and is deposited in the University Library to be made available to borrowers under rules of the Library.

Brief quotations from this dissertation are allowable without special permission, provided that accurate acknowledgment of source is made. Requests for permission for extended quotation from or reproduction of this manuscript in whole or in part may be granted by the head of the major department or the Dean of the Graduate College when in his judgment the proposed use of the material is in the interests of scholarship. In all other instances, however, permission must be obtained from the author.

SIGNED: Mary Vaughan Price

ACKNOWLEDGMENTS

I warmly thank those who by word, deed, or example contributed to the completion of this graduate apprenticeship. I am especially grateful to my doctoral committee members--Dr. E. L. Cockrum, Dr. H. R. Pulliam, Dr. E. A. Stull, Dr. W. Van Asdall, and Dr. C. T. Mason, Jr.--for their advice, criticism, and support. Discussions with graduate student colleagues, notably N. Waser, G. Pyke, J. Cross, C. Hoagstrom, E. Roth, and the members of the O.T.S. Tropical Biology 73-1 course, were invaluable, as was the cheerful assistance in field work donated by N. Waser, P. Warren, J. Wondolleck, and the "hole in the wall gang." The manuscript was critically edited and greatly improved by my committee members and by Dr. J. H. Brown.

This work was supported by an N.D.E.A. Title IV Fellowship, an N. S. F. Doctoral Dissertation Improvement Grant, and a Sigma Xi Grant-in-Aid-of-Research.

TABLE OF CONTENTS

	Page
LIST OF TABLES	vi
LIST OF ILLUSTRATIONS	vii
ABSTRACT	viii
 INTRODUCTION	 1
HETEROMYID RODENT COMMUNITIES	3
THE STUDY AREA	6
PRELIMINARY STUDY: MICROHABITAT USE IN A FOUR-SPECIES HETEROMYID RODENT COMMUNITY	9
Methods	9
Results	10
EXPERIMENT ONE: DYNAMICS OF MICROHABITAT USE IN RESPONSE TO CHANGING COMPETITOR DENSITY	20
Methods	21
Results	26
Niche Expansion	26
Niche Contraction and Shift with Increasing Competitor Density	29
Discussion	33
EXPERIMENT TWO: MICROHABITAT AVAILABILITY AS A DETERMINER OF DESERT RODENT ABUNDANCE	35
The Study Area and Experimental Design	36
Phase 1: Pre-manipulation Rodent-Microhabitat Relationships	37
Methods: Microhabitat Censuses	37
Methods: Rodent Censuses	39
Statistical Methods	40
Results: Pre-manipulation Dependence of Rodent Abundance on Microhabitat Availability	41

TABLE OF CONTENTS--Continued

	Page
Phase Two: Rodent Responses to Microhabitat Manipulation	47
Methods	47
Results	48
DISCUSSION	56
APPENDIX A: CAPTURE DATA FROM ENCLOSURE EXPERIMENTS	59
APPENDIX B: SUMMARY OF DATA FROM INTACT COMMUNITY STUDY	63
APPENDIX C: SUMMARY OF RODENT CAPTURE AND VEGETATION CENSUS DATA FROM MANIPULATION STUDY	65
LITERATURE CITED	68

LIST OF TABLES

Table		Page
1.	Microhabitat use overlap in four coexisting heteromyid rodents	17
2.	Experimental design and timetable of enclosure experiments	25
3.	The effect of competitor density on use of four microhabitats by four heteromyid species	28
4.	Principle components of vegetation variables	42
5.	Summary table of stepwise multiple linear regressions of pre-manipulation rodent densities on microhabitat principle component amplitudes	44
6.	Pre-manipulation partial regression of rodent density against significant ($F \geq 1$) principle components of vegetation variables	46
7.	Mean and standard deviation of per-station rodent density at control and manipulated stations before and after manipulation	50
8.	Species composition of manipulated and control sites before and after manipulation	54

LIST OF ILLUSTRATIONS

Figure	Page
1. Body size and morphology of the four heteromyid rodent species studied	7
2. Diagram of sampled microhabitat categories	11
3. Summer microhabitat distributions from the four-species assemblage	13
4. Summer and winter microhabitat distributions of <u>Dipodomys merriami</u> and <u>Perognathus baileyi</u>	14
5. The effect of competitor density on capture frequencies of four heteromyid species in four microhabitat categories .	30
6. The response of <u>Dipodomys merriami</u> to augmentation of large open spaces	51

ABSTRACT

Competition is thought to play a central role in determining patterns of resource use and species abundances in natural communities. Nevertheless, as yet we do not know the extent to which competition for scarce resources can determine the structure of multi-species assemblages. Enclosure experiments were used to see if resource use by seed-eating desert rodents changed in response to the presence of competitors. In addition, perturbation experiments assessed the effect of resource availability on species abundances. Results suggest that competition maintains interspecific differences in the microhabitats from which seeds are gathered, and that the availability of appropriate microhabitats determines patterns of species abundances on a local scale.

INTRODUCTION

What are the forces determining the structure of multi-species assemblages? This is a question central to community ecology, and one whose answer could, in the least tractable of all possible worlds, be different for each system examined. The hope of ecologists is that a relatively few factors can be built successfully into a predictive general theory of the number and relative abundance of species.

Much recent theoretical treatment of community structure (Levins 1968, MacArthur 1972, May 1973) has followed the pioneering work of Volterra (1928), Lotka (1925), and Gause (1934) in postulating interspecific competition for scarce resources as a force that commonly limits natural populations. This assumption yields a series of predictions about the structure of assemblages of resource competitors. Because it is difficult to identify the resource(s) for which species compete and the manner in which they interact, competition theory--and the community structure predictions to which it leads--has only recently begun to be examined in a careful and quantitative way (e.g., Yeaton and Cody 1974, Pulliam 1975, Werner and Hall 1976, Titman 1976). Further tests of competition theory are clearly necessary before we can assess the extent to which competitive interactions explain patterns of species diversity.

The research presented here was designed to see whether resource use by seed-eating desert rodents fits patterns predicted by

competition theory (Preliminary Study and Experiment 1) and whether variations in species abundances might be accounted for by spatial variations in resource availability (Experiment 2). Results support the notion that competition between rodents for foraging micro-habitat in large part maintains patterns of resource use and species abundances in desert rodent communities.

HETEROMYID RODENT COMMUNITIES

The Heteromyidae comprises a New World family of kangaroo rats, pocket and kangaroo mice that reaches high diversity in arid regions of North America and Mexico, with five or more species often co-occurring at one locality. Heteromyids live in burrows during the day and emerge at night to forage, storing individual food items in external cheek pouches presumably until enough volume has been gleaned to be worth transport to a cache. Many species rely almost exclusively on the seeds of desert annuals, which are produced sporadically in response to rains. Seeds undoubtedly limit desert rodent populations, because on a large geographic scale rodent densities are correlated with annual rainfall and vegetation productivity (Brown 1975) and because local densities increase dramatically following years of high seed production (Pearson 1975, Whitford 1976, Cockrum 1975). Competition theory would predict that subdivision of this limiting resource is critical to prolonged coexistence of heteromyid species. However, it seems that direct subdivision of food by size or species is either rare or of minor importance to coexisting heteromyids, perhaps because seed densities fluctuate too unpredictably to allow prolonged specialization. Riechman (1975), Brown and Lieberman (1973), and Smigel and Rosenzweig (1974) found very high overlap in the seed types gathered and eaten by co-occurring heteromyids. Although one species, Dipodomys microps, has

become a specialized herbivore (Kenagy 1972), this is an exceptional case, and most heteromyids are granivorous.

Despite the lack of obvious seed partitioning, there remains considerable evidence that heteromyid assemblages are competitively structured in North American deserts (Rosenzweig and Winakur 1969, Brown and Lieberman 1973, Rosenzweig 1973, Brown 1975, Wondolleck 1975). Brown (1973, 1975) noted regular spacing of body size among coexisting species and inferred that coexistence is allowed by subdivision of a body size related resource. Rosenzweig and Winakur (1969) correlated rodent species diversity with "complexity" of soil and vegetation in Arizona study sites. From this they postulated that the rodents subdivide a habitat resource, and that increased habitat complexity promotes rodent diversity by providing many habitat "niches." Hoagstrom (1976) also noted species-specific habitat affinities for five heteromyids in southern Arizona.

Rosenzweig (1973) demonstrated that species-specific habitat preferences occur by noting a differential change in the abundances of two species in response to augmentation of two habitat types. Wondolleck (1975) and Brown and Lieberman (1973) also documented microhabitat differences between coexisting heteromyids. Interspecific differences in the place from which seeds are gathered would result in effective partitioning of the soil's seed reserves, and a very plausible hypothesis emerging from these early studies is that coexistence of heteromyid rodent species is mediated at least partially through allocation of horizontal foraging microhabitats. Indeed, Lemen (1976; see also

Rosenzweig, in press) found that two coexisting species collect seeds from distinct microhabitats.

Non-heteromyid rodents also compete for space (Grant 1972, Douglass 1976), and habitat selection appears to be a major mechanism allowing regional coexistence of cricetid species with similar nutritional requirements (Grant 1972, M'Closkey and Fieldwick 1975). It would not be surprising to find that heteromyids similarly subdivide a space resource, although the scale on which the discrimination occurs (if it exists) must be extremely fine, for the large-scale and often clear-cut habitat segregation observed among coexisting cricetids is not as pronounced among desert-dwelling mice.

The research reported here was designed to study microhabitat use by four heteromyid species that coexist 30 miles south of Tucson, Arizona, in hopes of establishing clearly that interspecific differences in microhabitat use are being maintained (and were thus probably initiated) by interspecific competitive interactions; and that microhabitat availability in part determines the abundances of rodent species. The experiments constituted a qualitative test of current theoretical propositions (MacArthur 1972, May 1973) that competition and resource availability can determine the structure of natural communities.

THE STUDY AREA

The study area was situated in Lower Sonoran Desert--Desert Grassland transition (Lowe 1964) at 950 meters elevation. All experiments were conducted 30 miles south of Tucson, Arizona, on ungrazed portions of the Santa Rita Experimental Range, United States Department of Agriculture. Vegetation of the study area is complex and patchy, containing scattered trees (Prosopis juliflora and two Cercidium species), large shrubs (Celtis pallida, Acacia greggii, Ephedra trifurca), small shrubs (Haplopappus tenuisectus, Baccharis brachyphylla, Zinnia pumila), grass clumps (e.g., Aristida species, Andropogon barbinodis, Tricachne californica, Stipa neomexicana), and periodically abundant low annual plants. Surface soil also is heterogeneous in this region, as washes dissect the bajada, and pebbly wash banks alternate with fine sandy soil on small plateaus.

Four heteromyid rodent species are abundant in the area and are so distributed that all of the four can be caught at a single trap position. They include a kangaroo rat (Dipodomys merriami) and three pocket mice (Perognathus amplus, P. penicillatus, and P. baileyi), which conform qualitatively to the body size spacing pattern discussed by Brown (1973, 1975). See Figure 1. Several cricetid rodents (Neotoma albigula, three Peromyscus species, Onychomys torridus, and three Reithrodontomys species) and one sciurid rodent (Ammospermophilus

Dipodomys merriami

HEAD AND BODY LENGTH = 95 mm
 TAIL LENGTH = 130 mm
 HIND FOOT LENGTH = 36.1 mm
 BODY WEIGHT = 33 g

Perognathus baileyi

HEAD AND BODY LENGTH = 92 mm
 TAIL LENGTH = 109 mm
 HIND FOOT LENGTH = 26.3 mm
 BODY WEIGHT = 27 g

Perognathus penicillatus

HEAD AND BODY LENGTH = 76 mm
 TAIL LENGTH = 94 mm
 HIND FOOT LENGTH = 22.6 mm
 BODY WEIGHT = 17 g

Perognathus amplus

HEAD AND BODY LENGTH = 69 mm
 TAIL LENGTH = 73 mm
 HIND FOOT LENGTH = 19.3 mm
 BODY WEIGHT = 12 g

10 CM

Figure 1. Body size and morphology of the four heteromyid rodent species studied. -- Body measurements are the means of ten specimens taken from the Santa Rita Experimental Range.

harrisii) occur in the area but altogether comprised less than ten percent of all individuals caught. The most numerous cricetid species, Onychomys torridus, is locally abundant, but being primarily insectivorous was excluded from study because its diet is very different from that of the granivorous heteromyids.

PRELIMINARY STUDY: MICROHABITAT USE IN A FOUR-SPECIES HETEROMYID RODENT COMMUNITY

Because of the proposed importance of habitat selection to heteromyid community structure (Brown and Lieberman 1973, Rosenzweig 1973, Rosenzweig and Winakur 1973, Wondolleck 1975), habitat utilization patterns of four heteromyid rodents were quantified at the Santa Rita study site as baseline data for the experimental studies to follow. Interspecific differences in microhabitat use would suggest that divergence in use of this resource has been critical to the coexistence of these heteromyids.

Methods

Six qualitatively distinct microhabitats were sampled at each of 33 stations spaced at approximate 15-meter intervals on an irregular grid. This was accomplished by placing one 25 x 10 x 10 cm rectangular metal live trap in each of six microhabitats within a circular area of five meter radius and then moving at least ten meters away to establish, in a similar manner, the next six-trap station. Microhabitat categories were chosen to sample the full range of structural vegetation features that might be distinguished by a mouse while foraging for seeds on the ground. They were defined as follows: "large open" (a space of bare ground at least two meters in diameter), "small open" (a space of bare ground less than 0.50 meter and greater than 0.25 meter in diameter),

"small bush" (under a shrub between 0.6 and 1.0 meter in height and one meter away from another shrub), "clump" (in a grass clump or patch of low shrubs with continuous canopy of at least one meter diameter), "large bush" (under a shrub at least one meter in height and two meters in diameter), and "tree" (under a Prosopis or Cercidium individual at least 1.7 meters tall). All traps were placed carefully to sample as consistently as possible the defined microhabitat categories (Figure 2).

Rodent microhabitat use was sampled for 12 nights during the period 6 June to 30 September 1974, and for six nights during the following November, December, and January. Traps were baited with rolled oats at dusk, checked for captured animals at 10:00 p.m. and 6:00 a.m. the following morning, and then closed for the day. During the winter months traps were checked and closed at 10:00 p.m. to avoid torpor deaths. Animals were toe-clipped for individual recognition and each capture event was coded for species, individual, station, and microhabitat. The intensity with which each species used each microhabitat was measured as the relative number of captures, and of individuals, recorded in each category during a season's trapping.

Results

Proportional use of microhabitats was calculated for each species from the distribution of captures and of individuals. Chi-square homogeneity tests show that there was no heterogeneity between microhabitat distributions resulting from the two measures ($\chi^2 < 2$ and $p > .5$ for all species). Because "individuals" data yield small sample sizes, only

Figure 2. Diagram of sampled microhabitat categories.

microhabitat preferences reflected in capture frequencies are considered in the remaining analyses (Figures 3 and 4).

During the summer season, no species was equally active in the six microhabitat categories (chi-square goodness-of-fit tests of observed vs. random expected capture distributions yielded $\chi^2 > 15$ and $p < .025$ in all four cases). All seemed to have one or two preferred microhabitats (Figure 3). Dipodomys merriami was caught most frequently in large open traps and least often under trees. Perognathus amplus also used open spaces, but small open and small bush traps were favored over those placed in large open patches. Perognathus penicillatus and P. baileyi were most active in large bush and tree categories.

Not only were the mice selectively active in certain microhabitats, but each species was most active in a unique category. The four summer microhabitat distributions (Figure 3) were significantly heterogeneous at the .005 level (homogeneity $\chi^2 = 141.86$); and pairwise comparisons of the species most similar in microhabitat use (D. merriami and P. amplus; P. penicillatus and P. baileyi) show that even in these cases the preferred category of one species was used significantly less by the other. A one-tailed t-test of difference between proportions (Sokal and Rohlf 1969, p. 607) suggests D. merriami used large open spaces more than did P. amplus ($t = 1.87$; $p = .03$) and P. penicillatus used large bushes more than P. baileyi ($t = 1.57$; $p = .06$).

Microhabitat use changed seasonally (Figure 4). Winter capture distributions of both D. merriami and P. baileyi (the only winter-active species) were significantly different ($\chi^2 > 15$; $p < .01$ in both cases)

Figure 3. Summer microhabitat distributions from the four-species assemblage. -- Relative frequencies-of-capture in six microhabitats are given for four heteromyid species. Numbers above bars are the total number of captures obtained in each habitat category during the June-September 1974 sampling period. LO = large open; SO = small open; SB = small bush; C = clump; LB = large bush; T = tree.

Figure 4. Summer and winter microhabitat distributions of *Dipodomys merriami* and *Perognathus baileyi*. -- Relative frequencies-of-capture of two species in six microhabitats are compared for the periods June-September 1974 (stippled bars) and November 1974-January 1975 (white bars). Microhabitat abbreviations and numbers above bars are as in Figure 3.

from summer utilizations. Despite shifts in activity, both species maintained, as in the summer, separate most-used microhabitats: D. merriami was primarily active in small open spaces while P. baileyi confined its activity to large bushes. The direction of the shift is interesting, for both species increased, in winter, use of a microhabitat that had been usurped previously by another species: P. amplus had used small open and P. penicillatus large bushes. Such a change could result from seasonal shifts in food dispersion, or from the thermal advantages accruing to an animal protected in winter from radiant energy loss to a cold sky. It also could be due to reduced interspecific competition from hibernating species.

To summarize, each species maintained utilization patterns distinct from those of other simultaneously-active rodents, even though microhabitat specificity changed seasonally. These and previous data (Rosenzweig and Winakur 1969, Brown and Lieberman 1973, Rosenzweig 1973, Wondolleck 1975) confirm that heteromyid rodents partition structural space, and that habitat discrimination can be remarkably fine-tuned. Nevertheless, it is not immediately obvious how and why microhabitat is discriminated by these rodents, nor whether the degree of interspecific difference observed is sufficient by itself to allow their coexistence.

If indeed joint use of foraging microhabitats is the major mechanism of competition, could observed interspecific differences theoretically allow heteromyid coexistence? MacArthur (1972) estimated, for three species whose resource utilizations differ only in their position along a one-dimensional resource continuum, that the

maximal overlap consistent with coexistence is approximately .54 when overlap is measured as

$$\alpha_{ij} = \frac{\sum_k p_{ik} p_{jk}}{\sum_k (p_{ik})^2}$$

where p_{ik} is the proportional use of resource state k by species i .

Overlap values can be calculated according to this formula from the microhabitat distributions given in Figure 3 and compared to this theoretical limit. However, overlaps calculated from a raw resource use matrix like the one derived from Figure 3 might be biased by unequal resource state (microhabitat) distinctiveness, especially if resource categories were arbitrarily chosen as in this study. It is very possible, for instance, that heteromyids might not distinguish small bushes from small open spaces. To correct for such potential biases, Colwell and Futuyma (1971) have derived a resource state weighting factor based on the relative contribution of each resource to total heterogeneity of species distributions with respect to resource states. A resource use matrix in which each resource state is multiplied by the relative weighting factor yields niche overlap values that are corrected for variation in resource distinctiveness. Overlap values in Table 1 were calculated from such an expanded microhabitat use matrix. Entries in the table can be interpreted as the amount of resource usurped by one species (j) at the expense of another (i), relative to species i 's effect on itself.

Table 1. Microhabitat use overlap in four coexisting heteromyid rodents.

Species i	Species j			
	<u>Dipodomys merriami</u>	<u>Perognathus amplus</u>	<u>Perognathus penicillatus</u>	<u>Perognathus baileyi</u>
<u>Dipodomys merriami</u>	1.00	.936	.504	.563
<u>Perognathus amplus</u>	.998	1.00	.550	.606
<u>Perognathus penicillatus</u>	.622	.637	1.00	.937
<u>Perognathus baileyi</u>	.763	.769	1.03	1.00

Most observed overlap values exceed MacArthur's .54 estimate of limiting similarity, some by a considerable amount. This finding could mean that coexistence of heteromyids is not simply a function of their divergence in microhabitat use, or that the resource continuum is neither linear nor one-dimensional. However, these values might be underestimates of real interspecific differences for the following reasons: (1) Trapping methods do not measure foraging microhabitat precisely because a mouse can be captured while traveling through an inappropriate habitat patch to reach an appropriate one; actual foraging microhabitat specificities are thus probably higher than these data suggest. (2) The range of categories defined for this study is unlikely to coincide with all those discriminated by the rodents. For example, D. merriami probably utilizes open spaces larger than those sampled, and therefore its high overlap with P. amplus (.998 and .936) may be an artifact of sampling only one portion of the microhabitat states actually used by D. merriami. In addition, surface soil texture, one aspect of microhabitat discriminated by the rodents, was not sampled quantitatively. Between-station variation in P. baileyi density was correlated most highly with the abundance of large bushes ($r = .56$ in summer and $.43$ in winter), and secondarily with pebbly surface soil ($r = .30$ in summer and $.32$ in winter). See Appendix A. High overlap values between P. baileyi and the other three species may be due largely to this unsampled aspect of microhabitat.

Another possibility is that the microhabitat categories studied are not being subdivided actively by the rodents, but instead are

imperfectly correlated with the real resource being partitioned. Since the degree of activity measured by trapping in a microhabitat is directly related to the amount of foraging effort a heteromyid expends in that microhabitat (Price, in press), space partitioning may represent divergent specialization for foraging on particular densities and dispersions of seed resources. Seeds are not randomly scattered in desert soils (Goodall and Morgan 1974). On the contrary, their dispersal by wind and by water result in patterns of density and clumping that coincide with local variations in soil surface microtopography and structural features of vegetation (Reichman 1976). It may well be that patterns in morphological divergence between coexisting heteromyids (Brown 1973, 1975) parallel observed divergence in microhabitat use; the energetics of foraging may dictate that one particular body size and morphology is maximally efficient for a given seed array.

EXPERIMENT ONE: DYNAMICS OF MICROHABITAT USE IN RESPONSE TO CHANGING COMPETITOR DENSITY

Resource partitioning is often taken as evidence that competition in the past has caused evolutionary divergence between species in the use of limiting resources, and that competition presently maintains the niche separations. Certainly such phenomena are predicted by and therefore are consistent with competition theory, but alternate explanations could account for these patterns. By themselves, patterns do not identify mechanisms of their production and maintenance.

It is reasonable to expect that no species will exploit only one small portion of available food or habitat resources unless some force prevents niche expansion. This follows from predictions of optimal diet and optimal habitat use theories (MacArthur and Pianka 1966, Schoener 1971, Pulliam 1974) that an animal should take a more diverse array of food items or forage in a diverse array of patch types when the density of the best food type is low or the amount of food in the best patch is depleted. Since conspecifics prefer the same resources, an animal faced with intraspecific competitive pressures will tend to diversify its resource use when preferred resource types become scarce. Interspecific competition from individuals with similar resource utilizations could counteract the tendency of one species to expand its resource use by maintaining neighboring resources at a density too low to promote their use by the first species. In this way, competition could

account for the maintenance of divergent specializations among similar species. If this is indeed the mechanism which maintains the niche structure of communities, then resource use by one species should expand when competitors are absent and shift and/or contract when competitors are present (Vandermeer 1972).

Coexisting heteromyid rodents in North American deserts differ in their use of foraging microhabitats (Brown and Lieberman 1973, Wondolleck 1975) as well as in body size and morphology (Brown 1973, 1975). These observations have lead to the inference that communities of seed-eating desert rodents are competitively structured. Nevertheless, there is little direct evidence that competition occurs between co-existing heteromyids and indeed, there is some to the contrary (Schroeder and Rosenzweig 1975). Supporting evidence could be derived from a study of the dynamics of microhabitat use by one species as a function of the density of another. With this in mind, microhabitat use by four species of coexisting heteromyid rodents was measured when individuals of the other species were present and absent. Support for the hypothesis that microhabitat allocation reflects interspecific competition would consist of changes in the range, evenness, and types of resources used by one species in directions predicted theoretically for competitive situations.

Methods

A 1.84 hectare rectangular rodent enclosure built on the Santa Rita Range by Charles Lowe was partitioned into six contiguous enclosures ranging in area from .24 to .36 hectares. Partition walls consisted of

aluminum window screening 1.4 meters wide that was buried .5 meters in the soil and supported every 1.5 meters by aluminum poles. Outside enclosure walls were of the same height and buried similarly, but consisted of sheet metal supported by wooden stakes. The enclosures were situated adjacent to the study site where microhabitat partitioning among the four resident heteromyid rodent species was documented (see Preliminary Study page 9 of this dissertation).

All rodents resident in the enclosures were removed by snap-trapping prior to the start of the experiments. Adult animals were collected by live trapping within a mile of the study area and housed in cages at The University of Arizona under a 12-hour light-dark regime pending release in the enclosures. Individuals were maintained in the laboratory for varying lengths of time, but at most for two consecutive months, and most rodents were kept less than one month.

Microhabitat use by each of the four heteromyid species was sampled and tallied in a manner similar to that described for the Preliminary Study. In this case, though, four instead of six microhabitat categories were considered in order to maximize the sampling effort in each microhabitat type with a limited number of traps. To decide which of the six categories should be eliminated, Colwell and Futuyma's (1971) method for deriving weighting factors for a series of resource states was used with the rodent capture data from Figure 3. These weighting factors reflect the relative contribution of each microhabitat category to total between-habitat heterogeneity of rodent captures. Habitats were found to have relative weights as follows: large open, .105; small

open, .158; small bush, .141; clump, .126; large bush, .207; tree, .362. Aside from large open, clump was the least distinctive of the categories to the rodents, and so it was eliminated. Although large open had the lowest weighting factor, it represented the preferred microhabitat for at least one species--D. merriami--and I was loathe to exclude it. Small bush was therefore eliminated completely, for no species preferred it and its weight was small.

Eight traps were placed in each microhabitat category in each enclosure. Trap locations were rotated every four trap nights and individuals were released at least ten paces from point-of-capture to reduce recapture bias from individuals cuing on one trap site. Traps were baited with a pinch of rolled oats at dusk and checked twice nightly (except during April and May 1975, when they were closed at 10:00 p.m.). Microhabitat category, animal number (individuals were toe-clipped), and trap location were recorded for each capture event.

Each experimental treatment was continued until at least 35 legitimate (i.e., excluding within-night recaptures of an individual at a trap or captures of escapees from other enclosures) captures had been recorded in each enclosure. Densities were maintained approximately constant (they varied by $\pm$ two individuals per enclosure) by releasing new marked animals to replace known mortalities or individuals that had not been captured for four trapping sessions. Occasionally animals would escape from one enclosure to another. These were reestablished in the proper enclosure, or removed and replaced by new animals if they proved to be habitual fence-climbers. Unmarked animals appearing in

the enclosures (these were rare) were removed. After being added to enclosures at the start of a treatment, animals were allowed three to four days to equilibrate in the new environment before sampling resumed.

Microhabitat use (measured as capture frequency excluding within-night recaptures of an individual in one trap) was determined for each species first at a density of eight individuals per enclosure when no other species was present, secondly when four individuals of another species were added, and finally when the second species was present at high density (eight individuals). Dipodomys merriami, the largest of the four species, was chosen as the "competitor" to be placed with each of the three Perognathus species as it was likely that this kangaroo rat was behaviorally dominant over each of the smaller pocket mice (MacMillan 1964) and would elicit the most noticeable response. Table 2 summarizes details of experimental design.

Enclosure densities were chosen so as to be consistent with normal field densities for the area. In the summer of 1975, there were approximately 67 heteromyids per hectare on the Santa Rita Experimental Range (Courtney 1975), which would correspond to about 20 mice per enclosure. Highest enclosure densities used were 16 individuals per enclosure. Since the mice remained in excellent condition, and even gained weight during the experiments, these were reasonable densities.

Niche breadths for each species and each treatment were calculated by the formula:

$$\text{Niche Breadth} = -\sum_{i=1}^4 p_i \log_{10} p_i$$

Table 2. Experimental design and timetable of enclosure experiments. -- The species present at each time are given for each enclosure.

Enclosure Number	Treatment 1		Treatment 2		Treatment 3	
	April 11, 1975	May 22, 1975	June 21, 1975	July 3, 1975	July 20, 1975	
1W	8 <u>Dipodomys merriami</u>	8 <u>Perognathus amplus</u>	8 <u>P. amplus</u> + 4 <u>D. merriami</u>	8 <u>P. amplus</u> + 8 <u>D. merriami</u>		
2W	8 <u>D. merriami</u>	8 <u>P. amplus</u>	8 <u>P. amplus</u> (control)	8 <u>P. amplus</u>		
2E	8 <u>D. merriami</u>	8 <u>P. penicillatus</u>	8 <u>P. penicillatus</u> + 4 <u>D. merriami</u>	8 <u>P. penicillatus</u> + 8 <u>D. merriami</u>		
3W	8 <u>D. merriami</u>	8 <u>P. penicillatus</u>	8 <u>P. penicillatus</u> (control)	8 <u>P. penicillatus</u>		
1E	8 <u>D. merriami</u>	8 <u>P. baileyi</u>	8 <u>P. baileyi</u> + 4 <u>D. merriami</u>	8 <u>P. baileyi</u> + 8 <u>D. merriami</u>		
3E	8 <u>D. merriami</u>	8 <u>P. baileyi</u>	8 <u>P. baileyi</u> (control)	8 <u>P. baileyi</u>		

where p_i is the proportion of all captures occurring in microhabitat category i (Levins 1968). The maximum breadth for four habitat categories is .602, and would be obtained if a species were caught with equal frequency in all categories.

Results

Results were analyzed with regard to the following predictions from niche theory: (1) In the absence of competitors (Treatment 1), the breadth of microhabitat use should be higher than 1974 intact community values. In particular, each species is expected to increase use of microhabitats that it used only marginally in 1974 and decrease use of those it utilized highly in the intact community. (2) As the density of a single competitor species increases (Treatments 1 through 3), microhabitat niche breadth should decrease, and use of the competitor's preferred microhabitats should also decrease. Specifically, D. merriami's use of large open (its preferred category) should increase and its use of trees (the least preferred microhabitat) decrease in the presence of Perognathus. In turn, Perognathus should shift out of large open and into tree habitats when D. merriami are present (see Appendix B for raw data).

Niche Expansion

We can determine the effect of competitive release by comparing microhabitat use of a species in the four-species community studied in 1974 (see Preliminary Study) with the same species' behavior when alone in an enclosure. Sampling methods were equivalent in both studies, and

the density of heteromyids was, if different at all, higher in 1974. This would have the effect of biasing 1974 niche breadths in an upward direction.

For all species except D. merriami, microhabitat use when alone was significantly different from the intact community utilization (homogeneity $\chi^2 > 10$, $p < .025$ for each of the three Perognathus species). This change represented a niche expansion, for niche breadths of all species increased in Treatment 1 over the four-species intact community (Table 3). Although there is no good way of assessing the significance of a single change in this niche breadth index, a probability value for four changes in the same direction can be derived. If the niche breadth of one species changed randomly between treatments, we would expect an increase in niche breadth to be as likely as a decrease. Thus the probability of having all four species increase their niche breadths between the intact community and Treatment 1 by chance is only $(.5)^4 = .06$.

Shifts in microhabitat utilization patterns accompanying the niche expansion were as expected. Each species increased use of all microhabitats originally avoided (i.e., used with less than 25 percent frequency) and decreased use of microhabitats that were preferred (i.e., those having more than 25 percent of total captures) in the four-species community. Sixteen out of 16 possible shifts were in the proper direction.

Despite decreases in intensity of microhabitat preference due to interspecific competitive release, all species except P. penicillatus maintained distributions significantly different from random ($p < .05$

Table 3. The effect of competitor density on use of four microhabitats by four heteromyid species.

Species	Large Open	Small Open	Bush	Tree	Niche Breadth $\sum p_i \log_{10} p_i$
<u>Dipodomys merriami</u>					
Intact community	21 (.488)	13 (.302)	6 (.140)	3 (.070)	.510
Treatment 1	125 (.374)	92 (.275)	74 (.222)	43 (.129)	.574
Treatment 2 (actual)	24 (.308)	21 (.269)	23 (.295)	10 (.128)	.581
(expected)	29 (.369)	22 (.279)	17 (.219)	10 (.133)	.577
Treatment 3 (actual)	62 (.492)	29 (.230)	31 (.246)	4 (.032)	.497
(expected)	50 (.393)	34 (.264)	28 (.220)	15 (.121)	.568
<u>Perognathus amplus</u>					
Intact community	74 (.351)	83 (.393)	39 (.185)	15 (.071)	.537
Treatment 1	72 (.327)	61 (.277)	55 (.250)	32 (.145)	.585
Treatment 2 (actual)	15 (.254)	14 (.237)	12 (.203)	18 (.305)	.596
(expected)	19 (.320)	17 (.284)	15 (.247)	9 (.150)	.609
Treatment 3 (actual)	5 (.143)	11 (.314)	11 (.314)	8 (.228)	.583
(expected)	9 (.262)	11 (.299)	9 (.262)	6 (.179)	.595
<u>Perognathus penicillatus</u>					
Intact community	18 (.145)	12 (.097)	51 (.411)	43 (.347)	.538
Treatment 1	67 (.296)	40 (.177)	61 (.270)	58 (.257)	.595
Treatment 2 (actual)	25 (.410)	8 (.131)	16 (.262)	12 (.197)	.566
(expected)	18 (.295)	10 (.168)	17 (.271)	17 (.266)	.593
Treatment 3 (actual)	6 (.182)	10 (.303)	9 (.273)	8 (.242)	.595
(expected)	8 (.233)	6 (.176)	10 (.288)	10 (.300)	.593
<u>Perognathus baileyi</u>					
Intact community	22 (.195)	20 (.177)	37 (.327)	34 (.301)	.587
Treatment 1	58 (.195)	87 (.293)	89 (.300)	63 (.212)	.594
Treatment 2 (actual)	16 (.213)	21 (.280)	22 (.293)	16 (.213)	.597
(expected)	14 (.189)	23 (.297)	23 (.300)	17 (.221)	.595
Treatment 3 (actual)	3 (.081)	10 (.270)	5 (.135)	19 (.514)	.507
(expected)	6 (.173)	11 (.286)	11 (.295)	9 (.246)	.593

in three cases). Furthermore, the microhabitat distributions of the four species remained significantly heterogeneous (chi-square homogeneity test comparing Treatment 1 capture frequencies between species yields $\chi^2 = 28$, $p < .005$). This implies microhabitat preferences are not solely functions of competitive interactions, but are maintained to a lesser extent even in the absence of other species. This could be due to morphological specializations for efficient foraging in certain microhabitats, or could be an artifact of the short time scale of the experiments.

Niche Contraction and Shift with Increasing Competitor Density

By comparing changes in microhabitat use and niche breadth between Treatments 1, 2, and 3, we can see whether increased competitor density resulted in niche contraction (Table 3 and Figure 5).

The niche breadths of D. merriami, P. amplus, and P. baileyi increased between Treatments 1 and 2, and then decreased in Treatment 3. That of P. penicillatus decreased at first and then increased to the level of Treatment 2. Niche theory clearly predicts that niche breadths should decrease with increasing competitor density, but half of the niche breadth changes between Treatments 1 through 3 were in the "right" and half in the "wrong" direction.

Nevertheless, no species had a Treatment 3 niche breadth larger than its Treatment 1 value, and the changes of greatest magnitude (D. merriami and P. baileyi between Treatments 2 and 3) were in the "right" direction. It may be that the niche breadth index is

Figure 5. The effect of competitor density on capture frequencies of four heteromyid species in four microhabitat categories. -- Numbers above histogram bars are numbers of captures.

insensitive to shifts in microhabitat use, for when decreased use of one category is compensated by increased use of another, the "unevenness" of resource use remains the same.

Shifts in use of particular microhabitat categories were as predicted. Dipodomys merriami did not respond differentially to the three pocket mice (homogeneity chi-square between enclosures; $\chi^2 = 5.62$ for Treatment 2 and 4.34 for Treatment 3, $p > .1$ in both cases). However, the kangaroo rats changed their microhabitat use as a whole between treatments (lumping data from all enclosures within a treatment, the between-treatment $\chi^2 = 15.6$, $p = .017$). The change was due largely to their being captured more frequently in large open traps and less frequently in trees during Treatment 3.

No Perognathus showed significant over-all heterogeneity between the three treatments. However, P. amplus responded significantly to the low kangaroo rat treatment, decreasing use of large open and increasing use of other habitat types (lumped controls vs. Treatment 2 data were homogeneous, $\chi^2 < 3.21$ and $p > .1$ for two species; and heterogeneous for P. amplus, $\chi^2 = 16.9$, $p < .005$). At high D. merriami density, both P. baileyi and P. amplus significantly decreased their use of large open spaces from Treatment 1 levels while increasing use of other habitats (notably, trees) not preferred by the kangaroo rats (one-tailed t-test of the difference of large open capture proportions; P. amplus $t = 2.43$, $p = .008$; P. baileyi $t = 1.94$, $p = .026$). Perognathus penicillatus showed a statistically nonsignificant change in the direction of decreased use of large open ($t = 1.45$, $p = .074$).

Since in Treatment 2 D. merriami density was half the Treatment 1 density, possibly confounding its response to interspecific competition with changes due to decreased intraspecific competition, we look to changes between Treatment 1 and Treatment 3 in use of the large open and tree microhabitats to test niche theory predictions about the directionality of niche shifts. Expected changes are for D. merriami to increase large open use and decrease tree use in Treatment 3, while the three pocket mice should shift in the opposite direction. Seven out of eight changes in these habitats were in the predicted direction.

These effects cannot be artifacts of using traps to sample microhabitat use. It could be argued that observed changes in capture frequency could be due solely to a trap saturation effect. If one species (D. merriami, for example) preferentially entered traps in large open spaces before Perognathus could find them, they would decrease the availability to pocket mice of traps in large open spaces relative to other microhabitat types. This would result in decreased pocket mouse captures in large open spaces even if their use of microhabitat patches remained unchanged by the presence of D. merriami. In Table 3, this effect was considered in calculating "expected" capture frequencies for Treatments 2 and 3. To derive expected values, the Treatment 1 capture frequencies were multiplied by the relative number of traps left vacant by the competitor during a treatment and then normalizing these four joint probabilities. The actual capture distributions for Treatments 2 and 3 together were significantly different from expectations for D. merriami ($\chi^2 = 7.8$, $p = .051$) and P. amplus ($\chi^2 = 10.4$, $p = .017$)

but insignificant for P. penicillatus ($x^2 = 2.7$, $p > .1$). Perognathus baileyi's distribution differed significantly from expectation in Treatment 3 alone ($x^2 = 15.8$, $p < .005$). All Perognathus used large open less than expected; P. amplius and P. baileyi also used trees more than expected while D. merriami used large open more and trees less than expected. Similarly, expected niche breadth changes due only to changing trap availability are not as large as observed changes (Table 3). Thus, observed changes in capture frequencies reflect changing microhabitat use and not trap saturation phenomena.

In summary, the Perognathus shifted microhabitat use in response to the presence of D. merriami, and the shift tended to be out of Dipodomys-preferred habitat into others. Dipodomys merriami increased use of its preferred foraging space in response to the presence of pocket mice. These shifts were accompanied by variable responses of niche breadth indices to increasing competitor density, but no species increased its niche breadth at the highest competitor density relative to that in the noncompetitive situation.

Discussion

These observed changes in microhabitat use are generally consistent with predictions of niche theory, for when competitors are absent heteromyids are less selective in the microhabitats they frequent; and when competitors are present, the rodents become more selective with respect to foraging space and avoid each other's preferred microhabitats. This supports the hypothesis that foraging microhabitat is an object of

interspecific competition and that niche differentiation among species is maintained in part by competition. Were microhabitat preferences completely fixed with no present competition occurring between species, there should have been no consistent directional changes in resource use.

Three out of four species maintained significant microhabitat preferences even in the absence of interspecific competitors. This might indicate that to some extent competitive interactions acting over time have promoted behavioral and/or morphological specializations allowing efficient exploitation of seed resources in certain microhabitats but not others. Divergence in body size and other morphological characters is apparent among these and other coexisting heteromyids (Brown 1973, 1975), but the relationship between microhabitat preference and morphology is obscure. Studies elucidating energetic consequences of body size and morphology to a rodent foraging among different microhabitats (e.g., Rosenzweig and Sterner 1970; Taylor, Schmidt-Nielsen and Raab 1970) will be prerequisites to a complete understanding of the role competition plays in determining resource utilizations and thus its role in structuring heteromyid rodent communities.

EXPERIMENT TWO: MICROHABITAT AVAILABILITY AS A DETERMINER OF DESERT RODENT ABUNDANCE

It is becoming clear that competition can set an upper limit to the number of species which live together by limiting the similarity in their use of resources (see MacArthur 1972 and May 1973 for discussion of the limiting similarity concept, and Abrams 1975 for a critique of their work). However, species number is but one aspect of the structure of communities. It is important also to understand what determines the abundances of coexisting species.

Naturalists (e.g., Grinnell and Orr 1934) have long suspected that the availability of appropriate habitats and foods exerts control over the distribution and abundance of species. This notion was expressed in a clear and quantitative form by MacArthur (1970, 1972), who showed theoretically that competition can result in equilibrium assemblages of competitors whose combined resource utilizations provide the best least-squares fit to available resources. It follows that a proximate cause of particular species abundance relationships among coexisting competitors may be the shape of the spectrum of available resources.

Because the machinery of competition is elusive in most natural situations, this theory of species abundance has been applied with qualified success only in a few instances (Yeaton and Cody 1974, Cody 1974, Pulliam 1975), and further tests of the theory are desirable.

Seed-eating desert rodents may eventually prove amenable to quantitative tests of community theory, since foraging microhabitat has been identified as an important resource for which coexisting heteromyid species compete (Wondolleck 1975). These interactions between species result in microhabitat partitioning such that each species gathers its food preferentially in a microhabitat type distinct from those used by the others. Although a quantitative test of the theory with this system would be premature at present because the basis of microhabitat preferences is only crudely understood, a qualitative test would consist of finding that abundance of heteromyid rodent species is influenced by availability of appropriate microhabitat patches in the direction predicted by theory.

As the first step in establishing a relationship between microhabitat availability and density of rodent species, I determined whether variations in abundance of each species correlated with the abundance of its preferred microhabitat. Secondly I manipulated the availability of selected microhabitat types to test further whether microhabitat exerts control over rodent abundances.

The Study Area and Experimental Design

A large 7 by 7, 49-station grid with 30 meter intervals between stations was set up in July 1975 to study rodent-microhabitat relationships near sites where the microhabitat preferences of four coexisting heteromyid rodents had been previously studied. Vegetation and topographic heterogeneity of the Santa Rita Experimental Range was such that

the grid covered a mosaic of several intermixed habitat patches, and rodent abundances varied considerably between stations.

The study consisted of two phases. First, coverage of several microhabitats was measured and simultaneously rodents were censused at each of the 49 stations. Rodent densities were regressed on the principal components of microhabitat variables to see whether the abundance of each species' preferred foraging microhabitat seemed to determine its local abundance. Second, one microhabitat category (large open) was augmented at randomly-selected stations, and a second rodent census was made subsequently to see if the habitat manipulation changed the rodent community inhabiting treated stations in a predictable way. Short-term increases in the abundance of Dipodomys merriami, the large open specialist, would indicate that rodent assemblages can adjust to changes in the spectrum of available microhabitats.

Phase 1: Pre-manipulation Rodent- Microhabitat Relationships

Methods: Microhabitat Censuses

The previous experiments indicated that large open spaces, small open spaces, ground under large bushes and trees were preferred foraging microhabitats for Dipodomys merriami, Perognathus amplus, P. baileyi and P. penicillatus, respectively; and in addition that pebbly surface soil might be a distinguishing characteristic of P. baileyi's foraging areas. These aspects of habitat structure were censused in a circular area within ten meters of the center of each trapping station. First, maximum height and canopy diameter were recorded for each tree or shrub over

60 cm in height that occurred within a station. From these data total coverage in square meters (assuming circular canopies) was calculated for three vegetation height categories: 60-100 cm (small bushes), 101-200 cm (large bushes), and over 200 cm (trees). These values were estimates of the abundances at each station of bush and tree microhabitats. Secondly, radial line transects were made of the vegetation between zero and 15 cm above the ground. The amount (in meters) of various microhabitat categories intersecting the six transects was an estimate of the ground-level abundance of large open bare patches (vegetationless spaces at least one meter in diameter), small open bare patches (diameter less than .5 m), grass cover, and bushes. Use of 54 meters of transect (six radial lines starting one meter and ending ten meters from the center) reduced the standard error of the mean cm of large open habitat per meter of transect to a constant minimal value. Since large open had the largest standard deviation of microhabitats censused by transects, six radii were adequate sample sizes over-all.

In addition to these variables, estimates of topographic and surface soil characteristics of each station were made (1) by counting the number of washes or drainages intersecting the ten-meter radial area of each station; and (2) by sieving a composite of four surface soil samples and calculating the weight percent of "pebbles" (particles greater than 5.6 mm in diameter) and "coarse sand" (particles less than 5.6 mm and greater than 2 mm in diameter). The four samples were taken by moving six paces due north from the center of each station and collecting a trowel full of surface soil. This was repeated for the

three remaining points of the compass and the samples mixed in a paper bag.

A total of ten variables were thus measured to characterize the availability of various foraging microhabitats at each grid station.

Methods: Rodent Censuses

Rodents were censused on two occasions for seven nights. The first census occurred prior to manipulation from 27 June through 19 July 1975; the second occurred six weeks after the manipulation from 9-20 September 1975. Each station comprised four Sherman or National metal live traps placed at even intervals along the circumference of a six-meter radius circle around the station's center. Specific trap locations were rotated counterclockwise by two meters between trapping nights to reduce the tendency of some rodents to return repeatedly to a known reward location. At dusk traps were rotated, opened, baited with rolled oats; at dawn they were checked and closed for the day. Mice were toe-clipped for individual recognition; and for each capture event date, rodent species, sex, individual, and station were recorded. These data in final form were expressed, by station, as the total number of individuals of each species (density) that were caught at least once during the seven-night census. Data were not expressed as number of captures in order to reduce inequalities in between-station capture rates caused solely by repeated captures of "trap-happy" individuals at single locations. For the post-manipulation censuses the number of "new" individuals not previously recorded in the July census as well as the

number of "old" individuals disappearing between census periods were also tallied for each station.

Statistical Methods

Statistical techniques combining principle component analysis with multivariate regression were used for evaluating the dependence of rodent abundance on microhabitat availability. Because microhabitat variables were not strictly independent of each other in the study area--grass, for example, was associated with trees which in turn were clustered along sandy washes--a straightforward multiple linear regression technique was not used. Instead principle components (eigenvectors of the matrix of correlation coefficients between microhabitat variables) describing the major modes of behavior of the microhabitat variables were first extracted, and the normalized vegetation data of each station were converted without any loss of information (by vector multiplication with each principle component) to a new set of "amplitudes" or "station scores." These derived variables have the virtue of being uncorrelated, and are thus effective as independent variables in multiple regressions.

Stepwise multiple linear regressions of per station rodent density on the derived independent variables (amplitudes) were run. Variables with insignificant regression coefficients ($F < 1$) were excluded. The matrix of significant regression coefficients of principle components was then converted to a new set of coefficients associated with each of the original microhabitat variables by multiplying the regression coefficients by their corresponding principle components. This technique

is identical to that described in detail by Fritts et al. (1971) in analyzing tree ring-climate relationships.

Principle components and their amplitudes were extracted by a computer program (PRINCY) developed by the University of Arizona Laboratory of Tree Ring Research. Amplitudes and rodent densities were then run through an S.P.S.S. stepwise multiple linear regression program (Nie et al. 1975) and coefficients of significant variables were later converted by hand to regression coefficients associated with the original ten microhabitat variables.

Results: Pre-manipulation Dependence of Rodent Abundance on Microhabitat Availability

Density of each heteromyid rodent species was highly correlated with the abundance of that microhabitat previously found to be its preferred foraging patch (see Appendix C for summary data).

Vegetation principle components and their significance levels are in Table 4. Positively correlated variables are given weightings of the same sign in any one vector. For example, principle component 1 expresses a positive association between tree cover (-.34 weighting factor), large open spaces (-.44) and washes (-.43); and negative relationships between these factors and small open (.38), and small bushes (.30). Thus this first component pulled out as a major vegetation pattern a tree-wash-large open space association as distinct from the small bush-small open association. Principle component (2) describes a positive association between low and high bushes and negative association between bushes and trees. Although non-significant principle components (in this case,

Table 4. Principle components of vegetation variables. -- The percent of over-all variance in vegetation accounted for by each vector is given in the last column.

Vector Number	Eigenvectors										Percent of Variance
	Small Bush	Large Bush	Tree	Large Open	Small Open	Grass	Bush	Washes	Pebbles	Sand	
1	-.064	.008	-.339	-.436	.382	-.146	.303	-.428	-.388	-.313	.333
2	-.548	-.585	.378	-.200	.193	.232	-.270	-.010	-.057	-.083	.183
3	.084	-.064	.021	.242	-.527	.500	.096	-.224	-.278	-.514	.125
4	.399	-.162	-.080	-.449	.130	.614	.280	.242	.227	.157	.097
5	.163	-.382	.425	.054	-.200	-.307	.608	-.078	-.223	.291	.085
6	.013	-.452	-.290	.113	-.046	-.301	.192	.128	.572	-.475	.059
7	.696	-.286	.168	.032	.196	-.134	-.505	-.283	-.064	-.096	.050
8	-.134	-.124	-.246	.030	-.196	.137	-.034	-.720	.368	.439	.037
9	-.001	.421	.617	-.120	.151	.019	.175	-.290	.446	-.302	.029
10	.011	.052	.040	-.691	-.617	-.275	-.237	.058	.028	-.049	.001

components 4 through 10) usually are discarded prior to using their amplitudes in multiple regressions (Fritts et al. 1971), this was not done here because rodents may not have confined their attention to features that contribute major amounts to over-all patterns of variation in vegetation. Indeed, the regressions show that in some cases an insignificant principle component correlated significantly with rodent density: P. baileyi density, for example, was most strongly correlated with the ninth principle component.

Table 5 presents summary data of the rodent versus principle component regressions for the four heteromyid species resident on the study site. All regressions were significant at the .04 level at least, indicating that variations in microhabitat availability influenced rodent abundance.

If we convert the partial regression equations (those including only principle components entering the regression with an F-ratio greater than 1) to equations expressed in terms of the ten original microhabitat variables (Table 6) we can assess the relative importance of each species' preferred microhabitat category in determining its density.

In all four cases, variables with highest positive regression coefficients are those which reflect the availability of preferred foraging microhabitats: D. merriami was highly associated with stations having an abundance of open spaces; P. penicillatus was dense where there were trees (and therefore grass); P. baileyi was associated with tree and large bush cover next to pebbly washes; P. amplus was abundant where

Table 5. Summary table of stepwise multiple linear regressions of pre-manipulation rodent densities on microhabitat principle component amplitudes.

Step	Principle Component Entered	F to Enter	Significance	R Square	Simple R	Over-all F	Significance	Regression Coefficient
Dependent Variable: <u>D. merriami</u> Density								
1	4	10.698	.002	.185	-.430	10.698	.002	-.431
2	3	3.116	.084	.237	.227	7.148	.002	.201
3	10	1.936	.171	.268	-.178	5.508	.003	-1.453
4	2	1.462	.233	.292	-.153	4.539	.004	-.112
5	8	.833	.366	.306	.116	3.784	.006	.189
6	5	.617	.437	.316	-.100	3.228	.011	-.107
7	7	.423	.519	.323	.084	2.789	.018	.116
8	6	.223	.639	.326	.061	2.422	.031	.679
9	9	.191	.664	.330	.057	2.131	.050	.106
10	1	.051	.822	.330	.030	1.876	.080	.162
Dependent Variable: <u>P. amplus</u> Density								
1	1	23.270	.000	.331	.575	23.270	.000	.260
2	7	3.643	.063	.380	-.222	14.111	.000	-.258
3	9	3.275	.077	.422	-.205	10.964	.000	-.316
4	5	1.252	.269	.438	.126	8.582	.000	.113
5	2	.876	.355	.449	.106	7.021	.000	.064
6	10	.858	.360	.460	.105	5.975	.000	.717
7	6	.476	.494	.467	.079	5.125	.000	.085
8	4	.308	.582	.471	.064	4.448	.001	.053
9	3	.235	.630	.474	-.056	3.904	.001	-.042
10	8	.035	.853	.474	.022	3.430	.003	.030

Table 5. -- Continued

Step	Principle Component Entered	F to Enter	Significance	R Square	Simple R	Over-all F	Significance	Regression Coefficient
<u>Dependent Variable: P. penicillatus Density</u>								
1	1	16.218	.000	.256	-.506	16.218	.000	-.423
2	2	7.727	.008	.363	.327	13.133	.000	.369
3	3	4.735	.035	.424	.246	11.045	.000	.336
4	7	3.482	.069	.466	.206	9.611	.000	.442
5	6	1.290	.262	.482	-.125	7.998	.000	-.248
6	9	.991	.325	.494	.109	6.828	.000	.312
7	5	.826	.369	.504	.100	5.947	.000	.165
8	4	.479	.493	.510	-.076	5.197	.000	-.118
9	8	.361	.551	.514	-.067	4.586	.000	-.169
10	10	.063	.803	.515	-.029	4.035	.001	-.359
<u>Dependent Variable: P. baileyi Density</u>								
1	9	4.498	.039	.087	.296	4.498	.039	.626
2	8	4.725	.035	.172	-.292	4.790	.013	-.545
3	6	1.780	.189	.204	.177	3.841	.016	.262
4	2	1.025	.317	.222	-.135	3.138	.024	-.113
5	10	.402	.529	.229	.085	2.557	.041	.798
6	1	.131	.719	.232	.049	2.110	.072	.030
7	5	.079	.780	.233	-.038	1.780	.118	-.047
8	3	.043	.837	.234	-.029	1.526	.179	-.029
9	4	.011	.916	.234	-.015	1.324	.256	-.017

Table 6. Pre-manipulation partial regression of rodent density against significant ($F \geq 1$) principle components of vegetation variables. -- Regression coefficients in the table are associated with the original vegetation variables, and have been derived by multiplication of principle components with their associated regression coefficients, and addition of like terms. The largest positive and negative coefficients are underlined.

Species	Microhabitat Variable										Constant of Regression
	Lowbush Cover	Medium Bush Cover	Tree Cover	Large Open Space	Small Open Space	Grass and Forbs	Bush (Ground Level)	Number of Washes	Weight % Pebbles	Weight % Coarse Sand	
<u>Dipodomys merriami</u>	-.110	.047	-.061	<u>1.267</u>	.712	.210	.273	<u>-.232</u>	-.188	-.091	1.408
<u>Perognathus amplus</u>	-.177	-.100	<u>-.278</u>	-.078	-.021	-.044	<u>.222</u>	.045	-.251	.072	1.184
<u>Perognathus penicillatus</u>	.158	-.254	<u>.436</u>	.178	-.170	.331	<u>-.466</u>	-.055	-.120	.004	1.857
<u>Perognathus baileyi</u>	.138	.279	<u>.401</u>	-.039	.167	-.168	.209	.246	.235	<u>-.544</u>	1.980

scattered low bushes provided many small open areas between bush canopies.

These data suggest that in general the availability of preferred foraging microhabitat types seems to determine local abundance of heteromyid rodents.

Phase Two: Rodent Responses to Microhabitat Manipulation

Methods

Regressions relating static patterns of rodent density to vegetation suggest, but do not demonstrate, that variations in the dependent variable are direct responses to the independent variables. Evidence for such cause-effect relationships must come from perturbation experiments in which manipulation of independent variables results in subsequent changes in the dependent variable consistent with predictions from static regressions.

The simplest manipulation that would yield predictable changes in rodent densities was augmentation of large open spaces. Accordingly 24 of the 49 grid stations were chosen from a random numbers table for manipulation and the remaining stations left as controls for possible seasonal shifts in rodent activity or density. At each experimental station one-half of the bushes that were between 60 and 100 cm in height and had a canopy diameter greater than one-half meter were chopped off at ground level and the brush hauled at least 100 meters off the study grid. This was done by rotating a ten-meter string around the center of each station and removing every other appropriately-sized bush that the

string contacted in its sweep. The canopy diameter of each bush was recorded before its removal to provide an estimate of the total area of large open spaces added to each station. Because the removed bushes (mostly Haplopappus tenuisectus and Baccharis brachyphylla) began flowering in September during the seven-day post-manipulation rodent census, it was unlikely that the manipulation had any effect on over-all seed density on the grid, and any response by the rodents could be considered a response to changed microhabitat abundance. A six-week period elapsed before the post-manipulation rodent census was initiated.

Results

From the equations in Table 6, we predict that only D. merriami should have a strong response to the augmentation of large open microhabitat, as this factor weighted low in the regressions for the other three species. No species should respond directly to the removal of low bushes, because Lowbush and Bush (alternate measures of the same variable) always had coefficients of inverse sign and/or were weighted very low relative to other factors.

Since the brush removed from manipulated stations averaged 4.34 percent of their 3.14 m² area, an average of 2.34 meters (.0434 x 54 meters of transect) were added to the pre-manipulation mean of 15.23 meters of large open per manipulated station. The regression equation for D. merriami predicts this species should increase in density at these stations by 2.97 individuals per station (1.27 individuals per meter of large open x 2.34 meters of large open added).

Table 7 compares the mean number of individuals recorded at manipulated versus non-manipulated (control) stations during the pre- and post-manipulation censuses. As expected, because manipulated stations were initially chosen at random, t-tests of the difference of means show no significant pre-manipulation differences in density of any species between manipulated and control stations; nor did P. pencilatus, P. baileyi, or P. amplus densities vary significantly between control and manipulated stations after the manipulation. This was as predicted from the regressions. The mean number of D. merriami per manipulated station increased by .29 individuals between census periods even though the controls showed a slight decrease over the same time period. However, the post-manipulation means were not significantly different at the .05 level between station types.

Nevertheless, since the response of D. merriami was in the proper direction, a regression was run to see if there was a relationship between the change in D. merriami density at each manipulated site and the amount of large open space that had been added to the station. Figure 6 shows a graph of this relationship. The correlation coefficient ($r = .62$) is significant at the .01 level. No other species demonstrated a significant correlation with the degree of augmentation.

The slope of the regression line (1.43) indicates that 3.35 individuals were added to a station per 2.34 meters of large open--very nearly the change predicted by the original regression equation. Given these remarkably coincident slopes (1.27 vs. 1.43), it is not immediately clear why the actual mean changes in D. merriami density were so

Table 7. Mean and standard deviation of per-station rodent density at control and manipulated stations before and after manipulation.

Species	Pre-manipulation				Post-manipulation			
	Manipulated Stations		Control Stations		Manipulated Stations		Control Stations	
	$\bar{x}$	s	$\bar{x}$	s	$\bar{x}$	s	$\bar{x}$	s
<u>Dipodomys merriami</u>	1.417	.974	1.360	.995	1.708	.954	1.240	1.268
<u>Perognathus amplus</u>	1.167	.761	1.200	.913	0.750	.737	0.440	.651
<u>Perognathus penicillatus</u>	1.917	1.472	1.880	1.641	1.875	1.035	1.800	1.190
<u>Perognathus baileyi</u>	1.667	1.129	2.280	1.100	1.000	.722	1.360	.638

Figure 6. The response of *Dipodomys merriami* to augmentation of large open spaces. -- The change in *D. merriami* density is measured as the number of individuals counted at a station in the post-augmentation census minus the number recorded there prior to augmentation.

much smaller than predicted. The discrepancy can be resolved if we postulate that stations receiving less than a certain degree of perturbation produced no response in D. merriami. Indeed, from the regression equation given in Figure 6, we find that D. merriami density tended to decline at stations which received less than 2.15 additional meters of large open, and that only perturbations in excess of the 2.15 threshold seemed to be relevant to the rodents. Thus, the actual average effective perturbation was probably .19 meters (2.34-2.15). Replacing this value in the original regression equation, the predicted average density change should have been .24 individuals per station ($1.27 \times .19$). The actual difference between post-manipulation means at control and manipulated stations (correcting for the initial pre-manipulation difference between these means) was .41 D. merriami.

In part the decreased density at minimally perturbed stations was due to kangaroo rat movements from these into more perturbed areas. Some individuals relocated between pre- and post-manipulation censuses; six shifts to and six shifts from manipulated stations occurred. The mean perturbation of stations losing individuals was significantly less than that of stations gaining relocated individuals (2.07 vs. 2.79 meters; $t = 2.31$, $p = .023$, one-tailed).

By following the appearance in the post-manipulation census of individuals not previously recorded on the grid, we can examine in more detail D. merriami's response to augmentation. Increased kangaroo rat density at manipulated stations was a function of differential invasion of these sites by immigrants. Among the manipulated stations an average

of 1.13 new individuals appeared, as opposed to a mean of .72 new kangaroo rats visiting the control stations. These means are significantly different ($t = 1.86$, $p = .04$, one-tailed). Emigration rates were not significantly different between station types.

No other species exhibited differential immigration rates between station types. However, more P. baileyi individuals initially resident at control stations disappeared between censuses (1.32 vs. .83, $t = 1.98$, $p = .028$). This appears to be a density-dependent seasonal change in density or activity, as there was initially a higher average density of P. baileyi at the control stations, and the number disappearing from a station correlated significantly ($r = .72$, $p < .001$) with the initial pre-manipulation density for all stations. The differential response does not appear to be a function of the manipulation itself.

Perognathus amplus appeared to respond indirectly to the manipulation by avoiding stations with increased D. merriami density. Changes in P. amplus density between censuses were negatively correlated to changes in D. merriami density ($r = -.50$, $p < .02$), although there was no significant correlation to degree of augmentation.

The manipulation also had an effect on the relative proportions of species inhabiting manipulated stations as a whole. Table 8 totals the number and relative proportion of individuals of each species that were caught at manipulated as opposed to those caught at control stations during each census period. The proportion of D. merriami increased an additional five percent among manipulated sites as compared to control sites after augmentation, a change coincident with the

Table 8. Species composition of manipulated and control sites before and after manipulation. -- Total number (and, in parentheses, the relative proportion) of individuals recorded from each site. Categories are tallied by species and by census period.

Species	Control Stations		Manipulated Stations	
	Pre-manipulation	Post-manipulation	Pre-manipulation	Post-manipulation
<u>Dipodomys merriami</u>	28 (.19)	28 (.25)	25 (.21)	35 (.32)
<u>Perognathus amplus</u>	29 (.20)	11 (.10)	26 (.22)	15 (.14)
<u>Perognathus penicillatus</u>	38 (.26)	39 (.35)	33 (.28)	38 (.35)
<u>Perognathus baileyi</u>	49 (.34)	33 (.30)	35 (.29)	20 (.19)

average percent of each station's area that had been converted to large open microhabitat (four percent). The relative abundance of P. baileyi showed a reciprocal decrease at manipulated stations that was six percent larger than that at control stations. This may be an artifact of the five percent increase in D. merriami proportions because the absolute decrease in P. baileyi numbers was not greater than at control stations.

These results substantiate the hypothesis that species composition of this heteromyid rodent community can shift in a fine-tuned and consistent manner to accommodate small changes through time and through space in the shape of the spectrum of available resources. Only D. merriami responded directly to the augmentation of its preferred foraging microhabitat. This resulted in a rodent community having relatively more large open space specialists than before, and therefore one which provided a better fit to the changed microhabitat availability curve at manipulated stations.

DISCUSSION

It remains to be seen whether geographic patterns in diversity and species composition of heteromyid rodent communities parallel those found in these studies of variation on a local scale. If, for example, the geographic distribution of D. merriami turns out to be limited to regions in which there is a threshold availability of appropriately-sized patches of open ground, and if heteromyids in general are limited by similar microhabitat requirements, then we will have gained power to predict the distribution and abundance of an important group of desert granivores. There are some indications that extrapolating geographic patterns from "microscopic" studies of habitat preferences will be feasible. Beatley (1976) found D. merriami in Nevada to be abundant only where bush cover was sparse, and sites which had been denuded by nuclear detonations were invaded by D. merriami to the exclusion of other species. This result, and those of other studies of heteromyid density patterns (e.g., Reynolds 1950, Rosenzweig and Winakur 1969) could have been predicted, perhaps even quantitatively, from studies similar to these.

It is possible that non-heteromyid rodent assemblages are similarly structured by habitat availability, for strong macrohabitat affinities have been described for many rodents and appear to limit their distribution. Grant (1972) reviewed a large body of evidence that habitat use shifts in response to the presence of competitors, and

concludes that rodents commonly compete for space, habitat preferences arising in response to interactions among competitors. Sympatric species with similar nutritional requirements (Baker 1971) exhibit pronounced habitat differences (e.g., Wilson 1968, Larrison and Johnson 1973, Kaufman and Fleharty 1974, M'Closkey and Fieldwick 1975) that probably allow their coexistence, albeit by rendering them allopatric on a local scale. M'Closkey and Fieldwick (1975) attribute the unusual local sympatry of Peromyscus leucopus and Microtus pennsylvanicus in a southern Ontario wet prairie to the intermingling of two habitats.

This is, in broad outline, the same mechanism which allows coexistence of heteromyid rodents, although the "habitats" discriminated by cricetids may be a function of nest-site requirements and therefore may be fundamentally different from the foraging patch specializations of heteromyids.

Habitat affinities correlate with body morphology in many cricetid forms (Dice 1940, Baker 1971), and in a few cases the significance of a morphological trait has been identified. Pelage color which matches the soil helps in predator avoidance (Dice 1947); long-tailed Peromyscus are more adept at climbing than short-tailed forms (Horner 1954). In Peromyscus maniculatus bairdii and P. m. gracilis, the selection of appropriate habitats (woods vs. grass) is genetically controlled and preferences reinforced by early experience with the proper habitat (Harris 1952, Wecker 1963). Microhabitat preferences in the heteromyid species studied here may be in part genetically fixed also, for each species maintained a preferred microhabitat even in the absence of competitors.

As yet we do not understand how habitat structure might constrain the optimal morphology of a foraging heteromyid rodent, and therefore we do not know why D. merriami (for example) should prefer open patches, nor can we predict a priori which forms should supplant D. merriami in other microhabitats. Empirical observations would suggest that areas with very large openings between very sparse bushes should support bipedal forms larger than D. merriami (such as D. deserti, Congdon 1974, Brown 1975); and brushy regions with small open spaces should support smaller, quadrupedal rodents (such as Perognathus species, Rosenzweig and Winakur 1969). Once the functional relationship between microhabitat and morphology is understood, it will perhaps be possible to predict the "morpho-species" of resident rodents from measurements of vegetation structure. Brown (1975) has documented remarkably convergent patterns in the morphologies of the heteromyids coexisting in sandy-substrate environments throughout the deserts of North America. It remains to be seen whether such patterns can be predicted solely from a knowledge of variations in availability of different microhabitats in desert environments. If they can, we will have gained support for the general theory that competitive divergence and resource availability are principles structuring communities of competing species, as well as better understanding of the behavior of an important group of desert organisms.

APPENDIX A

CAPTURE DATA FROM ENCLOSURE EXPERIMENTS

Capture data are tallied by treatment, species, enclosure, microhabitat category, and trapping date. Numbers in columns are captures on sequential trap nights during a treatment; their totals are given to the right of the first column entry.

Table A-1. Treatment 1--eight individuals of one species.

Trapping Dates:		April 15, 17, 23, 25, 30 May 1, 2, 22 (1975)				May 28, 30 June 6, 15, 16, 18, 19, 21 (1975)											
Species:		<u>Dipodomys merriami</u>				<u>Perognathus amplus</u>				<u>Perognathus penicillatus</u>				<u>Perognathus baileyi</u>			
Micro-habitat:		Large Open	Small Open	Large Bush	Large Tree	Large Open	Small Open	Large Bush	Large Tree	Large Open	Small Open	Large Bush	Large Tree	Large Open	Small Open	Large Bush	Large Tree
Enclosure:																	
1W		2 17	1 10	0 19	2 9	4 26	3 22	5 22	1 15								
		0	2	4	1	8	0	4	1								
		3	2	3	1	4	1	2	1								
		3	0	3	1	2	0	3	1								
		2	1	4	0	3	3	1	2								
		2	1	1	2	2	6	2	4								
		4	2	2	0	2	5	3	4								
		1	1	2	2	1	4	2	1								
2W		4 25	1 20	1 13	1 5	2 24	3 20	2 15	2 6								
		0	6	2	0	5	4	0	1								
		4	3	0	1	3	2	3	1								
		3	3	2	0	3	3	1	0								
		2	3	2	1	3	5	3	0								
		6	1	1	0	2	1	3	0								
		3	2	2	1	4	1	3	1								
		3	0	3	1	2	1	0	1								
3W		4 23	1 15	0 11	1 9					4 14	0 7	2 16	4 16				
		1	2	2	2					3	1	3	1				
		2	1	3	0					3	1	2	3				
		4	2	0	1					2	1	1	1				
		3	3	1	1					1	1	2	1				
		4	2	1	2					1	2	2	1				
		4	2	1	1					0	1	0	3				
		1	2	3	1					0	0	4	2				
1E		5 20	1 8	0 9	0 7									1 13	4 31	4 26	1 21
		4	2	1	0									4	3	3	2
		2	0	1	1									3	2	4	3
		3	0	0	1									2	3	4	2
		1	0	0	3									0	6	3	4
		3	1	1	1									2	5	3	2
		1	2	3	1									0	6	3	5
		1	2	3	0									1	2	2	2
2E		1 19	5 23	1 12	0 4					6 24	2 20	3 22	0 14				
		1	4	1	1					4	3	4	1				
		2	3	0	1					1	4	2	2				
		5	1	1	0					5	1	2	1				
		3	1	3	1					2	1	4	4				
		4	4	1	0					2	3	1	1				
		2	1	3	1					3	3	4	3				
		1	4	2	0					1	3	2	2				
3E		3 21	2 16	1 10	1 9									3 20	4 24	3 25	2 15
		1	2	2	3									4	4	4	0
		2	3	1	0									5	3	3	2
		4	2	0	0									2	2	3	2
		2	3	0	1									2	2	3	2
		3	2	3	1									2	6	3	1
		3	2	2	0									1	2	3	4
		3	0	1	3									1	1	3	2

Table A-2. Treatment 2--four Dipodomys merriami plus eight Perognathus.

Trapping Dates: June 24, 25, 27, 28, 30 and July 1 (1975)																
Species:	<u>Dipodomys merriami</u>				<u>Perognathus amplus</u>				<u>Perognathus penicillatus</u>				<u>Perognathus baileyi</u>			
Micro-habitat:	Large Open	Small Open	Large Bush	Tree	Large Open	Small Open	Large Bush	Tree	Large Open	Small Open	Large Bush	Tree	Large Open	Small Open	Large Bush	Tree
1W	3 8	1 4	1 6	2 4	3 16	2 14	1 12	5 19								
	2	1	2	1	3	3	3	4								
	0	1	1	1	3	2	1	5								
	1	0	1	0	2	2	2	2								
	1	0	1	0	2	2	3	1								
	1	1	0	0	3	3	2	1								
2W (control)					2 14	2 12	2 12	1 7								
					2	2	1	2								
					2	2	2	2								
					2	1	3	2								
					2	3	2	0								
					4	2	2	0								
3W (control)									6 18	2 9	4 17	2 15				
									3	3	1	3				
									1	3	5	2				
									3	1	3	2				
									3	0	1	3				
									2	0	3	3				
1E	4 11	1 8	3 12	0 5									2 16	5 22	4 22	1 16
	1	2	2	1									3	4	1	5
	2	3	1	2									3	3	6	1
	2	2	1	1									4	1	6	3
	1	0	4	1									2	5	3	3
	1	0	1	0									2	4	2	3
2E	2 5	3 9	2 5	0 1					5 25	2 8	2 17	2 12				
	2	4	0	0					0	2	6	2				
	1	2	1	1					6	1	3	1				
	0	0	0	0					4	2	2	1				
	0	0	1	0					4	0	3	4				
	0	0	1	0					6	1	1	2				
3E (control)													3 16	3 18	3 20	2 16
													2	3	2	3
													2	4	3	1
													4	2	1	4
													3	2	7	2
													2	4	4	4

Table A-3. Treatment 3--eight Dipodomys merriami plus eight Perognathus.

Trapping Dates: July 9, 14, 15, 19 (1975)																
Species:	<u>Dipodomys merriami</u>				<u>Perognathus amplus</u>				<u>Perognathus penicillatus</u>				<u>Perognathus baileyi</u>			
Micro-habitat:	Large Open	Small Open	Large Bush	Large Tree	Large Open	Small Open	Large Bush	Large Tree	Large Open	Small Open	Large Bush	Large Tree	Large Open	Small Open	Large Bush	Large Tree
<u>Enclosure:</u>																
1W	5 24	2 10	4 12	1 3	1 6	2 11	5 11	0 8								
	5 3	2 1	1 1	4 3	1 4	3 1										
	8 2	2 0	1 2	4 1	3 3	2 2										
	6 3	4 1	3 2	4 3												
2W (control)					1 6	2 7	2 9	0 3								
					0 2	2 1										
					2 1	2 2										
					3 2	3 3										
3W (control)									4 11	0 4	2 7	3 13				
									2 0	2 2	3 3					
									1 1	2 2	4 4					
									4 3	1 1	3 3					
1E	4 15	1 10	3 11	0 0									0 3	4 10	1 5	4 19
	6 2	2 0	0 0										0 3	3 1	5 5	
	3 4	3 0	0 0										0 2	1 1	5 5	
	2 3	3 0	0 0										3 1	2 2	5 5	
2E	5 22	4 11	2 8	0 1					1 6	2 10	0 9	2 8				
	5 3	2 0	0 0						1 2	3 3	3 3					
	5 2	1 0	0 0						1 4	2 2	2 2					
	7 2	3 1	0 0						3 2	4 4	1 1					
3E (control)													2 9	4 14	3 20	4 11
													3 3	8 8	1 1	
													2 1	5 5	6 6	
													2 6	4 4	0 0	

APPENDIX B

SUMMARY OF DATA FROM INTACT COMMUNITY STUDY

For each trapping station, number of captures of four rodent species are given for summer (June-September 1974), winter (November 1974-January 1975) periods. Vegetation and soil parameters are also given for each station.

Table B-1. Summary of data from intact community study.

Station	Rodent Captures								Vegetation Census				Density of Trees and Shrubs					Surface Soil Texture*	Minimum Distance to a Wash (meters)
	D. merriami		P. amplus		P. penicillatus		P. baileyi		Cover on Four 8-meter radial line transects				No. of Trees	No. of Large Bushes	No. of Low Bushes	No. of Cholla	No. of Prickly Pear		
	Summer	Winter	Summer	Winter	Summer	Winter	Summer	Winter	Large Small Bush Grass										
									Open (m)	Open (m)	(m)	(m)							
1	2	4	13		7		12	1	14.4	6.5	2.7	6.9	3	2	11	3		2	8
2	0	2	13		12		5	0	12.6	6.7	3.7	7.5	2		6	2		2	9
3	2	1	17		6		5	3	12.4	6.2	4.7	8.7	3	3	1	4	1	1	8
4	0	2	17		0		1	2	14.0	9.4	3.8	4.5	4	6	2	5	1	3	3
5	1	0	14		5		6	1	8.0	17.5	1.3	3.4	3	8		3	1	3	5
6	2	6	9		1		5	0	0.0	19.9	8.7	3.9	1	1	2	4	7	2	>10
7	2	1	11		1		9	0	8.2	14.9	6.1	3.4	1	4	4	4		1	>10
8	0	1	16		4		2	0	5.1	20.2	3.6	2.4	2	4	2	2	2	3	6
9	0	0	9		2		8	5	2.8	15.5	6.1	7.6	2	2	4	2	1	3	0
10	0	3	15		1		3	0	1.3	16.8	4.7	9.1	2	5	4	2	2	2	0
11	0	0	10		0		3	1	3.3	16.1	5.7	6.8	1	3	2	3		1	>10
12	0	1	16		0		0	0	7.9	14.5	0.8	4.7	3		2	6	2	1	9
13	0	1	7		2		11	6	7.3	12.8	1.7	9.2	2	6		3	3	3	6
14	0	0	8		3		12	2	7.4	14.2	5.8	1.6	2	9	10	2	3	3	0
15	0	6	17		2		2	0	13.0	11.0	3.1	6.6	1	1	10	3	1	2	5
16	2	5	1		3		5	0	10.4	11.7	2.2	8.7	3	1	8		1	2	0
17	3	4	3		11		4	0	11.3	4.6	1.1	9.5	3	4	13	3	1	1	0
18	4	1	3		5		4	3	3.8	13.6	6.8	6.3	2	1	6	5		1	>10
19	2	5	15		10		21	5	24.1	0.5	1.6	4.6	1	14	8	2		1	4
20	6	1	4		15		0	2	5.7	11.3	2.5	12.5	3	3	5			1	10
21	2	0	12		0		6	1	3.1	15.6	6.5	6.2	3	2	1	2	1	1	>10
22	0	0	10		0		7	2	3.7	16.2	7.8	1.2	1	7	2	7	1	3	7
23	0	0	4		1		10	2	4.0	12.1	4.4	11.5	4	4	4	4	1	3	4
24	3	1	19		0		0	0	3.0	14.1	5.7	7.7	1	1	6	7		1	>10
25	4	4	16		1		0	1	7.6	13.2	4.7	6.5	2		1	4	2	1	>10
26	2	3	10		1		3	0	6.0	13.8	4.2	8.0	2	3	1	1		1	>10
27	5	3	7		6		1	0	10.2	13.9	1.5	3.2	3	5	2	1		1	>10
28	2	1	11		4		2	1	5.0	11.9	2.3	12.8	3	1		5	3	1	>10
29	2	1	8		19		3	0	15.7	8.8	3.4	1.8	2	3	5	3		1	>10
30	1	2	20		10		1	0	7.9	11.1	4.7	7.3	4	2	1	5	1	1	>10
31	0	2	2		19		7	2	6.4	10.2	6.1	8.3	4	4	8	1	1	2	0
32	1	1	9		10		0	1	7.3	5.8	1.5	17.4	4	8	7	1		2	2
33	11	5	11		6		1	0	10.3	11.9	9.8	0.0	2	4	10	3	1	1	>10

* 1 - fine; 2 = coarse; 3 = pebbly.

APPENDIX C

SUMMARY OF RODENT CAPTURE AND VEGETATION CENSUS DATA FROM MANIPULATION STUDY

Vegetation measurements, bush cover removed, and rodent censuses are given.

Table C-1. Summary of vegetation census data from manipulation study.

Station	Manipulation Bush cover removed (m ²)	Pre-manipulation Vegetation Census						Tree Cover (m ²)	Number of Washes	Weight % Pebbles	Weight % Sand
		Large Open (m)	Small Open (m)	Grass and Forbs (m)	Bush (m)	Low Bush Cover (m ²)	Large Bush Cover (m ²)				
A1	15.40	27.90	11.50	10.30	6.00	45.8	18.0	32.6	1	9.7	10.8
A2	15.12	7.50	25.20	11.50	6.80	9.6	15.2	0.0	0	0.4	5.9
A3	17.34	13.80	27.40	8.50	8.10	24.8	43.2	2.6	2	12.9	15.9
A4		10.90	25.80	12.90	4.40	18.6	13.0	12.2	0	2.2	7.1
A5	13.84	8.20	30.80	10.30	4.50	13.8	7.4	0.0	0	4.9	8.1
A6		12.50	27.30	8.70	5.70	22.2	1.0	19.6	0	4.6	11.8
A7		16.80	24.30	9.20	3.70	15.0	15.8	58.2	1	8.4	17.9
B1	19.34	11.70	24.70	9.90	7.50	38.6	32.7	7.1	1	5.6	17.5
B2		19.20	17.80	6.80	11.50	29.4	34.4	50.6	0	5.0	16.2
B3		17.50	20.50	5.50	9.50	20.9	17.5	15.9	0	5.8	14.9
B4	18.05	7.90	26.50	10.70	8.90	13.8	18.0	0.0	0	1.9	13.0
B5	12.38	12.90	25.90	9.40	4.70	19.6	9.1	0.0	0	0.9	10.8
B6		9.10	28.90	10.60	7.30	12.3	11.2	62.4	1	7.6	14.1
B7		20.40	19.20	12.20	2.20	11.5	4.2	81.9	1	13.9	13.7
C1		9.20	31.50	11.70	1.40	20.5	45.2	0.0	1	7.5	11.2
C2		13.90	22.00	8.40	8.60	9.4	14.2	13.8	0	4.2	11.8
C3	19.70	6.90	31.50	8.10	7.50	15.0	19.4	19.6	0	3.2	14.8
C4		4.60	32.90	8.60	9.40	9.5	24.9	0.0	0	1.2	10.5
C5	15.40	14.20	17.30	13.50	3.70	12.3	14.6	41.8	0	6.5	15.2
C6	13.43	18.80	20.00	15.10	1.40	7.3	5.4	94.4	1	6.9	10.8
C7	17.04	16.00	28.10	6.10	3.50	28.9	20.7	4.5	0	18.5	10.5
D1		13.20	22.30	10.30	9.90	28.4	18.4	23.0	0	1.8	9.3
D2	11.02	10.80	25.90	11.40	5.70	8.6	14.4	25.5	0	9.7	14.2
D3	18.76	11.20	23.80	5.80	13.20	19.2	13.5	0.0	0	1.4	8.6
D4	9.69	12.50	24.90	9.30	7.30	8.3	16.3	0.0	0	4.3	7.8
D5		4.80	33.40	9.50	5.30	8.1	7.0	0.0	0	2.9	10.5
D6		19.90	15.50	15.30	5.00	18.6	8.1	25.1	2	28.3	18.2
D7	14.18	7.70	34.50	8.70	3.10	15.0	11.3	40.2	0	7.0	17.2
E1	9.69	20.60	20.50	10.60	2.30	11.5	52.5	10.8	0	6.2	13.4
E2		2.20	28.80	11.90	9.30	16.2	7.8	18.5	0	3.0	9.9
E3		12.90	25.00	6.90	9.10	8.2	16.5	35.1	0	7.8	8.0
E4		25.30	16.10	8.10	4.50	5.4	25.8	20.8	1	6.0	7.8
E5	9.23	28.30	17.20	5.70	2.80	6.5	18.2	53.5	2	12.7	20.0
E6		18.10	21.80	10.50	3.60	25.2	31.5	29.2	1	6.9	14.8
E7		14.00	24.50	8.60	7.50	14.8	8.5	38.4	0	3.9	9.2
F1	11.72	26.60	14.70	8.00	4.70	29.7	65.2	0.0	0	7.5	13.7
F2		9.40	35.20	3.50	5.90	7.2	19.7	19.6	0	7.4	8.4
F3	10.22	25.50	21.60	4.90	3.70	12.8	32.3	22.2	0	2.9	10.7
F4		19.90	24.10	8.20	1.90	11.2	16.4	21.6	1	18.7	12.8
F5		19.90	20.20	10.70	2.20	3.6	0.0	68.7	1	5.4	14.8
F6	13.62	29.00	15.20	6.30	3.50	13.8	13.1	32.2	1	10.3	17.9
F7		14.60	28.50	6.10	3.80	18.5	28.3	17.5	0	7.8	8.1
G1	8.90	9.50	32.10	6.10	4.30	10.4	5.7	28.3	0	8.3	16.5
G2	7.76	10.80	32.40	7.60	2.90	4.7	12.7	28.3	0	1.9	12.5
G3		10.00	24.50	11.90	10.80	11.4	31.4	0.0	0	3.7	8.3
G4	13.58	20.10	19.70	11.50	2.70	17.1	3.3	47.9	1	6.0	10.0
G5		24.00	10.90	15.80	6.30	18.0	18.9	28.5	0	9.7	10.3
G6		24.40	16.00	11.60	2.00	13.3	9.0	35.5	1	10.0	11.1
G7	12.54	20.90	17.90	12.20	3.00	8.2	9.5	28.3	0	3.2	10.0
$\bar{x}$	13.66	15.23	23.80	9.49	5.56	15.77	18.38	24.81	.41	6.87	12.17
s	3.48	6.78	6.12	2.76	2.91	8.68	13.23	22.57	.61	5.15	3.44

Table C-2. Summary of rodent census data from manipulation study.

Station	Pre-manipulation				Post-manipulation			
	Rodent Density (No. Individuals)		Rodent Captures (No. Captures)		Rodent Density		Rodent Captures	
	Dm	Pa	Pp	Pb*	Dm	Pa	Pp	Pb
A1	2	0	4	2	6	0	6	5
A2	2	2	1	1	7	2	1	1
A3	0	0	0	3	0	0	0	9
A4	2	1	1	1	6	2	2	3
A5	2	2	2	2	4	2	2	4
A6	2	2	0	3	3	4	0	3
A7	2	0	4	0	3	0	9	0
B1	0	2	2	3	0	2	6	4
B2	2	1	0	2	3	1	0	8
B3	0	2	3	2	0	3	3	4
B4	1	2	1	0	4	3	1	0
B5	1	1	0	0	1	1	0	0
B6	1	1	2	2	4	2	3	8
B7	0	0	7	2	0	0	12	2
C1	1	1	1	4	6	3	2	6
C2	2	3	0	5	2	3	0	14
C3	0	3	1	2	0	6	1	3
C4	1	3	0	3	1	5	0	8
C5	0	1	3	1	0	1	4	2
C6	2	1	4	4	3	1	5	4
C7	2	1	1	3	8	1	1	6
D1	2	1	1	2	2	2	1	11
D2	1	1	1	1	1	1	2	3
D3	1	2	0	2	1	2	0	8
D4	2	1	0	3	7	3	0	5
D5	2	1	2	1	6	3	2	3
D6	1	1	2	1	2	1	5	1
D7	2	1	3	0	5	1	8	0
E1	2	1	1	1	2	2	1	2
E2	0	2	1	3	0	4	1	4
E3	1	1	2	4	3	1	2	11
E4	4	2	2	2	8	4	2	2
E5	1	1	1	1	2	4	1	1
E6	0	0	2	3	0	0	2	6
E7	3	1	2	2	4	2	4	4
F1	4	0	2	2	5	0	4	2
F2	1	2	3	2	2	2	5	4
F3	2	0	3	3	2	0	6	6
F4	1	0	0	3	1	0	0	9
F5	1	0	4	2	1	0	8	2
F6	2	1	3	1	4	2	7	1
F7	2	1	0	2	7	2	0	5
G1	2	1	1	2	7	2	1	2
G2	1	2	1	2	1	2	1	2
G3	1	2	2	1	1	4	4	2
G4	0	1	5	1	0	1	8	4
G5	2	0	3	2	2	0	4	2
G6	0	2	3	3	0	3	3	4
G7	2	1	4	0	6	2	5	0

* Dm = Dipodomys merriami; Pa = Perognathus amplus; Pp = Perognathus penicillatus;
Pb = Perognathus baileyi.

LITERATURE CITED

- Abrams, P. 1975. Limiting similarity and the form of the competition coefficient. *Theor. Pop. Biol.* 8:356-375.
- Baker, R. H. 1971. Nutritional strategies of myomorph rodents in North American grasslands. *J. Mammal.* 52:800-805.
- Beatley, J. C. 1976. Environments of kangaroo rats (*Dipodomys*) and effects of environmental change on populations in southern Nevada. *J. Mammal.* 57:67-93.
- Brown, J. H. 1973. Species diversity of seed-eating desert rodents in sand dune habitats. *Ecology* 54:775-787.
- Brown, J. H. 1975. Geographical ecology of desert rodents. In M. L. Cody and J. M. Diamond (eds.) *Ecology and evolution of communities*. New Haven, Conn.: Belknap Press of Harvard Univ. Press, pp. 315-341.
- Brown, J. H. and G. L. Lieberman. 1973. Resource utilization and co-existence of seed-eating desert rodents in sand dune habitats. *Ecology* 54:788-797.
- Cockrum, E. L. 1975. Professor of Biology, University of Arizona, unpublished data.
- Cody, M. L. 1974. *Competition and the structure of bird communities*. New Jersey: Princeton University Press.
- Colwell, R. C. and D. J. Futuyma. 1971. On the measurement of niche breadth and overlap. *Ecology* 52:567-576.
- Congdon, J. 1974. Effect of habitat quality on the distributions of three sympatric species of desert rodents. *J. Mammal.* 55:659-662.
- Courtney, M. 1975. Doctoral Candidate, Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, personal communication.
- Dice, L. R. 1940. Ecologic and genetic variability within species of *Peromyscus*. *Amer. Natur.* 74:212-221.
- Dice, L. R. 1947. Effectiveness of selection by owls of deer mice (*Peromyscus maniculatus*) which contrast in color with their background. *Contrib. Lab. Vert. Biol. Univ. Mich.* 34:1-20.

- Douglass, R. J. 1976. Spatial interactions and microhabitat selections of two locally sympatric voles, Microtus montanus and Microtus pennsylvanicus. *Ecology* 57:346-352.
- Fritts, H. C., T. J. Blasing, B. P. Hayden, and J. E. Kutzbach. 1971. Multivariate techniques for specifying tree growth and climate relationships and for reconstructing anomalies in paleoclimate. *J. Appl. Meteorol.* 10:845-864.
- Gause, G. F. 1934. The struggle for existence. Baltimore, Maryland: Williams and Wilkins.
- Goodall, D. W. and S. J. Morgan. 1974. Seed reserves in desert soils. *US/IBP Desert Biome RM 74-16*.
- Grant, P. R. 1972. Interspecific competition among rodents. *Ann. Rev. Ecol. and Syst.* 3:79-106.
- Grinnell, J. and R. T. Orr. 1934. Systematic review of the Californicus group of the rodent genus Peromyscus. *J. Mammal.* 15:210-220.
- Harris, V. T. 1952. An experimental study of habitat selection by prairie and forest races of Peromyscus maniculatus. *Contrib. Lab. Vert. Biol. Univ. Mich.* 56:1-53.
- Hoagstrom, C. W. 1976. Ecologic distribution of nocturnal rodents in a part of the Sonoran Desert. Unpublished doctoral dissertation, University of Arizona, Tucson (in preparation).
- Horner, B. E. 1954. Arboreal adaptations of Peromyscus, with special reference to use of the tail. *Contrib. Lab. Vert. Biol. Univ. Mich.* 61:1-85.
- Kaufman, D. W. and E. D. Fleharty. 1974. Habitat selection by nine species of rodents in North-central Kansas. *S. W. Natur.* 18:443-452.
- Kenagy, E. J. 1972. Saltbush leaves: excision of hypersaline tissue by a kangaroo rat. *Science* 178:1094-1096.
- Larrison, E. J. and D. R. Johnson. 1973. Density changes and habitat affinities of rodents of shadscale and sagebrush associations. *Great Basin Natur.* 33:255-264.
- Lemen, C. 1976. Doctoral Candidate, University of New Mexico, Albuquerque, personal communication.

- Levins, R. 1968. Evolution in changing environments. New Jersey: Princeton University Press.
- Lotka, A. J. 1925. Elements of physical biology. Baltimore: Williams and Wilkins.
- Lowe, C. H. 1964. The vertebrates of Arizona. Tucson: University of Arizona Press.
- MacArthur, R. H. 1970. Species packing and competitive equilibrium for many species. Theor. Pop. Biol. 1:1-11.
- MacArthur, R. H. 1972. Geographical ecology. New York: Harper and Row.
- MacArthur, R. H. and E. R. Pianka. 1966. On optimal use of a patchy environment. Amer. Natur. 100:603-609.
- MacMillan, R. E. 1964. Population ecology, water relations, and social behavior of a southern California semidesert rodent fauna. Univ. Calif. Publ. Zool. 71:1-59.
- May, R. H. 1973. Stability and complexity in model ecosystems. New Jersey: Princeton University Press.
- M'Closkey, R. T. and B. Fieldwick. 1975. Ecological separation of sympatric rodents (Peromyscus and Microtus). J. Mammal. 56:119-129.
- Nie, N. H., D. H. Hull, J. G. Jenkins, K. Steinbrenner, and D. H. Bent. 1975. Statistical package for the social sciences, 2nd edition. New York: McGraw-Hill, Inc.
- Pearson, O. R. 1975. An outbreak of mice in the coastal desert of Peru. Mammalia 39:375-386.
- Price, M. V. In press. Validity of live-trapping as a measure of foraging activity of heteromyid rodents. J. Mammal.
- Pulliam, H. R. 1974. On the theory of optimal diets. Amer. Natur. 108:59-75.
- Pulliam, H. R. 1975. Coexistence of sparrows: a test of community theory. Science 189:474-476.
- Reynolds, H. 1950. Relation of Merriam kangaroo rats to range vegetation in southern Arizona. Ecology 31:456-463.
- Reichman, O. J. 1975. Relation of desert rodent diets to available resources. J. Mammal. 56:731-751.

- Reichman, O. J. 1976. Dispersion Patterns of seeds in desert soils. IBP Desert Biome RM-76.
- Rosenzweig, M. L. 1973. Habitat selection experiments with a pair of coexisting heteromyid rodent species. *Ecology* 54:111-117.
- Rosenzweig, M. L. In press. Coexistence and diversity in heteromyid rodents. In B. Stonehouse and C. Perrins (eds.) David Lack Symposium. London: Macmillan.
- Rosenzweig, M. L. and P. W. Sterner. 1970. Population ecology of desert rodent communities: body size and seed husking as bases for heteromyid coexistence. *Ecology* 51:217-224.
- Rosenzweig, M. L. and J. Winakur. 1969. Population ecology of desert rodent communities: habitats and environmental complexity. *Ecology* 50:558-572.
- Schoener, T. W. 1971. Theory of feeding strategies. *Ann. Rev. Ecol. and Syst.* 2:369-404.
- Schroeder, G. E. and M. L. Rosenzweig. 1975. Perturbation analysis of competition and overlap in habitat utilizations between D. ordii and D. merriami. *Oecologia* 19:9-28.
- Smigel, B. W. and M. L. Rosenzweig. 1974. Seed selection in Dipodomys merriami and Perognathus penicillatus. *Ecology* 55:329-339.
- Sokal, R. R. and F. J. Rohlf. 1969. Biometry. San Francisco: W. H. Freeman and Co.
- Taylor, C. R., K. Schmidt-Nielsen and J. L. Raab. 1970. Scaling of energetic cost of running to body size in mammals. *Amer. J. Physiol.* 219:1104-1107.
- Titman, D. 1976. Competition between algae: experimental confirmation of resource-based competition theory. *Science* 192:463-465.
- Vandermeer, J. H. 1972. Niche theory. *Ann. Rev. Ecol. and Syst.* 3:107-132.
- Volterra, V. 1928. Variations and fluctuations of the number of individuals in animal species living together. In Royal Chapman (ed.) *Animal ecology*. New York: McGraw-Hill, pp. 409-448.
- Wecker, S. C. 1963. The role of early experience in habitat selection by the prairie deer mouse, P. maniculatus bairdi. *Ecol. Monog.* 33:307-325.

- Werner, E. E. and D. J. Hall. 1976. Niche shifts in sunfishes: experimental evidence and significance. *Science* 191:404-405.
- Whitford, W. G. 1976. Temporal fluctuations in density and diversity of desert rodent populations. *J. Mammal.* 57:351-369.
- Wilson, D. E. 1968. Ecological distribution of the genus Peromyscus. *S. W. Natur.* 13:267-274.
- Wondolleck, J. T. 1975. The influence of interspecific interactions on the forage areas of heteromyid rodents. Unpublished master's thesis, University of Arizona, Tucson.
- Yeaton, R. I. and M. L. Cody. 1974. Competitive release in island song sparrow populations. *Theor. Pop. Biol.* 5:42-58.