1 Implicit bias toward cervical cancer: provider and training differences

2 Juliana Liang¹, Katherine Wolsiefer², Colin A. Zestcott³, Dana Chase⁴, Jeff Stone^{2, *}

3 ¹University of Arizona College of Medicine – Phoenix

4 ²University of Arizona

5 ³State University of New York at Geneseo

6 ⁴University of Arizona College of Medicine – Phoenix and Creighton University at St. Joseph's

7 Hospital and Medical Center

8 *Corresponding author: Jeff Stone

9 Postal address: University of Arizona, Psychology Department, Room 454, Tucson, AZ 85721,

10 USA

11 Telephone number: (520) 626-2438

12 Fax number: (520) 621-9306

13 Email: jeffs@email.arizona.edu

14 Abstract

15 Objective

16 Implicit prejudice and stereotyping may exist in health care providers automatically without their
17 awareness. These biases can correlate with outcomes that are consequential for the patient. This
18 study examined gynecologic oncology care providers' implicit prejudice and stereotyping toward
19 cervical cancer.

20 Methods

21 Members of professional gynecologic oncology organizations were asked to complete two
22 Implicit Association Tests to determine if they implicitly associate cervical cancer with feelings
23 of anger (prejudice) and beliefs about culpability for the disease (stereotypes), compared to
24 ovarian cancer. Linear models and student t-tests examined average levels of implicit bias and
25 moderators of the implicit bias effects.

26 Results

27 One-hundred seventy-six (132 female, 43 male, 1 nonresponse; $\bar{X}_{\text{age}} = 39.18$ years, $SD_{\text{age}} = 10.58$
28 years) providers were recruited and the final sample included 151 participants (93 physicians and
29 58 nurses, $\bar{X}_{\text{age}} = 38.93$, $SD_{\text{age}} = 10.59$). Gynecologic oncology providers showed significant
30 levels of implicit prejudice, $\bar{X} = 0.17$, $SD = 0.47$, 95% CI: (0.10, 0.25), towards cervical cancer
31 patients. They also showed significant levels of implicit stereotyping of cervical cancer patients,
32 $\bar{X} = 0.15$, $SD = 0.42$, 95% CI: (0.08, 0.21). Whereas physicians did not demonstrate significant
33 levels of implicit bias, nurses demonstrated greater levels of implicit prejudice and implicit
34 stereotyping. Providers without cultural competency/implicit bias training demonstrated greater
35 bias than those who had completed such training ($p < 0.05$).

36 Conclusions

37 This study provides the first evidence that gynecologic oncology providers hold implicit biases
38 related to cervical cancer. Interventions may be designed to target specific groups in gynecologic
39 oncology to improve interactions with patients.

1. Introduction

The present research investigates one potential barrier to cervical cancer care: The negative emotions and beliefs that providers may associate with the women who contract the disease. Recent research, for example, shows that 37% of female community members attributed blame toward cervical cancer patients [1]. Another study suggests that women with cervical cancer are viewed more negatively compared to women with ovarian cancer [2]. These findings underscore the importance of understanding whether gynecologic oncology providers hold similar negative emotions and beliefs about women with cervical cancer that may impact patients' quality of cancer care and outcomes.

In contrast to self-reported, explicit emotions and beliefs, implicit emotions and beliefs come to mind relatively quickly, are less easy to control, and may influence behavior outside of conscious awareness [3,4]. In the domain of health care, several studies show that providers hold negative implicit associations toward a variety of patient groups [5–13], and that these negative implicit associations relate to lower quality care for patients [8, 24]. However, few studies to date have examined the prevalence of implicit bias toward cancer patients. One study suggests that health care providers hold implicit negative attitudes toward patients with lung cancer [15], and another study shows that negative implicit attitudes are associated with lower quality of care for African-American cancer patients [12]. The purpose of the current study was to examine implicit biases that relate to cancers of the reproductive system like cervical cancer. Specifically, the current study measured the degree to which the implicit biases toward cervical cancer reflect negative stereotypes about risky health behavior and negative emotions such as anger and frustration.

This study used the well-established Implicit Association Test (IAT) [16] to examine two forms of implicit bias toward cervical cancer versus ovarian cancer among gynecologic oncology

providers. The IAT uses respondents' reaction times or error rates to calculate the degree to which individuals associate particular target groups with specific characteristics and attributes. The present study also investigated factors that may moderate the degree to which providers hold implicit bias, such as provider's area of specialty (nurses vs. physicians), previous training in cultural competence or implicit bias, demographic variables (sex, ethnicity), and providers' explicit internal and external motivations to avoid bias.

2. Methods

2.1. Participants and procedure

A total of 176 (132 female, 43 male, 1 non-response; $M_{age} = 39$, $s_{age} = 11$) participants were recruited primarily through emails to listservs for healthcare providers in gynecologic oncology¹. Participants were invited to participate in a grant-funded study examining perceptions of patients. After consenting to participate, participants completed the study individually on a computer. First, participants completed two measures of implicit associations: one assessing prejudice and the other assessing stereotyping of cervical cancer patients. The IAT is designed to assess relative associations [16]. Building on previous research [1, 2], implicit prejudice and stereotypes towards cervical cancer were assessed relative to another prevalent cancer of the reproductive organs that is not acquired through sexual behavior: ovarian cancer. Following the implicit measures, participants provided demographic information and ratings of external and internal motivation to respond without prejudice toward women with cervical cancer. All participants were thanked, debriefed, and compensated with a \$50 Amazon gift card.

¹ Listservs included gynecologic oncology fellowship programs, Society of Gynecologic Oncology, Society of Gynecologic Nurse Oncologists, US Oncology Network of Gynecologic Oncologists, and Western Association of Gynecologic Oncologists.

2.2. Measures

2.2.1. Measures of implicit associations

Two IATs were used to assess implicit associations to cancer— one assessed implicit prejudice (i.e., negative emotions) and the other assessed implicit stereotypes (i.e., negative beliefs). The prejudice and stereotype measures were designed to examine the extent to which individuals associated cervical cancer with negative emotions and beliefs that may extend from blame.

Specifically, the prejudice measure examined the extent to which providers associated cervical cancer with anger-related emotions and the stereotype measure examined the extent to which individuals associated cervical cancer with beliefs related to risky health behavior.

The IAT requires that participants sort words related to two opposing target categories while simultaneously sorting words related to two opposing attribute categories. The IAT sorting task is based on the premise that stimuli belonging to categories that are more strongly associated in one's mind (e.g., prejudice or stereotype congruent information) will be easier to sort when they share a response key; and stimuli belonging to words that are unassociated in one's mind (i.e., prejudice or stereotype incongruent information) will be more difficult to sort when they share a

response key. A person who associates cervical cancer with anger more than empathy would

then be faster and more accurate on trials for which cervical cancer and anger share a response

key and slower on trials for which cervical cancer and empathy share a response key. As such,

implicit bias on the IAT is quantified as the extent to which an individual is faster on prejudice or

stereotype congruent trials compared to prejudice or stereotype incongruent trials.

For both the stereotype and prejudice measures, the two target categories were cervical cancer versus ovarian cancer². For the prejudice IAT, the attribute categories were words related to anger (e.g., *hostile, resentful, indignant, infuriated, offended*) versus empathy (e.g., *sympathetic, compassionate, supportive, understanding, considerate*). For the stereotype IAT, the attribute categories were words related to high "risk" (e.g., *reckless, liable, defiant, negligent, blameable*) versus low risk or "compliance" (e.g., *exempt, excusable, forgivable, justifiable, sensible*). The prejudice and stereotype IAT measures were identical except for the attribute categories and stimuli.

For each IAT, words belonging to one of the target or attribute categories appeared on the screen one at a time. For example, in the prejudice IAT, participants saw words related to ovarian cancer, cervical cancer, empathy and anger appear on the screen one at a time. Participants were instructed to respond by pressing the "E" key if the word was related to either cervical cancer or anger and to press the "I" key if the word was related to either ovarian cancer or empathy. This block would be considered a bias "congruent" block because cervical cancer and anger share a response key. In the second half of the IAT, the response mappings were reversed. That is, in the second half of the task, participants were instructed to press the "E" key in response to cervical cancer words or empathy related words and to press the "I" key in response to ovarian cancer words or anger related words. This response configuration is considered bias "incongruent" because cervical cancer and empathy share a response key. If a participant incorrectly categorized one of the words, a red X appeared on the screen until they corrected their response. Participants completed two blocks (one with 20 trials and one with 40 trials) of the bias

² Stimuli related to ovarian cancer and cervical cancer was identical across the two implicit bias measures. Ovarian cancer stimuli were: *pelvic exam, genetics, CA-125, bloating, abdomen pain*. Cervical cancer stimuli were: *pap test, HPV, Gardasil, bleeding, discharge*.

congruent and two blocks of the bias incongruent tasks. The order of the blocks (i.e., whether the bias congruent or bias incongruent blocks appeared first) was counterbalanced across participants.

The measure of bias resulting from an IAT is called a d-score. These IAT d-scores were calculated following well-established scoring algorithms [17]. After cleaning each dataset for individual trials that were too fast (< 300 ms) or too slow ($> 10,000$ ms), d-scores were calculated for each participant. To calculate these d-scores, average latency to correct response on stereotype/prejudice congruent blocks was subtracted from the average response latency (to correct response³) on stereotype/prejudice incongruent blocks. This difference score was then standardized using the standard deviation in response latency on all critical blocks. This resulted in a single IAT d-score on each IAT for each participant with higher values indicating faster (greater facilitation of) correct responding on stereotype/prejudice congruent blocks. Thus, larger positive IAT d-scores were indicative of stronger implicit stereotypes or prejudice and IAT d-scores at 0 were considered indicative of a lack of implicit stereotypes or prejudice.

2.2.2. Demographic variables

After completing the IATs, participants reported the number of years in practice, area of specialty, job title, practice setting, medical school location, and whether or not they had training in cultural competency and implicit bias during medical school. Participants also reported their sex, age, ethnicity, race, birthplace, years living in the United States, and their first language.

2.2.3. External and internal motivation to respond without prejudice

³ Latencies on trials where participants initially responded incorrectly were included in the calculation of the d-score. However the response latency used in these cases was the amount of time it took from the start of the trial until the participant corrected their response. This allows for a “time penalty” for incorrect responses.

The External and Internal Motivation to Respond Without Prejudice Scale [18] was adapted to measure explicit motivations to be non-prejudiced against women with cervical cancer. Participants used 7-point Likert scales (1 = completely disagree to 7 = completely agree) to rate the degree to which they agreed with five external motivation scale (EMS) items (e.g., “If I acted prejudiced toward women with cervical cancer, I would be concerned that others would be angry with me”) and five internal motivation scale (IMS) items (e.g., “I am personally motivated by my beliefs to be non-prejudiced against women with cervical cancer”). Ratings were averaged to create separate individual scores for the external motivation scale (EMS, $\alpha = .74$) and the internal motivation scale (IMS, $\alpha = .74$).

2.3. Data analysis

The same analysis approach was used to examine both implicit prejudice and implicit stereotypes. First, a single sample t-test was estimated to examine whether participants, on average, demonstrated implicit associations that were statistically different from neutral (i.e., IAT d-scores significantly different from 0). Second, independent samples t-tests (for categorical variables) and simple regression models were estimated to examine whether implicit associations differed as a function of participant demographic variables, variables related to medical experience and training, or motivation to respond without prejudice. Results of these models are presented in Tables 2 and 3. When there was evidence of significant moderation, single sample t-tests were used to examine the implicit bias d-score at different levels of the moderator (reported in the body of the text). Although implicit stereotypes and prejudice were related, $r = .45$, 95% CI: (0.31, 0.57), $R^2 = .20$, the correlation was not high enough to assume that the two measures captured the same construct. As such, the results are presented separately for the two implicit bias measures.

3. Results

3.1. Study sample

Of the total 176 providers recruited, 12 of the participants were recruited at the 2016 Western Association of Gynecologic Oncologists Annual Meeting and the remaining participants were recruited through emails to listservs. Of these participants, 10 (5.7%) were excluded for exhibiting high error rates ($> 30\%$) on at least one IAT. An additional 15 participants were excluded from the analyses because they identified themselves as nurse practitioners – a category distinct from both physicians and nurses. This resulted in a final sample consisting of 151 (112 female, 39 male, $\bar{X}_{\text{age}} = 39$ years, $SD_{\text{age}} = 11$) gynecologic oncology care providers (93 physicians and 58 nurses). The average number of years in practice was 12 years with a standard deviation of 11 years. 67.5% of participants reported having completed cultural competency training and 41.1% of participants reported having completed implicit bias training. Table 1 contains additional demographic information for this final sample.

3.2. Implicit prejudice

The average prejudice d-score was significantly different from 0, indicating that gynecologic oncology providers showed significant levels of implicit prejudice towards cervical cancer patients, $\bar{X} = 0.17$, $SD = 0.47$, 95% CI: (0.10, 0.25). Providers associated cervical cancer with emotions related to anger/frustration and ovarian cancer with empathy to a greater degree than the reverse. Table 2 presents tests of the potential moderators of the implicit prejudice d-score as well as means and standard deviations of the implicit prejudice d-score at different levels of the moderators. On average, nurses demonstrated greater levels of implicit prejudice than physicians (see Table 2 for supporting statistics). Whereas physicians, on average, did not demonstrate significant levels of implicit prejudice, 95% CI: (-0.02, 0.18), nurses did exhibit significant

implicit prejudice, 95% CI: (0.21, 0.44). Cultural competency training also significantly moderated implicit prejudice effects. Although individuals associated cervical cancer with anger over empathy both with and without cultural competency training $p < 0.02$, the magnitude of implicit prejudice was greater for those who did not report having received such training. Older providers and providers with more experience in the field both held stronger implicit anger associations towards cervical cancer. There was some evidence that ethnicity moderated implicit prejudice with Hispanic providers demonstrating no evidence of implicit prejudice, 95% CI: (-.49, .14), and non-Hispanic providers demonstrating significant implicit prejudice, 95% CI: (0.13, 0.28). But, there were only 13 providers who identified as Hispanic (compared to 138 who did not). Responses on the IMS and EMS scales did not significantly moderate implicit prejudice scores, both $t < 1$.

3.3. Implicit stereotypes

On average, gynecologic oncology providers showed significant levels of implicit stereotyping of cervical cancer patients, $\bar{X} = 0.15$, $SD = 0.42$, 95% CI: (0.08, 0.21), suggesting that providers more strongly associated cervical cancer with words related to risky health behavior, and ovarian cancer with words related to lower risk behavior, compared to the association between ovarian cancer and risk, and cervical cancer and low risk. Table 3 presents tests of the potential moderators of the implicit stereotype effect as well as means and standard deviations of the implicit stereotyping d-score by moderator group (for categorical moderators). As seen in Table 2, Similar to implicit prejudice, provider type significantly moderated the implicit stereotype d-score such that physicians' implicit stereotypes were not reliably different from 0, 95% CI: (-0.05, 0.13), but nurses' scores were higher than 0, indicating significant average implicit stereotypes, 95% CI: (0.22, 0.41). Although providers with and without cultural competency

training demonstrated significant implicit stereotypes, $p_s < 0.03$, implicit stereotypes were stronger for providers who reported never having participated in such training. Providers who reported never completing any implicit bias training associated cervical cancer and ovarian cancer with compliance, 95% CI: (0.01, 0.17), while providers who reported completing implicit bias training did not demonstrate significant implicit cervical cancer stereotypes, 95% CI: (-0.04, 0.16). Similar to their moderating effects on implicit prejudice, older providers and providers with more years of experience both exhibited stronger implicit stereotypes. Unlike the implicit prejudice effect, the implicit stereotype effect was marginally moderated by provider gender. Whereas males did not demonstrate significant implicit cervical cancer stereotype effects, 95% CI: (-0.09, 0.19), females did, 95% CI: (0.10, 0.26). As with the implicit prejudice measure, responses on the IMS and EMS scales did not significantly moderate implicit stereotyping scores, both $t_s < 1$.

4. Discussion

4.1. Implicit biases

The results of this study suggest that relative to ovarian cancer, gynecologic oncology providers hold implicit prejudice and stereotyping toward cervical cancer. Specifically, providers more strongly associated negative emotions related to anger and frustration with cervical cancer and positive emotions related to empathy with ovarian cancer, than they associated these negative emotions with ovarian cancer and positive emotions with cervical cancer. In addition, providers expressed stronger associations between beliefs about risky health behavior and cervical cancer than between beliefs about risky health behavior and ovarian cancer. The levels of implicit prejudice and stereotyping expressed by providers were modestly correlated, suggesting that

although the two forms of bias share common variance, each form of bias may play a distinct role in how providers perceive and treat cervical cancer patients.

4.2. Demographic moderators

Not all gynecologic oncology providers hold the same level of implicit bias toward patients with cervical cancer. In our study nurses expressed significantly greater prejudice and stereotyping of cervical cancer patients compared to physicians, which is consistent with the current literature [10], and suggests the need to understand why specific groups report higher implicit bias than others. For example, nurses may have more extended contact with cervical cancer patients than physicians, but given the evidence that contact can also be effective for reducing explicit and implicit bias toward a group and its members [19], including in a clinical context [20], we suspect that if the balance of the contact between certain providers and cervical cancer patients is negative, it will promote the development of negative implicit associations toward the patient group.. In addition, older providers and those with more years of experience reported significantly higher implicit bias compared to younger providers and those who have less experience. But, the very high correlation between age and years of practice obscures a clear interpretation of these relationships. As noted above, through negative contact, more experience caring for cervical cancer patients may engender more negative emotions and beliefs. Finally, consistent with previous research [21], there was also evidence that provider ethnicity and gender moderated the level of implicit prejudice and stereotyping, with Hispanic and male providers showing less implicit bias than non-Hispanic and female providers, respectively. These findings, however, should be interpreted with caution because they were based on small samples and only occurred on one measure.

4.3. Cultural competency and implicit bias training

Gynecologic oncology providers who report having had training in cultural competency or implicit bias expressed lower implicit bias than providers who did not. Whereas this could reflect the effectiveness of training providers to recognize the role that culture and their own biases can play in the care they provide to patients, the correlational nature of the finding could also reflect a third variable like motivation to appear low in prejudice. However, the correlations illustrated in Figure 1 show that the relationship between internal motivation to avoid prejudice and self-reports of having completed cultural competence and implicit bias training are small and not significant. The same is true of the relationship between external motivation to avoid prejudice and self-reports of training in cultural competence; only the relationship between external motivation to control prejudice and self-reports of having completed training in implicit bias, while small, is significant ($r = 0.22$, 95% CI: (0.06, 0.38)). This suggests that the relationship between cultural competency or implicit bias training and the level of implicit bias toward cervical cancer is not likely due to the desire to appear unbiased for internal or external reasons. However, the correlational nature of the finding does not permit firm conclusions that the training was a causal factor in the lower implicit bias scores.

4.4. Limitations and future directions

The data for the present study were drawn from a convenience sample of gynecologic oncology providers who are members of several professional gynecologic oncology organizations. The sample did not include those who do not belong to these organizations, and therefore, may not reflect the implicit attitudes and beliefs of all gynecologic oncology providers. The present study also did not examine other potential emotions and beliefs that gynecologic oncology providers may hold toward cervical and ovarian cancer patients. Future research should examine the degree to which providers implicitly associate other emotions and beliefs to cervical and ovarian cancer

patients that may impact their care. Another question for future study is to determine if differences in the way that nurses and physicians interact with cervical and ovarian cancer patients, above and beyond their differences in training and role in patient care, makes a unique contribution to the development and expression of implicit bias.

Another limitation is whether implicit bias toward cervical cancer reflects implicit bias towards ethnic or racial minority patients. The incidence of cervical cancer is disproportionately higher among African-American and Hispanic women than White women [22]. If gynecology care providers strongly associate cervical cancer with race or ethnicity, it is possible that the results on the IAT measures in the present study reflect biases based on these categorizations, rather than type of cancer. Future studies should separate these influences by varying the race or ethnicity of targets with cervical versus ovarian cancer.

Another future direction for research on implicit bias among gynecologic oncology providers is to examine if and when the biases influence judgment and behavior toward cervical cancer patients [12,23]. A recent study reported that the mortality rate for black women was 10.1 per 100,000 compared to 4.7 per 100,000 for white women [24]. Whereas the role that provider behavior plays in creating disparities in the survivorship of racial and ethnic women is currently unknown, if providers feel anger and frustration in the presence of a minority cervical cancer patient, or blame the patient for engaging in risky health behavior, it is possible that these emotions and beliefs may influence treatment decisions and interactions with the patient that ultimately affects the way they respond to diagnosis and treatment [20].

Finally, receiving training in cultural competency and implicit bias may reduce the tendency to automatically associate cervical cancer with negative emotions and beliefs about culpability for the disease. Nurses and older providers with more experience may especially benefit from the

training, but only if the training is effective for helping gynecologic oncology providers change their behaviors.

4.5. Conclusion

Although it may be reasonable to assume that health care providers are particularly motivated to be egalitarian [25], they hold similar intergroup implicit biases to the general public. This study provides the first evidence that gynecologic oncology providers hold negative implicit biases related to cervical cancer. Interventions may be designed to target specific groups in gynecologic oncology such as nurses, older providers, and providers who have been practicing for a longer amount of time to improve interactions with patients.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Funding

This work was supported by the Gynecologic Oncology Research Fund, St. Joseph's Foundation, and the Merck Foundation Alliance to Advance Patient-centered Cancer Care.

Author contribution

JL contributed to the study design, assisted in data collection, and worked on the final manuscript and revisions. KW completed the data analysis, wrote the results section, and edited the manuscript and revisions. CZ was instrumental in setting up the Implicit Association Test and provided feedback on the manuscript and revisions. DC contributed to the study design, facilitated data collection, and worked on the final manuscript and revisions. JS contributed to the study design, executed data collection, completed initial analyses of the data, wrote the first draft of the manuscript, and oversaw the revisions. DC, JS and JL secured funding for this study.

References

- [1] L.A. Marlow, J. Waller, J. Wardle, Variation in blame attributions across different cancer types, *Cancer Epidemiol. Biomark. Prev. Publ. Am. Assoc. Cancer Res. Cosponsored Am. Soc. Prev. Oncol.* 19 (2010) 1799–1805. doi:10.1158/1055-9965.EPI-09-1298.
- [2] M.A. Shepherd, M.A. Gerend, The blame game: Cervical cancer, knowledge of its link to human papillomavirus and stigma, *Psychol. Health.* 29 (2014) 94–109. doi:10.1080/08870446.2013.834057.
- [3] A.G. Greenwald, M.R. Banaji, Implicit Social Cognition: Attitudes, Self-Esteem, and Stereotypes, *Psychological Review.* 24 (1995) 4-27 .
- [4] B. Gawronski, G.V. Bodenhausen, Associative and propositional processes in evaluation: An integrative review of implicit and explicit attitude change., *Psychol. Bull.* 132 (2006) 692–731. doi:10.1037/0033-2909.132.5.692.
- [5] J.A. Sabin, R.G. Riskind, B.A. Nosek, Health Care Providers' Implicit and Explicit Attitudes Toward Lesbian Women and Gay Men, *Am. J. Public Health.* 105 (2015) 1831–1841. doi:10.2105/AJPH.2015.302631.
- [6] I.V. Blair, J.F. Steiner, D.L. Fairclough, R. Hanratty, D.W. Price, H.K. Hirsh, L.A. Wright, M. Bronsert, E. Karimkhani, D.J. Magid, E.P. Havranek, Clinicians' Implicit Ethnic/Racial Bias and Perceptions of Care Among Black and Latino Patients, *Ann. Fam. Med.* 11 (2013) 43–52. doi:10.1370/afm.1442.
- [7] M.G. Bean, J. Stone, G.B. Moskowitz, T.A. Badger, E.S. Focella, Evidence of Nonconscious Stereotyping of Hispanic Patients by Nursing and Medical Students:, *Nurs. Res.* 62 (2013) 362–367. doi:10.1097/NNR.0b013e31829e02ec.
- [8] A.R. Green, D.R. Carney, D.J. Pallin, L.H. Ngo, K.L. Raymond, L.I. Iezzoni, M.R. Banaji, Implicit Bias among Physicians and its Prediction of Thrombolysis Decisions for Black and White Patients, *J. Gen. Intern. Med.* 22 (2007) 1231–1238. doi:10.1007/s11606-007-0258-5.
- [9] G.B. Moskowitz, J. Stone, A. Childs, Implicit stereotyping and medical decisions: Unconscious stereotype activation in practitioners' thoughts about African Americans, *Am. J. Public Health.* 102 (2012) 996–1001.
- [10] N. Hagiwara, L.A. Penner, R. Gonzalez, S. Eggly, J.F. Dovidio, S.L. Gaertner, T. West, T.L. Albrecht, Racial attitudes, physician–patient talk time ratio, and adherence in racially discordant medical interactions, *Soc. Sci. Med.* 87 (2013) 123–131. doi:10.1016/j.socscimed.2013.03.016.
- [11] J.A. Sabin, M. Marini, B.A. Nosek, Implicit and Explicit Anti-Fat Bias among a Large Sample of Medical Doctors by BMI, Race/Ethnicity and Gender, *PLoS ONE.* 7 (2012) e48448. doi:10.1371/journal.pone.0048448.
- [12] L.A. Penner, J.F. Dovidio, R. Gonzalez, T.L. Albrecht, R. Chapman, T. Foster, F.W.K. Harper, N. Hagiwara, L.M. Hamel, A.F. Shields, S. Gadgeel, M.S. Simon, J.J. Griggs, S. Eggly, The Effects of Oncologist Implicit Racial Bias in Racially Discordant Oncology Interactions, *J. Clin. Oncol.* 34 (2016) 2874–2880. doi:10.1200/JCO.2015.66.3658.
- [13] W. von Hippel, L. Brener, C. von Hippel, Implicit Prejudice Toward Injecting Drug Users Predicts Intentions to Change Jobs Among Drug and Alcohol Nurses, *Psychol. Sci.* 19 (2008) 7–11. doi:10.1111/j.1467-9280.2008.02037.x.
- [14] T.S. Peris, B.A. Teachman, B.A. Nosek, Implicit and Explicit Stigma of Mental Illness: Links to Clinical Care, *J. Nerv. Ment. Dis.* 196 (2008) 752–760. doi:10.1097/NMD.0b013e3181879dfd.

- [15] N. Sriram, J. Mills, E. Lang, H.K. Dickson, H.A. Hamann, B.A. Nosek, J.H. Schiller, Attitudes and Stereotypes in Lung Cancer versus Breast Cancer, *PLoS ONE*. 10 (2015). doi:10.1371/journal.pone.0145715.
- [16] A.G. Greenwald, D.E. McGhee, J.L.K. Schwartz, Measuring Individual Differences in Implicit Cognition: The Implicit Association Test, *Journal of Personality and Social Psychology*, 17 (1998) 1464-1480.
- [17] A.G. Greenwald, B.A. Nosek, M.R. Banaji, Understanding and using the Implicit Association Test: I. An improved scoring algorithm., *J. Pers. Soc. Psychol.* 85 (2003) 197–216. doi:10.1037/0022-3514.85.2.197.
- [18] E.A. Plant, P.G. Devine, Internal and external motivation to respond without prejudice., *J. Pers. Soc. Psychol.* 75 (1998) 811-832.
- [19] T.F. Pettigrew, L.R. Tropp, A meta-analytic test of intergroup contact theory., *J. Pers. Soc. Psychol.* 90 (2006) 751–783. doi:10.1037/0022-3514.90.5.751.
- [20] S.E. Burke, J.F. Dovidio, J.M. Przedworski, R.R. Hardeman, S.P. Perry, S.M. Phelan, D.B. Nelson, D.J. Burgess, M.W. Yeazel, M. van Ryn, Do Contact and Empathy Mitigate Bias Against Gay and Lesbian People Among Heterosexual Medical Students? A Report from Medical Student CHANGES, *Acad. Med. J. Assoc. Am. Med. Coll.* 90 (2015) 645–651. doi:10.1097/ACM.0000000000000661.
- [21] E.N. Chapman, A. Kaatz, M. Carnes, Physicians and Implicit Bias: How Doctors May Unwittingly Perpetuate Health Care Disparities, *J. Gen. Intern. Med.* 28 (2013) 1504–1510. doi:10.1007/s11606-013-2441-1.
- [22] R.L. Siegel, K.D. Miller, A. Jemal, Cancer statistics, 2016, *CA. Cancer J. Clin.* 66 (2016) 7–30. doi:10.3322/caac.21332.
- [23] L.A. Cooper, D.L. Roter, K.A. Carson, M.C. Beach, J.A. Sabin, A.G. Greenwald, T.S. Inui, The Associations of Clinicians' Implicit Attitudes About Race With Medical Visit Communication and Patient Ratings of Interpersonal Care, *Am. J. Public Health.* 102 (2012) 979–987. doi:10.2105/AJPH.2011.300558.
- [24] A.L. Beavis, P.E. Gravitt, A.F. Rositch, Hysterectomy-corrected cervical cancer mortality rates reveal a larger racial disparity in the United States, *Cancer.* 123 (2017) 1044–1050. doi:10.1002/cncr.30507.
- [25] D.J. Burgess, S.S. Fu, M.V. Ryn, Why Do Providers Contribute to Disparities and What Can Be Done About It?, *J. Gen. Intern. Med.* 19 (2004) 1154–1159. doi:10.1111/j.1525-1497.2004.30227.x.

Table 1. Characteristics of Final Sample

Characteristic	Number	Gender, % Female	Age (Years), $\bar{X}(SD)$
Overall	151	74	39 (12)
Gender			
Male	39		38 (10)
Female	112		42 (11)
Race			
White	105	76	40 (11)
African American/Black	6	83	34 (6)
Asian or Asian American	25	64	37 (10)
Hispanic	6	667	32 (4)
Other	9	78	35 (5)
Ethnicity			
Hispanic	13	7	34 (6)
Non-Hispanic	138	68	39 (11)
Born in USA?			
Yes	129	63	39 (11)
No	22	11	40 (11)
English First Language?			
Yes	133	65	39 (11)
No	18	9	39 (11)
Provider Type			
Nurse	58	97	37 (10)
Doctor	93	57	41 (12)
Cultural Competency Training?			
Yes	102	80.4	35 (9)
No	49	61.2	47 (10)
Implicit Bias Training?			
Yes	62	79.0	42 (11)
No	89	70.8	35 (9)
Continuous Variables			
Characteristic	Number	$\bar{X}$ (SD)	
Age	151	39 (12)	
Years in USA	151	36 (12)	
Years in practice	151	12 (10)	
IMS	151	7 (1)	
EMS	151	4 (1)	

Note. This table contains descriptive statistics by race and gender for the final sample of 151 participants. The race and ethnicity items both included a “Hispanic” response so the category is reported twice. Number of responses, mean and standard deviation for continuous variables are provided in the bottom section of the table. $\bar{X}$ refers to sample mean and SD refers to the sample standard deviation.

Table 2. Moderators of the Implicit Prejudice Effects

Categorical Variables					
Moderator	$\bar{X}$ (SD)	df	t	95% CI	Cohen's D
Gender					
Male	0.10 (0.49)				
Female	0.20 (0.47)	149	-1.21	-0.28, 0.07	-0.21
Race					
White	0.21 (0.46)				
Minority	0.08 (0.50)	149	-1.54	-0.29, 0.04	-0.27
Ethnicity					
Hispanic	-0.17 (0.52)				
Non-Hispanic	0.21 (0.46)	149	2.83**	0.12, 0.65	0.80
Born in USA					
Yes	0.16 (0.48)				
No	0.27 (0.45)	149	-0.98	-0.32, 0.11	-0.23
English First Language					
Yes	0.17 (0.48)				
No	0.17 (0.42)	149	0.03	-0.23 0.24	0.00
Provider Type					
Nurse	0.32 (0.42)				
Doctor	0.08 (0.48)	149	3.16**	0.09, 0.40	0.51
Cultural Competency Training					
Yes	0.12 (0.48)				
No	0.28 (0.44)	149	-2.00*	-0.32, -0.002	-0.34
Implicit Bias Training					
Yes	0.13 (0.49)				
No	0.21 (0.46)	149	-0.98	-0.23, 0.08	-0.17
Continuous Variables					
Moderator	b	df	t	95% CI	R ²
Age	0.01	149	2.58*	0.002, 0.02	0.04
Years in USA	0.00	149	1.25	-0.002, 0.010	0.01
Years in Practice	0.01	149	2.93**	0.003, 0.018	0.05
IMS	0.02	149	0.37	-0.08, 0.12	0.00
EMS	-0.01	149	-0.21	-0.06, 0.05	0.00

Note. The race and ethnicity items both included a “Hispanic” response so the category is reported twice. Test statistics refer to tests of mean differences (categorical moderators) or tests of simple slopes (continuous predictors). $\bar{X}$ refers to sample mean and SD refers to the sample standard deviation. 95% CI refers to 95% confidence intervals of the mean differences (categorical predictors) or the 95% confidence interval for the slope (continuous predictors). * $p < .05$, ** $p < .01$,

*** $p < .001$.

Table 3. Moderators of the Implicit Stereotype Effects

Categorical Variables					
Moderator	$\bar{X}$ (SD)	df	t	95% CI	Cohen's D
Gender					
Male	0.05 (0.44)				
Female	0.18 (0.41)	149	-1.70+	-0.28, 0.02	-0.31
Race					
White	0.17 (0.43)				
Minority	0.08 (0.40)	149	-1.20	-0.23, 0.06	-0.21
Ethnicity					
Hispanic	0.01 (0.36)				
Non-Hispanic	0.16 (0.43)	149	1.19	-0.10, 0.39	0.36
Born in USA					
Yes	0.15(0.43)				
No	0.11(0.38)	149	0.45	-0.15, 0.24	0.14
English First Language					
Yes	0.15 (0.44)				
No	0.14 (0.28)	149	0.04	-0.20 0.21	0.02
Provider Type					
Nurse	0.32 (0.37)				
Doctor	0.04 (0.42)	149	4.14***	0.14, 0.41	0.67
Cultural Competency Training					
Yes	0.09 (0.40)				
No	0.26 (0.44)	149	-2.43*	-0.32, -0.03	-0.40
Implicit Bias Training					
Yes	0.06 (0.38)				
No	0.21 (0.44)	149	-2.15*	-0.28, -0.01	-0.36
Continuous Variables					
Moderator	b	df	t	95% CI	R ²
Age	0.01	149	2.85**	0.003, 0.015	0.05
Years in USA	0.01	149	2.36*	0.001, 0.011	0.04
Years in Practice	0.01	149	2.68**	0.002, 0.015	0.05
IMS	0.04	149	0.77	-0.056, 0.127	0.00
EMS	0.01	149	0.62	-0.032, 0.062	0.00

Note. The race and ethnicity items both included a “Hispanic” response so the category is reported twice. Test statistics refer to tests of mean differences (categorical moderators) or tests of simple slopes (continuous predictors). $\bar{X}$ refers to sample mean and SD refers to the sample standard deviation. 95% CI refers to the 95% confidence interval of the mean differences (categorical predictors) or the 95% confidence interval for the slope (continuous predictors). +p < .1, *p < .05, **p < .01, ***p < .001.