Title: Evidence for Multiple 4.0-3.7 Ga Impact Events within the Apollo 16 Collection

Running title: Evidence for Impact Events within A16 collection

Authors

Takafumi Niihara ^{1,2,3,4*}

Sky P. Beard ^{4,5}

Timothy D. Swindle ^{4,5}

Lillian A. Schaffer ⁶

Hideaki Miyamoto ^{1,2,}

David A. Kring ^{3,4}

1. Department of Systems Innovation, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan
2. University Museum, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
3. Center for Lunar Science and Exploration, Lunar and Planetary Institute, Universities Space Research Association, 3600 Bay Area Boulevard, Houston, Texas 77058, USA
4. SSERVI
5. Lunar and Planetary Laboratory, University of Arizona, 1629 E University Boulevard, Tucson, Arizona 85721-0092, USA.
6. Department of Earth and Atmospheric Sciences, University of Houston, 4800 Calhoun Rd., Houston, Texas 77204-5007, USA.

8 Figures

2 Tables

6 Supplements (1 Figure, 5 Tables)

Submit to Meteoritics and Planetary Science

*Corresponding Author

Takafumi Niihara (niihara@seed.um.u-tokyo.ac.jp)

Abstract

In a histogram of lunar impact ages from the Apollo 16 site, there is a spike circa 3.9 Ga that has been interpreted to represent either a large number of nearly synchronous events or an abundance of samples that were affected slightly differently by the event that produced the Imbrium basin. To further scrutinize those age relationships, we extracted six centimeter-sized clasts of impact melt from ancient regolith breccia 60016 and performed petrological and geochronological (^{40}Ar - ^{39}Ar) analyses. Three clasts have similar poikilitic textures, while others have porphyritic, apahnitic or intergranular textures. Compositions and abundances of relict minerals are different in all six clasts and variously imply Mg-suite and ferroan anorthosite target sequences. Estimated bulk compositions of four clasts are similar to previously defined group 1 Apollo 16 impact melt rocks, while the other two have higher Al_2O_3 and lower $\text{FeO}+\text{MgO}$ compositions. All six clasts have similar K_2O and P_2O_5 concentrations, which could have been derived from a KREEP-bearing component among target sequences. Eighteen $^{40}\text{Ar}/^{39}\text{Ar}$ analyses of the six clasts produced an age range from 3823 ± 75 to 4000 ± 23 Ma, consistent with estimates for the proposed late heavy bombardment. Four clasts have multiple temperature steps that define distinct plateau ages. These ages are distinct, so they cannot be explained by a single impact event, such as the one that produced the Imbrium impact basin. The conclusion that these represent distinct ages remains after considering the possibility of artifacts in defining plateaus.

Introduction

Samples of the Moon returned to Earth by the Apollo 11 astronauts showed that our satellite formed at nearly the same time as the Earth ($\sim 4.5 \times 10^9$ yrs ago) and that even its relatively young volcanic plains formed more than three billion years ago (3.7×10^9 years ago in the case of Mare Tranquilitatis) (LSAPT, 1970). Those initial analyses implied that the impact basins filled by the mare were also quite old, having formed between those two ages.

A series of geochronologic studies involving the Rb-Sr, K-Ar (using the Ar-Ar technique), and U-Pb radiometric systems revealed a series of impact-reset ages that clustered around 4.0 to 3.8 Ga (Papanastassiou and Wasserburg, 1971a,b; 1972a,b; Turner et al., 1973; Tera et al., 1974) in a pattern that has survived decades of additional scrutiny (e.g. Bogard, 1995; Merle et al., 2014; Norman and Nemchin, 2014; Norman et al., 2016). That distribution of ages was interpreted to represent a terminal lunar cataclysm involving at least the latter third of the lunar basins (Tera et al., 1974) and, when integrated with the ages of impact events captured in meteorites, interpreted to mean the impact bombardment affected the entire inner solar system (e.g. Bogard, 1995; Kring and Cohen, 2002). The hypothesis has been challenged and tested in a variety of ways. Bottke and Norman (2017) have provided a thorough review of many aspects of the subject, but we will mention a few of the studies that are most applicable to our investigation.

Some investigators have worked to determine the ages of specific basin-forming events (e.g., Dalrymple and Ryder, 1993, 1996), arguing that a cataclysm that may have involved most if not all of the lunar basins in a window of time as short as 20 million years (Ryder, 1990). Haskin et al. (1998) performed studies on mafic Th-rich melt breccias (which they defined as 19-29 % $\text{TiO}_2 + \text{FeO} + \text{MgO}$, 14-22% Al_2O_3) and suggested that some Ar-Ar ages might represent various degrees of partial resetting by Imbrium, rather than a number of discrete impact events. As reviewed by Bottke and Norman (2017), assigning ages to specific basins still remains a significant challenge, with the possible exception of the youngest basin on the near side, Imbrium, whose age will be discussed below.

Other alternative hypotheses are represented by the idea that a steady decline, rather than an abrupt spike, in impact flux erased evidence of older events (through regolith gardening and by depositing heat, which is what is required for resetting the age in diffusion-based radiometric chronometers used to date lunar samples) and that the Apollo sample sites may have been re-surfaced by Imbrium ejecta, influencing radiometric ages (Hartung, 1974; Hartmann, 2003). Fernandes et al. (2013) argued that Ar-Ar ages of 4.2-4.3 Ga in some samples have been misinterpreted as igneous or metamorphic events. The existence of impacts at such a time does not preclude the possibility of a late heavy bombardment, but does raise questions about the relative flux of impacts on the Moon at various times. Even more serious questions are raised by U-Pb analyses in zircon grains, some of which (Grange et al., 2009, 2011) show evidence for impacts at 4.2-4.3 Ga, but not at ~ 3.9 Ga. Norman et al. (2016) also interpreted the noritic anorthosite 67955 to be from a differentiated melt sheet with a ~ 4.2 Ga ^{40}Ar - ^{39}Ar age in the Procellarum-KREEP Terrane. Mercer et al. (2015) and Crow et al. (2017) interpreted strong signals in Ar-Ar of Apollo 17 impact melt breccias at 3.81-3.83 Ga and in U-Pb ages of lunar zircons at 3.9-4.1 Ga as signatures of Imbrium, once again providing ages in the range of interest,

but not settling the question of whether Imbrium was the only contributor to the apparent ages in that range. Zellner (2017), too, has suggested the duration of lunar bombardment extends from 3.5 to 4.2 Ga, based on a collation of sample and remote sensing data with dynamic models of impactor fluences.

Other details of the collisions that occurred during the basin-forming epoch have been surfacing in parallel with these geochronologic debates. Geochemical (Ganapathy et al., 1972; James, 1994, 1995, 1996; Korotev, 1994; Norman et al., 2001; Kring and Cohen, 2002; Puchtel et al., 2008) and mineralogical (Joy et al., 2012) remnants of the projectiles imply the impactors were dominated by asteroids. The size distribution of the craters also points to a population of impactors dominated by asteroids (Strom et al., 2005) that was sampled in a size-independent way (although a recent study by Orgel et al., 2018, concluded that the size distribution was not distinct in the basin-forming epoch). That suggests resonances swept through the asteroid belt when Jupiter's orbit shifted, an idea that has been explored in a series of dynamical studies (Gomes et al., 2005; Morbidelli et al., 2005; Morbidelli, 2010). It also appears that the oldest basin on the Moon, the South Pole-Aitken Basin, may have formed from a different population of objects during the accretional epoch of the Earth-Moon system (Marchi et al., 2012) prior to ~4.3 Ga (Morbidelli et al., 2012). Thus, any late heavy bombardment began after that time.

Boehnke and Harrison (2016) have taken a different approach, producing a model that yields an increase Ar-Ar ages at ~4 Ga despite a steadily decreasing flux in impactors with time. However, that result is a consequence of the assumption that the lunar crust would have been too hot to retain radiogenic Ar until several hundred Ma after the Moon's formation. Such a scenario would explain the increase in lunar ages once the lunar crust began retaining Ar, and would have important implications for the history of the Moon's surface. In fact, Bottke and Norman (2016) suggest that the thermal evolution of the lunar crust could affect the available Ar-Ar results, but only by explaining the paucity of lunar melt rocks >4.2 Ga. Keeping the lunar crust too hot to retain Ar until ~4 Ga would seem to be at odds with arguments mentioned above that there is a significant number of lunar Ar-Ar ages at 4.2-4.3 Ga (Fernandes et al., 2013; Norman et al., 2016). This model would also fail to explain asteroids which produce meteorites that have higher numbers of Ar-Ar ages at ~4 Ga than before or after (Bogard and Garrison, 2003; Swindle et al., 2009), despite there being no obvious reason why such small bodies would have such hot crusts so late in their histories.

To continue probing the timing of those events, we still need to determine if the population of highland impact ages c. 3.9 Ga from the Apollo 16 landing site reflects the overprint of a single impact event, the Imbrium basin impact event (an artificial age spike due to sampling bias), or if evidence of multiple impacts, consistent with the late heavy bombardment hypothesis, is represented in Apollo samples. Norman et al. (2006) was the first to address this issue by examining a suite of group 1, 2, 3, and 4 impact melts (using the classification scheme of Korotev, 1994) that have both a range of textures and a range of Ar-Ar ages. Among 25 whole-rock samples, they found ages from 3.75 to 3.96 Ga and suggested those ages were produced by at least four different impact events at 3.771, 3.836, 3.865, and 3.859 Ga. If it can be confirmed, it implies the cluster of ages between 4.0 and 3.8 Ga was not produced by a single (Imbrium) impact event, but reflects a large number of impact events that occurred in a short duration of time. Fernandes et al. (2013) obtained Ar-Ar ages between 4.3-3.3 Ga for 24 rock fragments

(1-3 mm sized) from an Apollo 16 soil sample, plus chips from three Apollo 16 and 17 samples which include crustal rocks, breccias, and impact melt rocks and suggested that the younger resetting events have occurred at 4.0, 3.9, 3.6, and 3.3 Ga, together with 4.3 and 4.2 Ga peaks. Merle et al. (2014) obtained high-precision U-Pb ages of 3938 Ma for Ca-phosphate in Apollo14 breccias and concluded that several impacts occurred during a 20-30 Ma period around 3900 Ma by comparison with zircon ages from Apollo12 and lunar meteorite samples. Later, Snape et al. (2016) also argued for multiple impacts in that period, based on U-Pb analyses of phosphate grains from Apollo 14.

An additional complication has been that several data sets suggest a late bombardment, but with a much broader range in ages. HED meteorites (Bogard and Garrison, 2003), H chondrites (Swindle et al., 2009), and L chondrites (Swindle et al., 2014) all show increases in the number of impact ages between ~ 4.1 Ga and ~ 3.4 Ga. This difference from the Apollo samples could plausibly be attributed to the asteroidal meteorites' different locations. More significantly, a suite of recrystallized impact melts from lunar meteorites (Cohen et al., 2000, 2005) has ages ranging from ~ 3.9 Ga to ~ 3 Ga, though with rather large uncertainties. It is not clear whether the difference between the Apollo samples and the lunar meteorite impact melts is a result of the different locations (the chemistry of the lunar meteorites studied shows that they are not from the central near side) or the different sizes (the Apollo samples studied have been primarily hand samples, while the lunar meteorite samples were less than a millimeter in length). Finally, it is possible that the differences simply represent the sizes of impacts involved. Whereas the lunar record is dominated by basin-size impact events, comparable collisions among asteroids would completely disrupt the targets. Thus, meteoritic melts we analyze from asteroids are representative of smaller events, which would persist after large impacts had stopped.

At a more fundamental level, as the Ar-Ar technique has matured, questions have been raised about the criteria that have been used to pull ages out of the data (e.g., Jourdan, 2012; Jourdan et al., 2007). In particular, the technique generates a series of apparent ages from extractions at different temperatures (as in this study) or places within a rock (as in Mercer et al., 2015). It is critical to understand how those apparent ages relate to the timing of actual events. In particular, it is critical to understand how partial resetting (e.g., the Imbrium scenario described above), would be reflected in the data. We will see that there may be more useful information than has sometimes been recognized.

With all this in mind, we selected additional impact melt samples from the Apollo 16 collection of an intermediate size (centimeter-sized) in an ancient regolith breccia. We petrographically characterized them to discern any similarities and differences that may occur among them, and performed Ar-Ar experiments on several splits of each sample to determine if they formed in a single event or in a series of discrete impacts. This is complementary to the study done by Norman et al. (2006), who studied single homogenized splits from large scale (> 5 cm) rock samples.

We also modeled the diffusive release of argon under a variety of conditions to understand the likely magnitude of analytical artifacts in apparent plateaus in the Ar-Ar system. In particular, we explored some of the situations that might arise when analyzing samples with multiple diffusion domains, as can be the case for "whole rock" samples such as those we analyzed (see also Swindle and Weirich, 2017).

Samples and their geologic context

The Apollo 16 landing site is within the ancient cratered highlands of the Moon. Photogeologically, it appears to contain a significant amount of impact ejecta from the Imbrium basin and potentially from the Nectaris basin. The site has two geologic formations (Hodges et al., 1973; Wilhelms et al., 1987; Ulrich, 1973; Fig. 1). The Cayley Formation extends from the northwest towards the southeast and represents a distal facies of Imbrium ejecta. It is superimposed on the Descartes Formation, which has at times been interpreted to be either another facies of the Imbrium ejecta (Hodges and Muehlberger, 1981) or potentially Nectaris ejecta (Spudis and Pieters, 1991). The Apollo 16 regolith is dominated by feldspathic material (Korotev, 1997), what would now be called FHT (Feldspathic Highlands Terrane) composition, but because of the low concentrations of radioactive elements such as K and U, the age of this material is difficult to determine.

We have focused our attention on ancient regolith breccia 60016. This sample is an immature regolith breccia that does not contain very many agglutinates (McKay et al., 1986). It was produced during the end of the basin-forming epoch and lithified, and, thus, closed to contributions from later events, about 3.74 Ga (Joy et al., 2011). We selected six centimeter-size impact melt clasts from the breccia. The melt clasts contain relict rock or mineral fragments, which can be used to evaluate the types of lithologies involved in each of their impact events. Petrological analyses of the impact melt can also be used to infer the types of rocks in the impact source region. We performed Ar-Ar analyses to determine the age(s) of the impact event(s) that produced them and evaluate any later geologic disturbances of that radiometric system. Collectively, the petrologic and Ar-Ar analyses of intermediate sized melt breccias are used to determine if they were produced by a single (Imbrium) impact event or several impact events that occurred at nearly the same time.

Analytical procedure

Petrological analyses

We were allocated six 1 to 2 cm-sized dark clasts that were removed from the light gray matrix of 60016: petrological thin sections (Fig. S1) of clasts we used are Clast 1 (,319), Clast 2 (,320), Clast 3 (,324), Clast 4 (,322), Clast 5 (,323), and Clast 6 (,321). We utilized an optical microscope, backscattered-electron (BSE) detectors on a Scanning Electron Microscope (SEM; JEOL JSM-5910LV), and a Field Emission Gun-SEM (JEOL JSM-7600F). Major element analyses were made with an Electron Probe Micro Analyzer (EPMA; CAMECA SX-100) using a 15 kV accelerating voltage, 20 nA beam current, and focused $<1\text{ }\mu\text{m}$ beam diameter for silicates, oxides, metal, sulfide, and phosphide analyses. Defocused beam (20 μm diameter) analyses (DBA) of representative areas for each clast were performed to determine bulk major element compositions.

Peak positions and intensities of individual elements were calibrated and monitored for silicates and glass using the following standards: Si with diopside, forsterite, and orthoclase; Al with oligoclase; K with orthoclase; Ca with diopside and feldspar (An_{74}); P with apatite; Ti with rutile; Cr with chromite; Fe with fayalite and hypersthene; Ni with NiO; Mn with rhodonite; Na with oligoclase; and Mg with forsterite and diopside. Elemental abundances in spinel, ilmenite, and armalcolite were analyzed with the same standards except Al, Fe, and Mg, which were calibrated with chromite. Metal, sulfide, and phosphide analyses were performed using the following standards: Ti, Fe, Zn, Cu, Cr, Mn, Co, Ni each with their own pure metal; S with troilite; Cl with tugtupite; P with apatite; Ca with apatite; Si and Mg with diopside. Interference of Fe K β on Co K α was corrected. Potential contamination by silicate inclusions was monitored with the Si and Mg signals. Major, minor, and trace elements of phosphates were analyzed using 15 kV accelerating voltage, 10 nA beam current, and 5 μm defocused beam using the following standards: Na with oligoclase; Si with topaz; Al with oligoclase; Mg with kaersutite; Y with YPO_4 ; Cl with tugtupite; Ca with apatite; Ti with rutile; P with apatite; S with apatite; Fe with kaersutite; La with LaPO_3 ; Ce with CePO_3 ; Sm with SmPO_3 ; and F with topaz (Stormer et al., 1993). Fluorine is calculated by the time zero intercept method (McCubbin et al., 2011). Rare earth element abundances were estimated according to McCubbin et al. (2011). Representative analysis data is shown in Table S1 and S2.

The compositions derived from DBA may show systematic difference in several elements due to density effects among minerals during EPMA analyses (Warren, 1997). Dowty et al. (1973) suggested the following correction factors for mare basalt: CaO, 0.93; FeO, 1.11; Al_2O_3 , 0.77; MgO, 1.06; and TiO_2 , 1.29. Warren (1997) obtained similar values from his method. We tested our DBA procedure with a certified terrestrial geostandard (BCR-2), which is a fine-grained basalt. We analyzed the standard in 5 different areas that seem to have slightly heterogeneous abundances of constituent materials (Table S3). Average oxides values are systematically higher in Al_2O_3 , MgO and P_2O_5 , and lower in FeO and K_2O than certified standard composition values (Table S4). We then calculated bulk compositions using Warren's correction scheme. The corrected values are consistent with certified values, although the observed heterogeneity among the 5 analyzed areas is evident (Table S4). The test indicates DBA can provide meaningful bulk major and minor element compositions of fine-grained impact melt clasts from 60016, although they may have small uncertainties.

Ar-Ar analyses

Samples from impact melt clasts located immediately adjacent to the petrological samples were split for Ar-Ar analyses: Clast 1 (,286; 49.7 mg), Clast 2 (,290; 48.6 mg), Clast 3 (,292; 43 mg), Clast 4 (,305 and ,314; 8 mg and 15 mg), Clast 5 (,299; 54 mg), and Clast 6 (,312; 35 mg).

Sample clasts for Ar-Ar analysis were divided into one to three splits, depending on the original clast size. The splits were irradiated at the Cadmium-Lined In-Core Irradiation Tube (CLICIT) facility at Oregon State University along with PP-20 Hb3gr Hornblende GSC as the primary irradiation monitor (1080.4 ± 1.1 Ma, Renne et al., 2010) to determine the J-factor; roughly 2×10^{-2} among our samples. A significant advantage of using the CLICIT facility is that the Cd lining significantly reduces the number of thermal neutrons, which in turn significantly reduces the production of ^{38}Ar by neutron capture on ^{37}Cl . Since these samples are volatile-poor (hence likely Cl-poor), this means that ^{36}Ar and ^{38}Ar are basically mixtures of only trapped (solar wind) Ar and Ar produced by cosmic-ray spallation, so the $^{36}\text{Ar}/^{38}\text{Ar}$ ratio can be used to accurately correct for spallation, making it possible to make much more accurate corrections for trapped Ar.

Samples were irradiated for 300 hrs in the linear-with-height portion of the reactor to provide a more evenly distributed neutron flux. Samples were then allowed to cool for a minimum of 10 weeks to allow for short-lived isotopes to decay before being placed in a glass storage tree above a double-vacuum, resistance-heated furnace. Despite the wait, the levels of radioactive ^{37}Ar were higher than we would normally admit into the mass spectrometer, so we sampled ^{37}Ar for two seconds per cycle for seven cycles, rather than the eight seconds per cycle for 10 cycles that we normally use. Instrumental backgrounds were monitored, but were not an issue. Argon was extracted from each sample using a computer controlled step-heating procedure. Typical temperature steps were from 300-1500 °C in 50 °C intervals, totaling 25 heating steps per split. The furnace temperature has an estimated accuracy of ± 5 °C, based on melting of pure metals. Each step was run for duration of 12 minutes and, after gettering, analyzed in a VG 5400 mass spectrometer. The computer program Mass Spec *Ver. 7.884* (Deino, 2001) was used for initial data collection and analysis, supplemented by software written by J. R. Weirich. The decay constants used are those reported in Renne et al. (2010). Use of the decay constants given by Steiger and Jäger (1977) would yield apparent ages that are approximately 10 Ma younger. Using the slight revision of Renne et al. (2011) would change the reported ages here by 6 Ma or less. Isotopic measurements were regressed to a time of zero using linear regression techniques, after subtracting baseline values in order to get the most accurate measurements. Uncertainties and corrections were applied to account for blanks, machine discrimination, spallation-produced isotopes, and interfering isotopes produced in the reactor from Ca and K. Isochron plots were evaluated to search for evidence of either terrestrial argon or Ar trapped on the Moon (which is a combination of ^{36}Ar and ^{38}Ar from the solar wind and ^{40}Ar that has been outgassed from the Moon, photoionized, and reimplanted). In most cases, no convincing isochrons were seen, in part because ratios of ^{36}Ar to ^{40}Ar were usually very low at this stage. Unless otherwise

stated, a trapped $^{40}\text{Ar}/^{36}\text{Ar}$ of 8 ± 8 was used, to span the values expected for either the modern (~ 1) or ancient (~ 15) lunar surface (Eugster et al., 1999).

Sample ages were determined based on ^{40}Ar - ^{39}Ar apparent age spectra (“plateau plots”). We follow the definition of a plateau as a region that contains at least 50 % of the ^{39}Ar and at least three consecutive temperature steps that agree at a 95 % confidence level (Fleck et al., 1977). It has been suggested that the definition of a plateau should require 70% of the ^{39}Ar (e.g., Jourdan, 2012), but Swindle and Weirich (2017) have shown that for a single-phase system that has undergone diffusive loss, a 50% plateau should be $\sim 2\sigma$ lower than the true age, while a 70% plateau should be $\sim 0.8\sigma$ lower than the true age, so even a 50% plateau should be meaningful (albeit slightly low). Furthermore, since we are analyzing multiminerallic systems, with multiple potassium sites with different diffusion parameters, those distinct minerals (or types of diffusion domains) may degas over distinct temperature ranges, so if a plateau includes only Ar from one mineral, 50% of the entire sample may represent $>50\%$ of the Ar from the degassing mineral. We will discuss these complications, and how we may disentangle some of them, below. Plateau ages reported are at the 2σ level and are calculated using the mean of all the steps in the plateau region, weighted by the amount of gas contained in each step. In some cases where there is a partial plateau, incorporating a smaller fraction of the ^{39}Ar , we will quote these as our best estimate of the age of the sample, but we will also evaluate the degree to which the apparent age of a partial plateau might differ from the actual age of degassing.

Results

Petrography and mineral compositions

Clast 1 (,319), a *clast-rich poikilitic melt*, contains ~45 modal % relict mineral fragments of plagioclase ($\text{An}_{97-95}\text{Or}_{0.6-0}$, $n=44$; 39 modal %), olivine (Fo_{91-69} , $n=59$; 5 modal %), pyroxene ($\text{En}_{77-58}\text{Wo}_{5.2-2.5}$, $\text{Mg\#}$ (mol% $\text{Mg}/[\text{Mg}+\text{Fe}]) = 79-59$, $n=21$; 1 modal %) and a small amount ($\ll 1\%$) of ilmenite and Mg,Al-spinel (Fig. 2a, b). Relict olivine grains (~300 μm) have a rounded shape indicating that these are not crystallization products from surrounding impact melts, and are completely or partly enclosed by melt products of fine-grained pyroxene and plagioclase grains showing poikilitic texture. Some olivine relict grains retain chemical zoning ($\text{Fo}_{89.8}$ at core, $\text{Fo}_{78.7}$ at rim). The relict olivine has a bimodal composition, $\text{Fo}_{91.0-77.9}$ and $\text{Fo}_{75.3-69.0}$. Relict pyroxene grains (~200 μm) usually show a corroded texture at the rim where the grains are in contact with the surrounding melt. Relict pyroxene also has a bimodal composition of $\text{En}_{77-72}\text{Wo}_{3.5-2.5}$ and $\text{En}_{69-58}\text{Wo}_{5.2-2.5}$, which is not in equilibrium with the olivine. Plagioclase relicts (~300 μm) are subhedral, anhedral to irregular morphology with slightly Na-rich ~2 μm -wide overgrowths (Na_2O contents are 0.4 wt. % at core and 1.2 wt. % at rim) with small Fe-Ni droplets ($<1\ \mu\text{m}$ in diameter). These plagioclase grains show undulatory extinction (up to 15°), but no evidence of maskelynitization.

Interstitial material consists of fine-grained pyroxene ($\text{En}_{82-46}\text{Wo}_{4.2-3.3}$, $\text{Mg\#}=85-76$; 25 modal %), plagioclase ($\text{An}_{95-83}\text{Or}_{1.7-0.3}$; 23 modal %), Fe-Ni metal ($\text{Ni}= \sim 6$ wt. %, $\text{Co}= \sim 0.4$ wt.%; 3.0 modal %), troilite, ilmenite, and glass. Pyroxene grains consist of high-Ca augite and low-Ca pigeonite, and enclose fine-grained euhedral to subhedral plagioclase and rounded relict olivine. Fe-Ni metal droplets (~44 μm) are fine-grained aggregates of few-micron-sized metal with interstitial P-bearing material ($\ll 1\ \mu\text{m}$; probably schreibersite; Fig. 3).

Clast 2 (,320), a *clast-poor poikilitic melt*, contains ~15 modal % relict grains of olivine ($\text{Fo}_{79.6-73.2}$, $n=30$; 12 modal %), plagioclase ($\text{An}_{97-93}\text{Or}_{0.4-0}$, $n=25$; 2 modal %), and pyroxene ($\text{En}_{84-78}\text{Wo}_{3.9-2.8}$, $\text{Mg\#}=86-80$, $n=19$; 1 modal %) (Fig. 2c, d). Relict olivine (~100 μm) and pyroxene grains are irregular to round in shape surrounded by pyroxene which crystallized from impact melt showing poikilitic texture. The composition of olivine falls into two groups of $\text{Fo}_{79.6-78.4}$ and $\text{Fo}_{77.0-73.2}$. Relict plagioclase grains (~200 μm) have subhedral to anhedral morphologies with overgrowth rims slightly enriched in Na (Na_2O contents are 0.4 wt. % at core and 1.3 wt. % at rim). Some relict plagioclase grains have undulatory extinction. The Na-enriched rims contain small Fe-Ni droplets ($<1\ \mu\text{m}$ in diameter). This clast has up to 2 mm pyroxene crystals ($\text{En}_{83-49}\text{Wo}_{4.1-2.9}$, $\text{Mg\#}=85-77$; 38 modal %) that enclose fine-grained euhedral plagioclase ($\text{An}_{96-84}\text{Or}_{1.3-0.2}$; 43 modal %) and are embedded in an olivine- and feldspar-bearing mesostasis. Opaque phases in this clast are Fe-Ni metal ($\text{Ni}= \sim 7$ wt.%, $\text{Co}= \sim 0.5$ wt.%; 2.1 modal %), troilite, and armalcolite. Fe-Ni metal droplets (~180 μm) have eutectic textures with schreibersite ($\text{Ni}=18$ wt.%) and are surrounded by rims of troilite.

Clast 3 (,324), a *clast-rich poikilitic melt*, contains ~40 modal % relict grains of plagioclase ($\text{An}_{97-82}\text{Or}_{2.6-0.1}$, $n=67$; 29 modal %), olivine ($\text{Fo}_{82.8-73.5}$, $n=32$; 8 modal %), and pyroxene ($\text{En}_{78-75}\text{Wo}_{4.8-4.0}$, $\text{Mg\#}=81-78$, $n=4$; 3 modal %) (Fig. 2e, f). Relict plagioclase (~5 μm) rims are, again, slightly enriched in Na (Na_2O contents are 0.4 wt. % at core and 1.2 wt. % at rim). The compositions of relict olivine grains (~100 μm) are clustered ($\text{Fo}_{78.7-73.5}$) and one grain has slightly higher Mg content ($\text{Fo}_{82.8}$). Clast 3 has up

to 1 mm pyroxene crystals ($\text{En}_{82-50}\text{Wo}_{40-3.0}$, $\text{Mg}\# = 85-79$; 36 modal %) enclosing euhedral plagioclase ($\text{An}_{97-88}\text{Or}_{0.9-0.3}$; 23 modal %) and relict olivine grains showing poikilitic texture similar to Clast 2. The pyroxene is embedded in an olivine- and feldspar-bearing mesostasis. Other constituent materials are Fe-Ni metal ($\text{Ni} = \sim 5$ wt.%, $\text{Co} = \sim 0.4$ wt.%; 0.4 modal %), troilite, armalcolite, whitlockite, and glass. Fe-Ni metal droplets (~ 80 μm) are fine-grained aggregates, but do not contain schreibersite or phosphorous materials. Whitlockite grains usually occur with Fe-Ni metal.

Clast 4 (322), a *clast-rich melt*, is petrologically heterogeneous with two types of melt. The bulk of the thin-section has an aphanitic texture with ~ 47 modal % relict grains of pyroxene (~ 50 μm ; $\text{En}_{81-78}\text{Wo}_{5.1-3.7}$, $\text{Mg}\# = 83-81$, $n = 34$; 29 modal %), plagioclase (~ 110 μm ; $\text{An}_{96-88}\text{Or}_{1.4-0.1}$, $n = 20$; 10 modal %) and olivine (~ 50 μm ; $\text{Fo}_{79.1-77.8}$, $n = 36$; 8 modal %), which show irregular to rounded morphologies (Fig. 2g, h). Matrix is filled with fine-grained (~ 10 μm) melt products of plagioclase ($\text{An}_{96-85}\text{Or}_{1.9-0.3}$; 17 modal %), Fe-Ni metal ($\text{Ni} = \sim 8.9$ wt.%, $\text{Co} = \sim 0.4$ wt.%; 3.5 modal %), troilite, and interstitial pyroxene. Fe-Ni droplets (~ 136 μm) have P-bearing materials (< 1 μm) in their mesostasis. Fine-grained shock-disseminated Fe-Ni metal droplets (< 1 μm) are responsible for the appearance of dark bands across the thin section dominating different types of melt. Adjacent to one of those bands is a second melt with an intergranular texture. Its matrix is coarser-grained (~ 20 μm) and consists of plagioclase and intergranular pyroxene. In this portion of the sample, few metal particles are found. This melt fraction can be distinguished from the aphanitic melt by both its grain size and abundances of metal droplets. Mineral compositions of relict and melt products in this intergranular melt are, however, the same as in the other melt.

Clast 5 (323), a *clast-rich intergranular glassy melt*, contains ~ 34 modal % relict grains of plagioclase (~ 150 μm ; $\text{An}_{97-82}\text{Or}_{0.4-0}$, $n = 47$; 34 modal %) and a trace of olivine (~ 100 μm ; $\text{Fo}_{90.8-93.7}$, $n = 25$; less than 1 modal %) which have an irregular to rounded shape (Fig. 2i,j). Plagioclase relicts have a thin overgrowth (< 1 μm). Matrix is composed of melt products of plagioclase ($\text{An}_{97-93}\text{Or}_{0.5-0}$; 24 modal %), Fe-Ni metal droplets, sulfide, and interstitial glass. Fe-Ni metal droplets (~ 68 μm) are fine-grained aggregates of kamacite ($\text{Ni} = \sim 4$ wt.%, $\text{Co} = \sim 0.5$ wt.%; 1.5 modal %) and schreibersite ($\text{Ni} = 10$ wt.%, with troilite rims (Fig. 3).

Clast 6 (321), a *clast-poor porphyritic melt*, contains ~ 15 modal % relatively large-sized grains of anhedral olivine (~ 50 μm ; $\text{Fo}_{86.4-80.7}$, $n = 28$; ~ 9 modal %) and plagioclase (~ 50 μm ; $\text{An}_{98-90}\text{Or}_{1.0-0}$, $n = 85$; 6 modal %) with groundmass (Fig. 2k,l). Some plagioclase grains have slightly Na-rich overgrowths (Na_2O contents are 0.4 wt. % at core and 0.7 wt. % at rim). Minor clots of phenocrysts produce a subtle glomeroporphyritic texture. The fine-grained matrix is composed of euhedral olivine ($\text{Fo}_{86.1-82.0}$; 9 modal %), plagioclase ($\text{An}_{96-90}\text{Or}_{0.6-0.1}$; 25 modal %), interstitial dendritic pyroxene, Fe-Ni metal droplets, troilite, schreibersite, and glass. Most metal droplets (4 to 160 μm in diameter) have rounded shapes and are a fine-grained mixture of Fe-Ni metal ($\text{Ni} = \sim 5$ wt.%, $\text{Co} = \sim 0.5$ wt.%; 2.2 modal %) and schreibersite ($\text{Ni} = \sim 11$ wt.% up to 16 μm in size) with rims of troilite.

Major and minor element compositions of the six impact melt clasts were estimated from the averaged values of defocused beam analyses (DBA) by EPMA (Table. 1).

Averaged DBA-derived compositions for the melt clasts in 60016 have a narrow range of K_2O (0.3-0.5 wt.%) and P_2O_5 (0.4-0.5 wt.%) concentrations. On the other hand,

Mg# (69-78 mol.%), FeO+MgO (10-23 wt.%) and Al₂O₃ (16-26 wt.%) concentrations vary widely. Clast 1 and Clast 5 have the lowest values of MgO+FeO (< 12 wt. %).

Ar-Ar ages

All apparent age spectra, or “plateau plots,” are disturbed in some way, typically with diffusive loss in low temperatures steps resulting in the lowest apparent ages at the lowest extraction temperatures (Fig. 4). However, Clasts 5 and 6 have a few steps with distinct K/Ca at low temperature steps and higher apparent ages. Most samples yield consistent ages over ~50% of the Ar release or more, but then have evidence of lower or higher ages at higher extraction temperatures that could be released from a different mineral phase, often with a different K/Ca. The amount of ³⁹Ar released in the plateau or that we use to determine a best age of a sample, is stated as the range released in the region of interest among all of the splits. Because each sample shows evidence of multiple mineral phases (based on K/Ca and release characteristics), quoted plateau widths (% of Ar released) represent a minimum width for that mineral phase. Including these effects generally increases the plateau width ~10-20%, with respect to the mineral phase from which it was released. This is not discussed in detail because it is not clear how to properly account for these effects for every sample split and therefore we only report the minimum width. Lower ages at high extraction temperatures can be the result of either recoil of ³⁹Ar during neutron irradiation at high temperatures or degassing of domains with distinct diffusion parameters (e.g., different mineral phases). For most samples (with some exceptions discussed below), we conclude the data are very good estimates of the age of a significant thermal event. These ages are summarized in a probability plot in Fig. 5.

The minimum ages reported are the range of the lowest ages from all splits analyzed for each sample (Table 2 and S5). They are presented for a comparison to the work done by Shuster et al. (2010), who point out that the minimum apparent ages may contain useful information about the overall impact history beyond that provided by the plateau ages alone. We will refer to these as “minimum ages”, even though they are defined by the minimum *apparent* ages and actually represent an upper limit (or maximum) to the time of the partial degassing event that produced them. We will discuss their full implications, and potential utility, in the Discussion.

The apparent age spectra, “plateau plots,” for each of the three splits of Clast 1 (,286; 49.7 mg) look very similar (Fig. 4). All three splits give a series of concordant ages through 40 to 55% of the ³⁹Ar released (600-900 °C temperature steps). Split SB16 does not meet our preferred definition of a “plateau” as having more than 50%, but another 20% of the ³⁹Ar released is consistent with an age only 10 Ma higher. The ‘best ages’ determined from all three splits average to 3877.5 ± 13.8 Ma, our best estimate of the age of this sample. Minimum ages range from 1756 to 3410 Ma.

We have more confidence in the age of Clast 2 (,290a; 48.6 mg) than that of most of our other results (Fig. 4). Plateau ages (550-850 °C temperature steps, ranging from 53.3 to 67.9% of the total ³⁹Ar in the samples) agree within error for two splits and very nearly agree for the third (3972.5 ± 4.1 , 3980.5 ± 8.3 , and 3957.3 ± 4.9 Ma). Averaging all three together gives our best estimate for the age of this sample at 3970.1 ± 12.8 Ma, which agrees with all three splits within error. Each split also agrees with the age determined by the standard isochron, giving us even more confidence in the plateau age. Because the

plateau ages (which often contain more than 70% of the gas from their host mineral) and standard isochron ages are in agreement, we feel confident in our results for Clast 2. The minimum ages range from 2170 to 2577 Ma.

For impact melt Clast 3 (,292; 43 mg), we analyzed 3 splits, using a trapped $^{40}\text{Ar}/^{36}\text{Ar}$ correction of 8 ± 2 , which was obtained from its reverse isochron. All three splits have concordant ages through 24-35 % of the ^{39}Ar released (Fig. 4). Although no sample meets the requirement of a plateau age, the closest thing (temperature steps varying from 600-900 °C), gives ages ranging from 3978.8 ± 12 to 4023 ± 6 Ma with an average age of 4000.0 ± 23.2 Ma, our best age estimate. The K/Ca ratios suggest that there may be separate minerals degassing at both higher and lower temperatures than those of the “plateaus”, and even the intermediate temperatures where the apparent ages are closest to concordant yield a pattern that is clearly disturbed. The minimum ages of the three splits range from 1500 to 2275 Ma. Finally, we note that after the end of the partial plateau at high temperature steps (starting at 1150 °C or later), there is another set of consecutive steps that give similar ages to each other, yet result in a lower age than that given by previous, lower temperature steps (3830.1 ± 24.61 Ma compared to 4000.0 ± 23.2 Ma). We interpret this as argon being released by another mineral phase or diffusion domain with both a higher activation energy and a higher pre-exponential factor (Cassata et al., 2011)

For Clast 4, we had only one split of ,305 (8 mg) and two of ,314 (15 mg) because of sample size. The apparent age plots of Clast 4 appear to be very disturbed, having an oscillating pattern after an initial region of diffusive loss (Fig. 4). Not only are the apparent age plots disturbed, they differ significantly more in both appearance and age between splits than our other samples do. The closest thing to a plateau for the ,305 sample, 3866.0 ± 3.9 Ma, comes from a ‘flat’ region in steps 1250-1350 °C, which contains 26.2 % of the ^{39}Ar released. If this were part of a diffusion pattern, the fraction of the ^{39}Ar contained is small enough that the age of the “plateau” would be expected to be significantly (several sigma) lower than the actual age of the sample (Swindle and Weirich, 2017). The K/Ca data for each split shows evidence for a distinct mineral phase at temperatures >1200 °C (above ~65% ^{39}Ar release), which would increase the relative amount of argon (plateau width) released from the high temperature phase to ~65%, making it a true plateau. Summing all of the argon, which might be a better determination of the age of the sample due to its disturbed age spectra, and is certainly a lower limit to the age, is 3822.2 ± 6.8 Ma, with a minimum of 2393.7 ± 23.2 Ma. The two splits from the other sample from Clast 4 (,314; 15 mg) yield disturbed plateau plots reminiscent of ,305, but different in detail. As was done in ,305, the best age is from a flat region near the end of the temperature steps (~1200-1450 °C, with 19.7-24.9 % ^{39}Ar), this time giving an age of 3787.2 ± 16.0 Ma. Similar arguments to those presented for ,305 apply – it could represent a true plateau, but could represent significant loss from a disturbed system. The overall summed age of the two ,314 splits is 3748.35 ± 15.7 Ma. The minimum age ranges from 2852 to 3427 Ma. Given the uncertainties, our best estimate of the age of Clast 4 would be 3823 ± 75 Ma, incorporating all of the plausible ages. Given the variety of disturbed behaviors, and the small sizes of the “plateaus”, even that may underestimate the uncertainty. In particular, diffusive loss would lead to a small “plateau” with an “age” that is younger than the true age, and since this estimate is at the

young end of our age distribution, we will not use this age as a part of our ultimate analysis.

The three splits of Clast 5 (,299; 54mg) are in good agreement with each other, even though there was not a plateau from every split. Each of the three apparent age plots shows the same trend (Fig. 4). There are very high apparent ages in the low temperature steps (300-650 °C), presumably a result of terrestrial contamination. Two of the three splits have a plateau, with the third not quite meeting our definition (agreement is within 3 sigma instead of 2). The plateaus of two splits (1150-1500 °C temperature steps, containing 51.3 to 56.0 % of ^{39}Ar released) agree within error to 3922.9 ± 3.7 Ma. If we include the 3 sigma range of the third split (1100-1400 °C steps, 55 % ^{39}Ar), our best age for this sample is 3926.5 ± 7.3 Ma. Another testament that shows how well these three splits agree is shown by looking at the entire summed age (minus the air contaminated steps, and containing over 90 % of the ^{39}Ar), which is 3913.1 ± 3 Ma for all three splits (3916.1 , 3912.0 , 3911.3 ± 5.3 Ma for each split), only slightly lower than the plateau ages. The minimum ages are only ~ 3700 Ma, so there is little evidence for any later diffusive loss. Because of the good agreement in ages for every split of this sample, we are quite confident in the results. Our best estimate for the age of Clast 5 is 3926 ± 7.3 Ma.

Splits of Clast 6 (,312; 35mg) also have apparent age spectra that agree very well with one another (Fig. 4). Over 25 % of the argon was contained in just one step at 1300 °C in two of the splits, so for the third split, we changed the number of temperature steps from 25 to 33 to help clarify the age represented by this region. The first two splits (25 heating steps) have plateaus (1200-1450 °C temperature steps, with 58.4-59.5 % of released ^{39}Ar) that agree within error at 3906.2 ± 2.2 Ma. The flat portion (1220-1350 °C steps) of the third split (33 heating steps) does not quite contain 50 % of the ^{39}Ar , but gives an age of 3906.6 ± 3.2 Ma, agreeing with the other splits. In all three, the plateaus appear to represent at least 60% of the gas of the primary K host phase, so we suspect they are all meaningful. Therefore we include the results of this split in our final best estimation of 3906.3 ± 2.4 Ma. If we include all the gas and sum up the age, they agree within error to 3885.7 ± 10.7 Ma, only slightly younger than the previous result. The minimum step ages range from 2846-3369 Ma, but have large errors (50-133 Ma). Our best estimate for the age of Clast 6 is 3906.2 ± 2.2 Ma.

Discussion

Origins of the centimeter-sized impact melt clasts

Clasts 1, 2, and 3 have similar poikilitic textures (Fig. 2). Moreover, Clasts 2 and 3 both have poikilitic pyroxene enclosing plagioclase and are embedded in a relict olivine- and plagioclase-bearing mesostasis, similar to the textures of the Apollo 16 mafic poikilitic group reported by Norman et al. (2006). However, the constituent relict materials and their abundances differ between all clasts. Clasts 4, 5 and 6 do not have poikilitic textures but have intergranular, aphanitic or porphyritic textures (Fig. 2). These differences in texture and abundance of relict materials imply the clasts have different post-shock conditions (e.g. post-shock temperature, cooling condition).

Relict minerals can be used to infer pre-impact target materials as long as they are not severely affected by shock-metamorphic overprints. Relict plagioclase laths in melt clasts do not have evidence for transformation into maskelynite, nor do olivine and pyroxene have any evidence of mosaicism or planar deformation features, implying the grains are reliable indicators of target sequences.

All six clasts contain different amounts of relict materials, plagioclase, pyroxene and/or olivine, which have irregular to rounded shapes surrounded by melt products.

Abundances of relict materials differ among the clasts; Clasts 1, 3, 4 and 5 contain higher abundance (~34-50 %) of relict materials, while the other two melts contain less than 15%. The different amounts of relict grains might reflect a difference in post shock conditions such as post shock temperature and degree of melting of precursor materials.

The cores of relict plagioclase grains with Na-enriched overgrowth rims have not equilibrated with the melt (Meyer et al., 1974; Thornber and Huebner, 1980). This is similar to a finding by Meyer et al., (1974) who analyzed major, minor, and trace element compositions in zoned plagioclase (50- 500 μm) in Apollo 16 poikilitic rocks (Simonds et al., 1973; Albee et al., 1973). Not only were the cores distinct from their rims, they were also different from the compositions of fine-grained euhedral plagioclase grains crystallized from the melt matrix and indicated relict grain cores that preserved target compositions of a KREEPy materials (Hubbard et al., 1973). Compositions of relict plagioclase are within the range of plagioclase in Mg-suite rocks and ferroan anorthosite (Warren 1993; Arai et al. 2008; Yamaguchi et al., 2010; Fig. 6). Relict olivine grains in Clast 1, 2 and 3 vary in Mg content. In particular, those in Clast 1 have significantly different Fe-Mg compositions. Clast 5 contains a few olivine grains which preserve chemical zoning (core $\text{Fo}_{90.2}$, rim $\text{Fo}_{88.2}$). We found, however, irregular-shaped olivine grains in Clast 6 that appear to be relict fragments of a target assemblage, but whose compositions could not be distinguished from newly crystallized olivine grains in the melt matrix and may have equilibrated with surrounding melt.

Relict pyroxene and olivine grains in Clast 1 have bimodal ($\text{En}_{77-72}\text{Wo}_{3.5-2.5}$ and $\text{En}_{69-58}\text{Wo}_{5.2-2.5}$; Fo_{91-77} and Fo_{75-69} ; Fig. 6) and Fe-rich compositions compared to those in other clasts. Because the pyroxene is not in equilibrium with the olivine, they may be derived from three or possibly four target sequences. Relict pyroxene grains and newly crystallized grains are similar in Mg#, but distinct in Al and Cr compositions in Clasts 2 and 3; relict grains have lower values of Al and Cr relative to melt products. This implies that the Fe-Mg composition of pyroxene might be disturbed by the surrounding melt. Relict olivine and pyroxene grains in Clasts 2, 3, and 4 have similar ferroan compositions (Fig. 6) and could represent the same target sequence. Relict olivine grains in Clast 5 and

6 have more magnesian compositions than other melt clasts in 60016, which could be derived from other target sequences (potentially similar to the more magnesian of the two olivine-bearing sequences in Clast 1). Even if relict pyroxene and olivine grains are equilibrated with surrounding melt, the compositional variation in six clasts cannot be explained by equilibrium with a single impact melt. Compositional differences in relict minerals, especially olivine grains, imply that those clasts originated from at least two different target sequences (high Mg# and low Mg# in olivine grains).

Fe-Ni metal droplets are found in all of the melt clasts, although the abundances of metal are different among the six clasts (0.4 to 3.5 %). All cm-sized clasts contain abundant Fe-Ni metal grains of meteorite origin (Goldstein and Yakowitz, 1971), as implied by the Ni/Co ratio which are within the range of meteorite composition (Korotev, 1987a,b, 1996). Osmium isotopes and highly siderophile element compositions for adjacent samples with our specimens (splits, 286b and 290b), and other Apollo 16 impact melt clasts indicate multiple, diverse impactors ranging from chondritic to suprachondritic (Liu et al., 2015). Metal droplets in Clasts 2 and 5 (probably Clasts 1, 4 and 6 also, but they cannot be characterized due to small grain size) have eutectic textures with schreibersite (El Goresy et al., 1971; Doan and Goldstein 1970; Gooley et al. 1973). Schreibersite is the main P-phase in those impact melt clasts. Clast 3 does not contain schreibersite, but instead contains phosphate. All melt clasts have similar P_2O_5 concentrations, but there is no correlation with the abundance of meteoritic Fe-Ni metal. Thus, the phosphorus component of schreibersite and phosphate is possibly derived from the lunar target rock and not from projectile contamination.

To compare the bulk compositions of our 6 clasts with previously reported impact melt rocks, we conducted multivariate analysis using a compilation of 95 published analyses with data for 9 elements (Si, Ti, Al, Fe, Mg, Ca, Na, K, and Cr), which were previously classified using trace element information using the methodology of Korotev (1994). We conducted principal component analysis (PCA), a method of dimension reduction of data set, to identify and interpret compositional variations and obtaining PC scores. The first principal component (PC1) reflects maximum variation, followed by a second (PC2) component. Using this method we obtained results consistent with Korotev (1994) and can separate chemical group 1, 2, 3, and 4 (Fig. 7). Moreover, our PCA method can also identify sub-types of group 1 melt (1M and 1F). PC1 scores show a positive correlation with the value of Al+Na+K and negative correlation with Mg+Fe. Thus, PC1 separates mafic from felsic melts. PC2 scores are correlated with the Mg/Fe ratio, which highlights the types of variations seen among group 2 melts (2M, 2Mo, 2DB, 2NR, and 2F). The six clasts from 60016 processed with the same PCA methods can be separated. Although the method utilizes measurements obtained with DBA techniques that, thus, have compositional uncertainty mostly in Mg/Fe ratio, the six clasts from 60016 appear to be from 2 or 3 groups based on PC1 scores (clast 1 and 5 have higher values relative to other clasts), which suggests the clasts originate from 2 or 3 different pre-impact target materials. DBA compositions of Clasts 1, 5 and 6 (either averaged DBA and corrected values) plot in the regions of group 2 melts and Clast 3 plots in the region of Group 1 melt. Korotev (1996) analyzed bulk trace elements of Sm and Sc for 1-3 mm sized mafic clasts in ancient regolith breccias include 60016. These clasts have compositions that range among group 1 and 2 melts (Korotev, 1994). Our centimeter-sized clasts have similar compositions to these clasts although we did not analyze trace elements.

We also plotted (Fig. 8) the major element compositions of the six clasts and compared them to previously classified Apollo 16 impact melt rocks (e.g., Korotev, 1994 and references therein). K_2O and P_2O_5 concentrations of all six clasts are similar to group 1 melt rocks (Korotev, 1994), although Clast 1 and Clast 5 have higher Al_2O_3 and lower $MgO+FeO$ compositions relative to group 1. The bulk compositions (lower Al_2O_3 and higher $MgO+FeO$) of Clast 2, 3, 4 and 6 are similar to thorium-rich mafic impact melt breccias reported by Haskin et al., 1998. These differences in Al_2O_3 and $MgO+FeO$ values for our six impact melt clasts are significant even if the values contain systematic errors. They appear to reflect higher amounts ($>30\%$) of relict plagioclase in Clasts 1 (39 modal %) and 5 (34 modal %) compared to Clasts 2 (2 modal %), 3 (29 modal %), 4 (10 modal %), and 6 (6 modal %).

We could not find any correlation between petrographic texture and bulk major element compositions of melt clasts. The compositional difference might be implying (1) all six impact melt clasts are created by a single impact into heterogeneous target sequences or (2) individual impact events have occurred on different targets.

Norman et al. (2006) classified Apollo 16 impact melt rocks into 8 petrographic groups (mafic poikilitic group, aluminous subophitic group, troctolitic vitrophyres group, aluminous poikilitic group, dimict, meta-poikilitic, North Ray Crater, and unclassified) on the basis of petrographical signatures. In terms of compositional groups (Korotev, 1994), Group 1 melts have poikilitic textures with lower Al_2O_3 (17-23 wt.%) components and Ar-Ar ages of 3865 Ma. Four of our centimeter sized clasts (Clasts 2, 3, 4 and 6) have similar major element compositions to group 1 melt (Korotev, 1994) and Clasts 2 and 3 have similar poikilitic textures to the mafic poikilitic group (Norman et al., 2006). On the other hand, Clasts 4 and 6 have aphanitic or porphyritic texture, respectively, nevertheless major element compositions are similar to group 1 melts. Clast 1 has a poikilitic texture with a high aluminum content ($Al_2O_3 = 26.4$ wt.%). Although the texture of Clast 1 is similar to that of Clasts 2 and 3, bulk compositions are significantly different (higher Al_2O_3 and lower $MgO+FeO$ compositions). These data indicate the melts reflect a diversity of target sequences. Lawrence et al. (2013) reported that ferroan anorthosite (FAN) and Mg-suite (MGS) rocks contribute to fragment-rich and fine-grained Apollo 16 impact melts, implying both types of lithologies existed in the pre-target region, and the relative abundances of FAN and MGS materials in targets of Apollo 16 and Apollo 17 impact melt breccias were different.

Fe-Ni metal droplets are found in all of the melt clasts, although the abundances of metal are different among the six clasts (0.4 to 3.5 %). Metal droplets in Clasts 2 and 5 (probably Clasts 1, 4 and 6 also, but they cannot be characterized due to small grain size) have eutectic textures with schreibersite (El Goresy et al., 1971; Doan and Goldstein, 1970; Gooley et al., 1973). Schreibersite is the main P-phase in those impact melt clasts. Clast 3 does not contain schreibersite, but instead contains phosphate. All melt clasts have similar P_2O_5 concentrations, but there is no correlation with the abundance of meteoritic Fe-Ni metal, although Ni and Co compositions are within the range of meteoritic materials. Korotev et al. (2003) measured siderophile element signatures of metallic grains in Apollo 16 and 17 impact melt breccias and found high concentrations of W up to several hundred over concentrations than meteoritic metals. Therefore, the phosphorus component originated not from impactors but the lunar target rock (Wlotzka et al., 1972; 1973). The bulk K_2O (0.3-0.5 wt.%) and P_2O_5 (0.3-0.5 wt.%) abundances in

the melt clasts are higher than the local regolith ($K_2O = \sim 0.12$ wt.%, $P_2O_5 = \sim 0.12$ wt.%; Korotev 1997, Lucey et al., 2006; Table 1). That requires an origin in an impact event beyond the landing site region, possibly the Procellarum KREEP Terrain (PKT) (Haskin et al., 1998; Jolliff et al., 2000; Korotev, 2000; Wieczorek and Phillips, 2000), which is >400 km away. Interestingly, the melt clasts in 60016 are also enriched relative to two of the best candidates for Imbrium impact melt, samples 15445 and 15455. Those two samples have K_2O and P_2O_5 of ~ 0.2 wt.% (Table 1).

In summary, the petrological evidence suggests the six clasts represent at least 4 target sequences which have different bulk major element compositions and mafic mineral compositions. Clast 1 appears to come from a terrain with high bulk Al_2O_3 and bimodal compositions in mafic minerals (Mg-suite and ferroan anorthosite), Clasts 2, 3 and 4 may come from a terrain with low Al_2O_3 composition and relatively ferroan composition in mafic minerals (Mg-suite), Clast 5 may have come from a terrain with high bulk Al_2O_3 and high Mg# in mafic minerals (Mg-suite), and Clast 6 may have come from a terrain with low bulk Al_2O_3 and high Mg# in mafic minerals (Mg-suite). In all cases, a component of KREEP is required. Compared to the classic samples of Imbrium impact melt, 15445 and 15455, the clasts have a larger K and similar P concentrations (Table 1). The Mg-suite and ferroan anorthosite components in the clasts are also seen in Imbrium melts.

Multiple impact events

One of our goals was to determine if the ejecta material covering the Apollo 16 site could be explained by one impact (Imbrium) or if it would be better explained by several impacts over a short period. This work is important in order to better understand the timing of impactor populations in the Apollo 16 site, to look for evidence of the late heavy bombardment contained in retention ages of intermediate (cm) sized clasts, and for the expansion and comparison of the Apollo impact breccias. Our intent was to gain more insight into the different age distributions and different impact histories that are implied by different sources and sizes of lunar impact material. Upon comparing the impact age distributions of our intermediate sized melt breccias, we find they follow the same trend as more massive Apollo melt rocks, rather than the sub-millimeter sized impact melt clasts in lunar meteorites (Cohen et al., 2005) or lunar impact glasses (Culler et al., 2000; Levine et al., 2005; Zellner et al., 2009). We find evidence for at least four distinct impact ages for our six clasts (Fig. 5 – the age of Clast 4 is poorly defined, and that of Clast 3 is difficult to interpret, and our best estimate is not well-resolved from the age of Clast 2), despite some interpretations that this area is overlain with Imbrium or Nectaris ejecta (Hodges and Muelberger, 1981; Spudis and Peiters, 1991), and it does not appear that the spread in ages could have been from a single event. This is consistent with the petrological analysis, which shows different textures and mineral compositions from one clast to another. Although all of the plateaus are small enough that the ages may underestimate the true age, they are all large enough ($>50\%$ of the ^{39}Ar from the host phase) that they should be within $\sim 2\sigma$, the stated uncertainty, based on modeling described below (and in an abstract by Swindle and Weirich, 2017).

With one exception, the different splits of a single sample followed the same general trend in their apparent age spectra, as well as agreeing in age. The exception was Clast 4, represented by one split of 60016,305 and two of 60016,314. Each of its splits showed a

disturbed apparent age spectra and resulted in different ages ranging about 50 Myr apart from one another. Clast 4 must be a very heterogeneous sample, even on the mm scale, as the samples (,305 and ,314) were located adjacent to one another and still showed different apparent age patterns, which may reflect the petrological complexity described above. Proposed impact ages for all samples range from 3877.5 ± 13.8 to 4000.0 ± 23.2 Ma, not counting Clast 4 (3823 ± 75 Ma).

Until recently, the quoted age of Imbrium was usually 3.85 ± 0.02 Ga (e.g., Wilhelms, 1987; Ryder, 1990, 1994; Stöffler and Ryder, 2001). If taken at face value, that would imply our clasts represent older impact melts that may have been excavated by Imbrium and deposited as part of the Cayley Formation at the Apollo 16 site or, alternatively, older impact melts produced elsewhere and ejected on top of the Cayley Formation after the Imbrium impact event.

As discussed by Nemchin et al. (2009), however, Imbrium may be older than 3.85 Ga. U-Pb analyses of apatite in Apollo 14 samples of Imbrium ejecta suggest an age of 3926 ± 3 Ma (Merle et al., 2014; which agrees with Clast 5, 3926.5 ± 7.3). Similarly, U-Pb analyses of phosphate grains in 12013 give an indistinguishable age of 3924 ± 4 Ma (Thiessen et al., 2018). If Imbrium is 3926 ± 3 Ma, then both of the scenarios above are possible, but the second one becomes easier as there is more time for mixing to occur after the Imbrium impact event occurs and before the breccia is closed. Since the K_2O and P_2O_5 abundances in the melts are higher than those in the local regolith, those other impacts probably occurred far from the Apollo 16 site, perhaps 400 km or farther.

In either case, the ages of the melts are broadly consistent with estimates of the latter stages of the basin-forming epoch, as tabulated by Norman (2009), including Orientale (~ 3750 Ma), Imbrium (3850 Ma, or, based on the recent U-Pb, 3925 Ma), Serenitatis (3890 Ma), and perhaps Nectaris (3900–4200 Ma). These ages are in rough agreement with the Ar-Ar analyses of group 1 and 2 (Korotev, 1994) clasts by Norman et al. (2006), who analyzed 25 Apollo 16 impact melt rocks. They reported four clusters with each cluster having common ages, mineralogy and chemistry, with cluster ages ranging from 3771 Ma to 3913 Ma. Three of our samples, with different mineralogy and chemistry, cover the same age range, and again have distinct ages, while our fourth (Clast 2) is at a slightly higher age (Fig. 5). Assigning specific ages to specific basins is critical, but is also notoriously difficult, and may remain controversial until appropriate samples are returned in future missions. However, the existence of multiple robust ages within the same sample, as seen here (as well as in Ar-Ar ages of other Apollo 16 melt rocks by Norman et al, 2006 and in U-Pb ages of Apollo 14 phosphates by Snape et al. (2016), suggests that there were multiple events in a rather short period of time.

All of our samples show partial resetting of the ^{40}Ar - ^{39}Ar system, which is very common among lunar samples. Shuster et al. (2010) point out that the partial resetting may contain useful information to the overall impact history beyond that provided by the plateau ages alone. They argue that since the initial steps in the apparent age spectra represent the maximum age of an event in which significant ^{40}Ar loss occurred, it could thus provide a more quantitative contribution to the determination of the impact history than previously considered. The results of Shuster et al. (2010) show a minimum apparent age of 3300–3400 Ma for five of seven mm-sized rocks from regolith sample 63503 (collected at a different station than 60016), in some cases giving what they referred to as a “sub-plateau”. They performed detailed modeling that showed that an

impact at 3300-3400 Ma could account for the partial argon loss from their samples, and indeed could correspond to the age of an impact event. We do not see, however, the 3.3-3.4 Ga event reported by Shuster et al. 2010 in their analyses of seven 2-4 mm-diameter particles in sample 63503, or any other well-defined event. We note that 63503 is a soil that has been open to lithologic contributions up to the present day, whereas 60016 is a consolidated regolith breccia that was closed to input 3.74 Ga (Joy et al., 2011), so we would expect that all of the clasts would have the same thermal history after 3.74 Ga. Since we see no clustering of minimum ages or other post-3.74 Ga ages, the most likely scenario is that all are reflecting varying degrees of minor ($\sim 10\%$ or less, based on our modeling) ^{40}Ar loss, at ~ 1.5 Ga (the lowest minimum, in Clast 3) or more recently.

Robustness of plateau ages

Taken at face value, the plateaus suggest that there are at least four different ages, and hence at least four different impact events, represented. A key question, then, is how robust the plateaus are. As mentioned above, there have been different definitions of what constitutes a plateau applied in different studies, so we will first consider the idealized case of a diffusive event affecting a single phase. We will then consider some of the complications introduced by the multi-minerallic samples that we (and most other studies of lunar impact melts) deal with. While there are a number of complicating factors that can influence apparent ages, including secondary mineralization, undegassed clasts, recoil during irradiation, and trapped (or “excess”) ^{40}Ar (see discussions in Jourdan, 2012; McDougall and Harrison, 1999), we believe the primary effect in most of our samples is diffusive loss in multiple events.

Motivated by the results in this work, Swindle and Weirich (2017) modeled a monominerallic system that experienced partial diffusive loss, varying the time of resetting and the amount of partial loss to determine what conditions generate apparent plateaus, and how those plateaus relate to the time of the resetting event or the time of the age before resetting (what we will refer to as the original age). In any case with partial diffusive loss, the highest apparent age, in the last release step, will be lower than the original age, and shape of the release pattern on a plateau plot will be flattest at the highest temperatures. They showed that in a monominerallic system that only experienced mild diffusive loss, plateaus representing 50-70% of the total ^{39}Ar released produce ages $\sim 1-2\sigma$ lower than the original age. Subsequent modelling we have done shows that for samples experiencing near-complete loss, plateaus of that width are $\sim 1-2\sigma$ higher than the resetting age. Since the samples analyzed here are polyminerallic, as suggested by the behavior of the plateau plots and confirmed by the K/Ca ratios, this adds a complication, but it is the fraction of the ^{39}Ar coming from a specific host phase that matters, and the release patterns of different phases are often well separated, so most of the ages given are based on plateaus of 50% or more. Furthermore, the lowest-temperature ages are significantly lower than the plateaus, suggesting that we are not in the regime of near-complete resetting. Detailed modelling using the techniques of Swindle and Weirich (2017) suggest loss amounts of 10% or less in most cases. While the results of Swindle and Weirich (2017) indicate that these ages probably underestimate the true ages (as might be obtained by, for example, the U-Pb system), the fact that they are similar width plateaus suggests that they probably underestimate all the ages by similar amounts, so the distinction between the different ages obtained remains. In

addition, a numerical simulation of the deposition of basin-size ejecta (Fernandes and Artemieva, 2012) suggest that temperatures in Imbrium ejecta deposited downrange near the Apollo 16 landing would have had temperatures no greater than 130 °C, which is far below the closure temperature (c. 300 °C; Cassata et al., 2010). Thus, it would not have been hot enough to produce different ages among regions of entrained melt.

To summarize, we believe that our estimated ages are systematically lower than the true formation ages of the clasts, but by $1-2\sigma$ (<25 Ma). Thus, while detailed comparison with other chronometers would be problematic, we believe that since our estimated ages are distinct from one another by up to several tens of Ma, they are not all the result of different degrees of diffusive loss in a large later thermal event such as Imbrium, but instead represent samples that experienced complete loss in events a few tens of Ma apart (one of which might be Imbrium), then varying degrees of mild resetting ($<10\%$) in a later, smaller, event (or events).

Delivery mechanism of the impact melt clasts

Our six impact melt clasts found in same ancient regolith breccia 60016 (within ~10 cm) contain KREEPy materials and have different impact ages around 4.0 Ga to 3.7 Ga. These compositions and impact ages imply that the clast-forming impact events might have occurred on or near the PKT region and that the clasts were then delivered to Apollo 16 site. The Apollo 16 landing site is located on highland crust, at least 400 km away from the PKT region. The impact melt clasts must have been delivered such a long distance into the same area, and then lithified as an ancient regolith breccia. One simple delivery system would be that each impact event dispersed melt clasts, and that these happened to all land in the same place, the Apollo 16 landing site, where they were lithified into a regolith breccia by a local shock event. The problem for this mechanism is that our sample contains no KREEP-poor impact melt clasts that might have originated from closer to the Apollo 16 site, outside of PKT region. The population of impact events should be same between PKT and other places, thus non-PKT origin impact melt should exist in these samples in this scenario. Another possible mechanism is that each impact melt clast was created by a small impact event on the PKT region. Those melt clasts were deposited near one another and then delivered to the Apollo 16 site by the Imbrium basin forming impact event. If the Apollo rocks experienced the latter scenario, the age of Imbrium forming event should be revised (to a younger value). We have examined only six impact melt clasts from a single rock, so, we cannot judge which scenario is most realistic. To solve the mystery of the delivery system, further data from more lunar impact melt rocks are required.

Conclusion

We have undertaken a petrological and geochronological study of six centimeter-sized impact melt clasts in 60016. Three clasts have similar poikilitic textures, but their mineral abundances, size, chemical compositions, and ages are different from each other implying they have originated from different impact events and had different cooling history after impact melt forming events. All of the melt clasts contain similar amounts of K_2O and P_2O_5 contents that might be derived from a KREEP component in their target regions. The apparent ^{40}Ar - ^{39}Ar ages agree from split to split of the same sample, but are distinct from clast to clast. Ages of our samples fall in the range proposed for the late heavy bombardment, though other studies have shown older ages exist (Norman et al., 2007). Four of the six clasts produced multi-step plateaus that range from 3877.5 ± 13.8 to 3970.1 ± 12.8 Ma. The age range of all clasts is from 3823 ± 75 to 4000.0 ± 23.2 Ma, with a cluster around 3900 Ma. This cannot be explained by a variable degrees of resetting in a single (Imbrium) impact, and requires more than one event within the short duration. In fact, our analysis suggests evidence for four distinct impact events, which are broadly consistent with estimates of some of the most recent impact basins (Norman, 2009). We do not have good enough evidence of closure ages from our partially reset ^{40}Ar - ^{39}Ar ages to additionally constrain the ages of lunar impacts.

Collectively, the petrological texture, mineral compositions, and Ar-Ar age data suggest these six cm-sized clasts in 60016 represent different impact events on the early history of the Moon. This means that attempts to interpret the pile-up of ages circa 3.9 Ga as an artifact of a single (Imbrium) impact are unwarranted. There seems to have been a large number of impact events in a short period of time.

Acknowledgments

Samples are allocated from Astromaterials Acquisition and Curation Office at the NASA Johnson Space Center (JSC). Axel Wittmann helped DAK when selecting samples in the Lunar Sample Laboratory Facility at JSC. We are grateful to Drs. C. N. Achilles, C. E. Isachsen, A. H. Peslier and D. K. Ross for support on laboratory work. We also thank Drs. M.D. Norman, K. H. Joy, D. Mittelfehldt and A. Wittmann for their useful comments and discussion. Constructive comments from R. L. Korotev and anonymous reviewer improved and clarified the manuscript. We also thank for A. J. T. Jull (editor) and A. Yamaguchi (associate editor) for handling of the manuscript. This work was supported by NASA Lunar Science Institute Cooperative Agreement NNA09DB33A and NASA Solar System Exploration Research Virtual Institute Cooperative Agreement NNA14AB07A (PI David A. Kring).

References

- Albee A. L., Gancarz A. J. and Chodos A. A. 1973. Metamorphism of Apollo 16 and 17 and Luna 20 metaclastic rocks at about 3.95 AE: Samples 61156, 64423, 14-2, 65015, 67483, 15-2, 76055, 22006, and 22007. *Proceedings of the Lunar Science Conference* 4:569–595.
- Arai T., Takeda H., Yamaguchi A. and Ohtake M. 2008. A new model of lunar crust: asymmetry in crustal composition and evolution. *Earth Planets Space* 60:433–444.
- Bence A. E., Papike J. J., Sueno S. and Delano J. W. 1973. Pyroxene poikiloblastic rocks from the lunar highlands. *Proceedings of the Lunar Science Conference* 4:597–611.
- Boehnke P. and Harrison T.M. 2016. Illusory late heavy bombardments. *Proceedings of the National Academy of Sciences of the United States of America* 113, 10802–10806.
- Bogard D. D. 1995. Impact age of meteorites: A synthesis. *Meteoritics* 30:244–268.
- Bogard D. D. and Garrison D. H. 2003. ^{39}Ar – ^{40}Ar ages of eucrites and thermal history of asteroid 4 Vesta. *Meteoritics & Planetary Science* 38:669–710.
- Bottke W. F. and Norman M. D. 2017. The Late Heavy Bombardment. *Annual Review of Earth and Planetary Sciences* 45:619–647.
- Cassata W. S., Renne P. R. and Shuster D. L. 2010. Argon diffusion in plagioclase and implications for thermochronometry: A case study from the Bushveld Complex, South Africa. *Geochimica et Cosmochimica Acta* 73:6600–6612.
- Cassata W. S., Shuster D. L., Renne P. R. and Weiss B. P. 2011. Evidence for shock heating and constraints on Martian surface temperatures revealed by $^{40}\text{Ar}/^{39}\text{Ar}$ thermochronometry of Martian meteorites. *Geochimica et Cosmochimica Acta* 74:6900–6920.
- Cohen B. A., Swindle T. D. and Kring D. A. 2000. Support for the Lunar Cataclysm Hypothesis from Lunar Meteorite Impact Melt. *Science* 290:1754–1756.
- Cohen B. A., Swindle T. D. and Kring D. A. 2005. Geochemistry and ^{40}Ar – ^{39}Ar geochronology of impact-melt clasts in feldspathic lunar meteorites: Implications for lunar bombardment history. *Meteoritics & Planetary Science* 40:755–777.
- Crow C. A., McKeegan K. D. and Moser D. E. 2017. Coordinated U–Pb geochronology, trace element, Ti-in-zircon thermometry and microstructural analysis of Apollo zircons. *Geochimica et Cosmochimica Acta* 202:264–284.
- Culler T. S., Becker T. A., Muller R. A. and Renne P. R. 2000. Lunar impact history from $^{40}\text{Ar}/^{39}\text{Ar}$ dating of glass spherules. *Science* 287:1785–1788.
- Dalrymple G. B. and Ryder G. 1993. $^{40}\text{Ar}/^{39}\text{Ar}$ age spectra of Apollo 15 impact-melt rocks by laser step-heating and their bearing on the history of lunar basin formation. *Journal of Geophysical Research* 98:13,085–13,095.
- Dalrymple G. B. and Ryder G. 1996. Argon-40/argon-39 age spectra of Apollo 17 highlands breccia samples by laser step heating and the age of the Serenitatis basin. *Journal of Geophysical Research* 101:26,069–26,084.
- Deino A. 2001. Mass Spec, Version 7.884. Special Publication. Berkeley Geochronology Center.
- Doan A. S. and Goldstein J. I. 1970. The ternary phase diagram, Fe–Ni–P. *Metallurgical Transactions* 1:1759–1767.

- Dowty E., Prinz M. and Keil K. 1973. Composition, mineralogy, and petrology of 28 mare basalts from Apollo 15 rake samples. *Proceedings of the Lunar Science Conference* 4:423–444.
- Duncan A. R., Erlank A. J., Willis J. P. and Ahrens L. H. 1973. Composition and interrelationships of some Apollo 16 samples. *Proceedings of the Lunar Science Conference* 4:1097–1113.
- Eugster O. 1999. Chronology of dimict breccias and the age of South Ray crater at the Apollo 16 site. *Meteoritics & Planetary Science* 34:385–391.
- El Goresy A., Ramdohr P. and Taylor L. A. 1971. The geochemistry of the opaque minerals in Apollo 14 crystalline rocks. *Earth and Planetary Science Letters* 13:121–129.
- Fernandes V.A., Artemieva N. 2012. Impact ejecta temperature profile on the moon-what are the effects on the Ar-Ar dating method? (abstract #1367). 43rd Lunar and Planetary Science Conference.
- Fernandes V.A., Fritz J., Weiss, B.P., Garrick-Bethell I., and Shuster D.L. 2013. The bombardment history of the Moon as recorded by ^{40}Ar - ^{39}Ar chronology. *Meteoritics & Planetary Science* 48:241–269.
- Fleck R. J., Sutter J. F., and Elliot D.H., 1977. Interpretation of discordant $^{40}\text{Ar}/^{39}\text{Ar}$ age-spectra of Mesozoic tholeiites from Antarctica. *Geochimica et Cosmochimica Acta* 41:15–32.
- Ganapathy R., Laul J. C., Morgan J. W., and Anders E. 1972. Moon: Possible nature of the body that produces the Imbrium basin, from the composition of Apollo 14 samples. *Science* 175:55–59.
- Gooley R. C., Brett R. and Warner J. L. 1973. Crystallization history of metal particle in Apollo 16 rake samples. *Proceedings of the Lunar Science Conference* 4:799–810.
- Gomes R. S., Levison H. F., Morbidelli A., and Tsiganis K. 2005. Origin of the cataclysmic Late Heavy Bombardment period of the terrestrial planets. *Nature* 435:466–469.
- Grange M. L., Nemchin A. A., Pidgeon R. T., Timms N., Muhling J. R. and Kennedy A. K. 2009. Thermal history recorded by the Apollo 17 impact melt breccia 73217. *Geochimica et Cosmochimica Acta* 73:3093–3107.
- Grange M. L., Nemchin A. A., Timms N., Pidgeon R. T., and Meyer. C. 2011. Complex magmatic and impact history prior to 4.1 Ga recorded in zircon from Apollo 17 South Massif aphanitic breccia 73235. *Geochimica et Cosmochimica Acta* 75:2213–2232.
- Haskin L. A., Helmke P.A., Blanchard D. P., Jacobs J. W. and Telander K. 1973. Major and trace element abundances in samples from the lunar highlands. *Proceedings of the Lunar Science Conference* 3:1275–1296.
- Haskin L. A., Korotev R. L., Rockow K. M., and Jolliff B. L. 1998. The case for an Imbrium origin of the Apollo thorium-rich impact-melt breccias. *Meteoritics & Planetary Science* 33:959–975.
- Hartmann W. K. 2003. Megaregolith evolution and cratering cataclysm models—lunar cataclysm as a misconception (28 years later). *Meteoritics & Planetary Science* 38:579–593.

- Hartung J. B. 1974. Can random impacts cause the observed Ar 39/40 age distribution for lunar highland rocks? *Meteoritics* 9:349.
- Hodges C. A. 1972. Geological map of part of the Descartes region of the Moon. Geologic Atlas of the Moon. Part of the Descartes region-Apollo 16. United States Geological Survey Map I-748 Part 2. Scale 1:50000.
- Hodges C. A., and Muehlberger W. R. 1981. Summary and critique of geologic hypotheses. In *Geology of the Apollo 16 Area, Central Lunar highlands*, edited by G. E. Ulrich, C. A. Hodges, and W. R. Muehlberger. United States Geological Survey Professional Paper 1048:215-230.
- Hodges C. A., Muehlberger W. R. and Ulrich G. E. 1973. Geologic setting of Apollo 16. *Proceedings of the Lunar Science Conference* 4:1-25.
- Hubbard N. J., Rhodes J. M., Gast P. W., Bansal B. M., Chih C.-Y., Wiesmann H. and Nyquist L. E. 1973. Lunar rock types: The role of plagioclase in non-mare and highland rock types. *Proceedings of the Lunar Science Conference* 4:1297-1312.
- James O. B. 1994. Siderophile and volatile elements in Apollo 17 impact melt rocks (abstract). *Lunar and Planetary Science Conference* 25:617-618.
- James O. B. 1995. Siderophile elements in lunar impact melts: Nature of the impactors (abstract). *Lunar and Planetary Science Conference* 26:671-672.
- James O. B. 1996. Siderophile elements in lunar impact melts define nature of the impactors (abstract). *Lunar and Planetary Science Conference* 27:603-604.
- Jolliff B. L. 1998. Large-scale separation of K-frc and REEP-frc in the source regions of Apollo impact-melt breccias, and a revised estimate of KREEP composition. *International Geology Review* 40:916-935.
- Jolliff, B. L., Gillis, J. J., Haskin, L. A., Korotev, R. L. and Wieczorek M. A. 2000. Major lunar crustal terranes: surface expressions and crust-mantle origins. *Journal of Geophysical Research* 105:4197-4416.
- Jourdan F. 2012. The $^{40}\text{Ar}/^{39}\text{Ar}$ dating technique applied to planetary science and terrestrial impacts. *Australian Journal of Earth Science* 59:199-244.
- Jourdan F., Renne P. R., and Reimold W. U. 2007. The problem of inherited $^{40}\text{Ar}^*$ in dating impact glass by the $^{40}\text{Ar}/^{39}\text{Ar}$ method: Evidence from the Tswaing impact crater (South Africa). *Geochimica et Cosmochimica Acta* 71:1214-1231.
- Joy K. H., Kring D. A., Bogard D. D., McKay D. S. and Zolensky M. E. 2011. Re-examination of the formation ages of the Apollo 16 regolith breccias. *Geochimica et Cosmochimica Acta* 75:7208-7225.
- Joy K. H., Zolensky M. E., Nagashima K., Huss G. H., Ross D. K., McKay D. S. and Kring D. A. 2012. Direct detection of projectile relicts from the end of the lunar basin-forming epoch. *Science* 336:1426-1429.
- Korotev R. L. 1994. Compositional variation in Apollo 16 impact-melt breccias and inferences for the geology and bombardment history of the Central Highlands of the Moon. *Geochimica et Cosmochimica Acta* 58:3931-3969.
- Korotev R. L. 1996. On the relationship between the Apollo 16 ancient regolith breccias and feldspathic fragmental breccias, and the composition of the prebasin crust in the Central Highlands of the Moon. *Meteoritics & Planetary Science* 31, 403-412.
- Korotev R. L. 1997. Some things we can infer about the Moon from the composition of the Apollo 16 regolith. *Meteoritics & Planetary Science* 32:447-478.

- Korotev R. L. 2000. The great lunar hot spot and composition and origin of the Apollo mafic ("LKFM") impact melt breccias. *Journal of Geophysical Research* 105:4317–4345.
- Korotev R. L., Jolliff B. L., Campbell A. J., and Humayun M. 2003. Laser-Abration ICP-MS analyses of meteoritic metal grains in lunar impact melt breccias (abstract #1487). 34th Lunar and Planetary Science Conference.
- Kring D. A. and Cohen B. A. 2002. Cataclysmic bombardment throughout the inner solar system 3.9–4.0 Ga. *Journal of Geophysical Research* 107(E2)5009.
- Laul J. C., Jolliff B. L. and Schmitt R. A. 1974. Chemical studies of Apollo 16 and 17 samples. *Proceedings of the Lunar Science Conference* 5:1047–1066.
- Lawrence S. J., Taylor G. J., and Norman M.D. 2013. Trace element geochemistry of mineral, clasts in Apollo 16 impact melt breccias (abstract #2848). 44th Lunar and Planetary Science Conference.
- Levine J., Becker T. A., Muller R. A. and Renne P. R. 2005. $^{40}\text{Ar}/^{39}\text{Ar}$ dating of Apollo 12 impact spherules. *Geophysical Research Letters* 32:L15201.
- Lindstrom M. M., Nava D. F., Lindstrom D. J., Winzer S. R., Lum R. K. L., Schuhmann P. J., Schuhmann S. and Philpotts J. A. 1977. Geochemical studies of the white breccia boulders at North Ray Crater, Descartes region of the lunar highlands. *Proceedings of the Lunar Science Conference* 8:2137–2151.
- Liu J., Sharp, M., Ash, R.D., Kring, D.A., Walker, R.J. 2015. Diverse impactors in Apollo 15 and 16 impact melt rocks: Evidence from osmium isotopes and highly siderophile elements. *Geochimica et Cosmochimica Acta* 155:122–153.
- LSAPT. 1970. Summary of Apollo 11 lunar science conference. *Science* 167:449–451.
- LSPET (Lunar Sample Preliminary Examination Team) 1973. The Apollo 16 lunar samples: Petrographic and chemical description. *Science* 179:23–34.
- Lucey P., Korotev R. L., Gillis J. J., Taylor L. A., Lawrence D., Campbell B. A., Elphic R., Feldman B., Hood L. L., Hunten D., Mendillo M., Noble S., Papike J. J., Reedy R. C., Lawson S., Prettyman T., Gasnault O., and Maurice S. 2006. Understanding the lunar surface and space–Moon Interactions. In *New views of the Moon*, edited by B.L. Jolliff, M. A., Wiczorek, C. K. Shearer, and C. R. Neal. *Reviews in Mineralogy and Geochemistry* 60:83–219.
- Marchi S., Bottke W. F., Kring D. A., and Morbidelli A. 2012. The onset of the lunar cataclysm as recorded in its ancient crater populations. *Earth and Planetary Science Letters* 325–326:27–38.
- McCubbin F. M., Jolliff B. L., Nekvasil H., Carpenter P. K., Zeigler R. A., Steele A., Elardo S. M., and Lindsley D. H. 2011. Fluorine and chlorine abundances in lunar apatite: Implications for heterogeneous distributions of magmatic volatiles in the lunar interior. *Geochimica et Cosmochimica Acta* 75:5073–5093.
- McDougall, I., Harrison, T.M., 1999. *Geochronology and thermochronology by the $^{40}\text{Ar}/^{39}\text{Ar}$ method*. Oxford University Press, New York, pp. 212.
- McKay D. S., Bogard D. D., Morris R. V., Korotev R. L., Johnson P., and Wentworth S. J. 1986. Apollo 16 regolith breccia: Characterization and evidence for early formation in the Mega-Regolith. *Proceedings of the Lunar and Planetary Science Conference* 16:D277–D303.

- Mercer C. M., Young K. E., Weirich J. R., Hodges K. V., Jolliff B. L., Wartho J.-A., and Van Soest M. C. 2015. Refining lunar impact chronology through high spatial resolution $^{40}\text{Ar}/^{39}\text{Ar}$ dating of impact melts. *Science Advances* 1:e1400050.
- Merle, R.E., Nemchin, A. A., Grange, M. L., Whitehouse, M. J., and Pidgeon R.T., 2014 High resolution U-Pb ages of Ca-phosphates in Apollo 14 breccias: Implications for the age of the Imbrium impact. *Meteoritics & Planetary Science* 49:2241–2251.
- Meyer C., Anderson D. H., and Bradley J. G. 1974. Ion microprobe mass analysis of plagioclase from “non-mare” lunar samples. *Proceedings of the Lunar Science Conference* 5:685-706.
- Morbidelli A., 2010. A coherent and comprehensive model of the evolution of the outer Solar System. *Comptes Rendus Physique*, 11:651–659.
- Morbidelli A., Levison H. F., Tsiganis K., and Gomes R. 2005. Chaotic capture of Jupiter’s Trojan asteroids in the early Solar System. *Nature* 435:462–465.
- Morbidelli A., Marchi S., Bottke W. F., and Kring D. A. 2012. A sawtooth-like timeline for the first billion year of lunar bombardment. *Earth and Planetary Science Letters* 355–356:144–151.
- Morris R. V., See T. H., and Hörz F. 1986 Composition of the Cayley formation at Apollo 16 as inferred from impact melt splashes. *Proceedings of the Lunar and Planetary Science Conference* 17:E21-E42.
- Nemchin A. A., Pidgeon R. T., Healy D., Grange M. L., Whitehouse M. J., and Vaughan J. 2009. The comparative behavior of apatite-zircon U-Pb systems in Apollo 14 breccias: Implications for the thermal history of the Fra Mauro Formation. *Meteoritics & Planetary Science* 44:1717–1734.
- Norman M. D. 2009. The Lunar Cataclysm: Reality or “Mythconception”? *Elements* 5:23–28.
- Norman M. D. and Nemchin A. A. 2014. A 4.2 billion year old impact basin on the Moon: U-Pb datation of zirconolite and apatite in lunar melt rock 67955. *Earth and Planetary Science Letters* 388:387–398.
- Norman M. D., Bennett V. C., and Ryder G. 2001. Highly siderophile (Re-PGE) and lithophile element geochemistry of Apollo 17 LKFM impact melts (abstract #1418). 32nd Lunar and Planetary Science Conference.
- Norman M. D., Duncan R. A., and Huard J. J. 2006. Identifying impact events within the lunar cataclysm from ^{40}Ar – ^{39}Ar ages and compositions of Apollo 16 impact melt rocks. *Geochimica et Cosmochimica Acta* 70:6032–6049.
- Norman M. D., Shih C.-Y., Nyquist L. E., Bogard D. D., and Taylor L. A. 2007. Early impacts on the Moon: Crystallization ages of Apollo 16 melt breccias (abstract #1991). 38th Lunar and Planetary Science Conference.
- Norman M. D., Duncan R. A. and Huard J. J. 2010. Imbrium provenance for the Apollo 16 Descartes terrain: Argon ages and geochemistry of lunar breccias 67016 and 67455. *Geochimica et Cosmochimica Acta* 74:763–783.
- Norman M.D., Taylor L.A., Shih C.-Y., and Nyquist L.E. 2016. Crystal accumulation in a 4.2 Ga lunar impact melt. *Geochimica et Cosmochimica Acta* 172:410–429.
- Orgel C., Michael G., Fassett C. I., Van der Bogert C. H., Riedel C., Kneissl T., Hiesinger H. 2018. Ancient bombardment of the inner Solar System -

- Reinvestigation of the "fingerprints" of different impactor populations on the lunar surface. *Journal of Geophysical Research* 123:748-762.
- Papanastassiou D. A., and Wasserburg G. J. 1971a. Lunar chronology and evolution from Rb-Sr studies of Apollo 11 and 12 samples. *Earth and Planetary Science Letters* 11:37-62.
- Papanastassiou D. A., and Wasserburg G. J. 1971b. Rb-Sr ages of igneous rocks from Apollo 14 mission and age of Fra Mauro formation. *Earth and Planetary Science Letters* 12:36-48.
- Papanastassiou D. A., and Wasserburg G. J. 1972a. Rb-Sr age of a Luna 16 basalt and the model age of lunar soils. *Earth and Planetary Science Letters* 13:368-374.
- Papanastassiou D. A., and Wasserburg G. J. 1972b. Rb-Sr systematics of Luna 20 and Apollo 16 samples. *Earth and Planetary Science Letters* 17:52-64.
- Puchtel I. S., Walker R. J., James O. B., and Kring D.A. 2008. Osmium isotope and highly siderophile element systematic of lunar impact melt breccia: implications for the late for the late accretion history of the Moon and Earth. *Geochimica et Cosmochimica Acta* 72:3022-3042.
- Renne P. R., Mundil R., Balco G., Min K., and Ludwig K. R. 2010. Joint determination of ^{40}K decay constants and $^{40}\text{Ar}^*/^{40}\text{K}$ for the Fish Canyon sanidine standard, and improved accuracy for $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology. *Geochimica et Cosmochimica Acta* 74:5349-5367.
- Renne P. R., Balco G., Ludwig K. R., Mundil R., and Min K. 2011. Response to the comment by W.H. Schwarz et al. on "Joint determination of ^{40}K decay constants and $^{40}\text{Ar}^*/^{40}\text{K}$ for the Fish Canyon sanidine standard, and improved accuracy for $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology" by P.R. Renne et al. (2010). *Geochimica et Cosmochimica Acta* 75:5097-5100.
- Rose H. J. Jr., Cuttitta F., Berman S., Carron M. K., Christian R. P., Dwornik E. J., Greenland L. P., and Ligon D. T. Jr. 1973. Compositional data for twenty-two Apollo 16 samples. *Proceedings of the Lunar Science Conference* 4:1149-1158.
- Ryder G. 1990. Lunar samples, lunar accretion, and the early bombardment of the Moon. *EOS, Transactions of the American Geophysical Union* 71:313-323.
- Ryder G. 1994. Coincidence in time of the Imbrium basin impact and Apollo 15 KREEP volcanic flows: The case for impact-induced melting. In Large meteorite impacts and planetary evolution, edited by B. O., Dressler, R. A. F., Grieve, and V. L., Sharpton. *Geological Society of America Special Paper* 293:11-18.
- Ryder G. and Spudis P. (1987). Chemical composition and origin of Apollo 15 impact melts. *Proc. Proceedings of the Lunar Sci. Conf. Lunar and Planetary Science Conference* 17:, E432-E446.
- Shuster D. L., Balco G., Cassata W. S., Fernandes V. A., Garrick-Bethell I., and Weiss B.P. 2010. A record of impacts preserved in the lunar regolith. *Earth and Planetary Science Letters* 290:155-165.
- Simonds C. H., Warner J. L., and Phinney W. C. 1973. Petrology of Apollo 16 poikilitic rocks. *Proceedings of the Lunar Science Conference* 4:613-632.
- Spudis P. H., and Pieters C. 1991. Global and regional data about the Moon. In *Lunar Sourcebook: A User's Guide to the Moon*, edited by G. Heiken, D. Vaniman, and B. M. French. Cambridge:Cambridge University Press. pp. 595-632.

- Steiger R. H. and Jäger E. 1977. Subcommission on geochronology: Convention on the use of decay constants in geo- and cosmochemistry. *Earth and Planetary Science Letters* 36:359–362.
- Stöffler D and Ryder G. 2001 Stratigraphy and isotope ages of lunar geologic units: Chronological standard for the inner solar system. *Space Science Reviews* 96:9–54.
- Stöffler D., Bischoff A., Borchardt R., Burghelle A., Deutsch A., Jessberger E. K., Ostertag R., Palme H., Spettel B., Reimond W. U., Wacker K., and Wänke H. 1985. Composition and evolution of the lunar crust in Descartes highlands, Apollo 16. *Proceedings of the Lunar and Planetary Science Conference* 15:C449–C505.
- Snape J. F., Nemchin A. A., Grange M. L., Bellucci J. J., Thiessen F. and Whitehouse M. J. 2016. Phosphate ages in Apollo 14 breccias: Resolving multiple impact events with high precision U–Pb SIMS analyses. *Geochimica et Cosmochimica Acta* 174:13–29.
- Stormer J. C. Jr., Pierson M. L., and Tacker R. C. 1993. Variation of F and Cl X-ray intensity due to anisotropic diffusion in apatite during electron microprobe analysis. *American Mineralogist* 78:641–648.
- Strom R. G., Malhotra R., Ito T., Yoshida F., and Kring D. A. 2005. The origin of planetary impactors in the inner solar system. *Science* 309:1847–1850.
- Swindle T. D., and Weirich J.R. 2017. The effect of partial thermal resetting on ^{40}Ar – ^{39}Ar “plateaus” (abstract #1265). 48th Lunar and Planetary Science Conference.
- Swindle T. D., Isachsen C. E., Weirich J. R., and Kring D. A. 2009. ^{40}Ar – ^{39}Ar ages of H-chondrite impact melt breccias. *Meteoritics & Planetary Science* 44:747–762.
- Swindle T. D., Kring D. A., and Weirich J. R. 2014 ^{40}Ar – ^{39}Ar ages of impacts involving ordinary chondrite meteorites. In ^{40}Ar – ^{39}Ar dating: from geochronology to thermochronology, from archaeology to planetary sciences, edited by F. Jourdan, D. Mark, and C. Verati. London: Geological Society. pp. 333–347.
- Taylor S. R., Gorton M. P., Muir P., Nance W. B., Rudowski R., and Ware N. (1973). Composition of the Descartes region, lunar highlands. *Geochimica et Cosmochimica Acta* 37:, 2665–2683.
- Tera F., Papanastassiou D., and Wasserburg G. J. 1974. Isotopic evidence for a terminal lunar cataclysm. *Earth and Planetary Science Letters* 22:1–21.
- Thiessen F., Nemchin A. A., Snape J. F., Bellucci J. J. and Whitehouse M. J. 2018. Apollo 12 breccia 12013: Impact-induced partial Pb loss in zircon and its implications for lunar geochronology. *Geochimica et Cosmochimica Acta* 230:94–111.
- Thorner C. R., and Huebner S. 1980, An experimental study of the thermal history of fragmenta-laden “basalt” 77115. Proceedings, Conference on the Lunar Highlands Crust. pp. 233–252.
- Turner G., Cadogan P. H., and Yonge C. J. 1973. Argon selenochronology. *Proceedings of the Lunar Science Conference* 4:1889–1914.
- Ulrich G. E. 1973. A geologic model for north Ray crater and stratigraphic implications for the Descartes region. *Proceedings of the Lunar Science Conference* 4:27–39.
- Wänke H., Baddenhausen H., Dreibus G., Jagoutz E., Kruse H., Palme H., Spettel B., and Teschke F. 1973 Multielement analyses of Apollo 15, 16, and 17 samples and the

- bulk composition of the moon. *Proceedings of the Lunar Science Conference* 4:1461–1481.
- Wänke H., Palme H., Kruse H., Baddenhausen H., Cendales M., Dreibus G., Hofmeister H., Jagoutz E., Palme C., Spettel B., and Teschke F. 1976 Chemistry of lunar highland rocks: A refined evaluation of the composition of the primary matter. *Proceedings of the Lunar Science Conference* 7:3479–3499.
- Warren P. H. 1993. A concise compilation of petrologic information on possibly pristine nonmare Moon rocks. *American Mineralogist* 78:360–376.
- Warren P. H. 1997. The unequal host-phase density effect in electron probe defocused beam analysis: An easily correctable problem (abstract). 28th Lunar and Planetary Science Conference. 1497–1498.
- Wieczorek M. A., and Phillips R. J. 2000. The “Procellarum KREEP Terrane”: Implications for mare volcanism and lunar evolution. *Journal of Geophysical Research* 105:20,417–20,430.
- Wilhelms D.E., McCauley J. F., and Trask N. J. 1987. The geologic history of the Moon. Washington:United States Geological Survey, Department of the Interior. United States Geological Survey Professional Paper #1348.
- Wlotzka F., Jagoutz E., Spettel B., Baddenhausen H., Balacescu A., and Wänke H. 1972. On lunar metallic particles and their contribution to the trace element content of Apollo 14 and 15 soils. *Proceedings of the Lunar Science Conference* 3:1077–1084.
- Wlotzka F., Spettel B., and Wänke H. 1973. On the composition of metal from Apollo 16 fines and meteoritic component. *Proceedings of the Lunar Science Conference* 4:1483–1491.
- Yamaguchi A., Karouji Y., Takeda H., Nyquist L. E., Borgard D. D., Ebihara M., Shih C.-Y., Reese Y. D., Garrison D., Park J., and McKay G. A. 2010. The variety of lithologies in the Yamato-86032 lunar meteorite: Implications for formation process of the lunar crust. *Geochimica et Cosmochimica Acta* 74:4507–4530.
- Zellner N.E.B. 2017. Cataclysm no more: New views on the timing and delivery of lunar impactor. *Origins of Life and Evolution of Biospheres* 47: 261–280.
- Zellner N. E. B., Delano J. W., Swindle T. D., Barra F., Olsen E., and Whittet D. C. B. 2009. Evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ age of lunar impact glasses for an increase in the impact rate ~800 Ma ago. *Geochimica et Cosmochimica Acta* 73:4590–4597.

Figure Captions

Figure 1. Apollo 16 traverse route and sampling site. Data are overlain on mosaic of LROC NAC frames M106777343RE and M106777343LE. An ancient breccia, 60016 was collected at the landing site (Station 0) on Cayley formation (Hodges, 1972; Joy et al., 2011). Norman et al. (2006) reported Ar-Ar ages from 25 Apollo 16 impact melt rocks (ample numbers and ages are showing in figure) and suggested at least 4 different impact events are required; Mafic poikilitic (3865 Ma); Aluminous subophitic (3836 Ma); Aluminous poikilitic (3771 Ma); Troctolitic vitrophyres (3859 Ma).

Figure 2. Back scattered electron images of centimeter-sized impact melt clasts in 60016 (representative areas). Clast 1, 2 and 3 have poikilitic to sub poikilitic textures. Pyroxene grains are enclosing rounded-shaped olivine and euhedral plagioclase. Clast 4, 5, and 6 have aphanitic, intergranular and porphyritic textures, respectively. Images b, d, f, h, j and l are close-up of white box areas in images a, c, e, g, i, and k, respectively. Ol: Olivine. Px: Pyroxene. Pl: Plagioclase. Opq: Opaques (metal, troilite, ilmenite, chromite).

Figure 3. Back scattered electron images of representative metal-troilite-schreibersite particles. Metal: Fe-Ni metal. Tr: Troilite. Sch: Schreibersite.

Figure 4. Ar release spectra for six impact melt clasts in 60016. Best estimated ages of clasts range from 3823 ± 75 to 4000.0 ± 23 Ma.

Figure 5. Relative Ar age probability plot for impact melt clasts in 60016. Each clast's age is plotted as a Gaussian distribution with the same total area, with the centers and spreads of the Gaussian's given by the best age and the quoted uncertainty, respectively. The age range of all clasts is from 3823 ± 75 to 4000 ± 23 Ma, generally clustering around 3900 Ma.

Figure 6. An# in plagioclase vs. Mg# in low-Ca olivine and pyroxene grains for relict minerals. Bars indicate absolute range of An# and Mg#. Compositions of plagioclase in all impact melt clast are similar and within the range of Mg-rich or FAN materials. Mg# for olivine and plagioclase is scattered among clasts. $An\# = Ca/(Na+Ca+K) \times 100$ (atomic ratio). $Mg\# = Mg/(Mg+Fe) \times 100$ (atomic ratio). Reference region for alkaline-rocks, Mg-rich rocks and FAN are obtained from previously reported values from pristine rocks (Warren 1993; Arai et al. 2008; Yamaguchi et al., 2010).

Figure 7. Principal component analysis on 9 major elements. (a) PCA can express >98% of variety of data by PC2. PCA can distinguish groups of Apollo 16 impact melt rocks as well as trace element classification (Korotev, 1994). (b) Eigen vector for

PC1 and PC2, imply Al, Fe, Mg and Ca are important element to explain compositional diversity. (c) Clast 1 and 5 in 60016 have higher values in PC1 score. (d) PC1 is good correlate with Al+Na+K (positive) and Fe+Mg (negative). (e) PC2 is correlate with Mg/Fe ratio. Compositional data are compiled from (Haskin et al., 1973; Wänke et al., 1973, 1976; Morris et al., 1986; Korotev 1994; Jolliff 1998; Norman et al., 2010)

Figure 8. Al_2O_3 (wt %) vs. K_2O , $\text{Mg}^\#$, $\text{FeO}+\text{MgO}$, and P_2O_5 plots for six impact melt clasts. Bulk compositions are estimated by the average values of defocused beam analyses (>200 points). Every impact melt clast has high- K_2O and P_2O_5 values which are similar to group 1 impact melt rocks. Reference region for impact melt groups are obtained from previously reported Apollo impact melt rocks (Albee et al., 1973; Bence et al., 1973; Duncan et al., 1973; Haskin et al., 1973; Hubbard et al., 1973; LSPET 1973; Rose et al., 1973; Simonds et al., 1973; Laul et al., 1974; Lindstrom et al., 1977; Stöffler et al., 1985; Korotev 1994; Jolliff 1998).

Table captions

Table 1. Defocused beam analyses data for impact melt clasts in 60016 (Averaged value) and reference values for Apollo 16 (A16) local regolith and Apollo 15 (A15) impact melt rocks.

Table 2. Ar-Ar ages for centimeter-sized clasts in 60016.

Supplementary materials caption

Figure S1. Optical microscope images for 6 impact melt clasts in 60016 (Transmitted light). Clast1, 2, and 3 have poikilitic texture. Clasts 4, 5 and 6 have intergranular, aphanitic or porphyritic textures.

Table S1. Representative microprobe analyses (wt. %) of silicates, oxides, and phosphate in melt clasts.

Table S2. Representative microprobe analyses (wt. %) for metal, schreibersite and troilite in melt clasts in 60016.

Table S3. Modal abundance (%) for selected areas.

Table S4. Defocused beam analysis on certified geostandard BCR-2

Table S5. Ar-Ar analysis data for 6 clasts from 60016.

90x154mm (300 x 300 DPI)

182x67mm (300 x 300 DPI)

238x178mm (300 x 300 DPI)

778x588mm (96 x 96 DPI)

209x88mm (300 x 300 DPI)

205x168mm (300 x 300 DPI)

162x181mm (300 x 300 DPI)

Table 1. Defocused beam analyses data for impact melt clasts in 60016 (Averaged value) and reference values for Apollo 16 (A16) local regolith and Apollo 15 (A15) impact melt rocks.

Sample	60016												A16	A15	
	Clast 1 (,319)		Clast 2 (,320)		Clast 3 (,324)		Clast 4 (,322)		Clast 5 (,323)		Clast 6 (,321)		Mature soil ¹	15445 ²	15455 ³
	Averaged	Corrected	Averaged	Corrected	Averaged	Corrected	Averaged	Corrected	Averaged	Corrected	Averaged	Corrected		,243	,14
SiO ₂	46.5	45.6	47.9	47.5	48.4	48.6	46.6	41.1	47.7	47.2	48.5	45.1	44.9	46.3	47.3
TiO ₂	0.65	0.98	1.14	1.62	0.64	0.7	1.53	1.45	1.04	0.93	1.55	1.53	0.59	1.8	1.35
Al ₂ O ₃	26.4	23.6	18.5	15.8	20.4	17.8	16.4	13.3	23.9	24.8	18.6	15.2	26.7	19.0	17.1
Cr ₂ O ₃	0.11	0.15	0.23	0.26	0.19	0.23	0.24	0.25	0.14	0.13	0.22	0.25	0.11	0.16	0.26
NiO	0.02	0.03	0.02	0.03	0.01	0.01	0.06	0.05	0.01	0.01	0.01	0.01	N/A	N/A	N/A
FeO	4.11	5.04	6.89	7.88	5.6	6.41	8.66	9.71	5.44	4.97	7.52	8.57	5.44	8.5	8.79
MnO	0.07	0.08	0.12	0.13	0.11	0.13	0.14	0.15	0.09	0.08	0.12	0.13	0.07	0.13	N/A
MgO	5.98	7.06	13.66	15.34	10.86	12.4	14.37	16.4	6.87	6.25	11.35	13.0	6.0	12.4	13.3
CaO	15.0	13.4	10.2	8.97	12.3	11.0	9.8	7.98	14.0	14.3	10.8	8.84	15.3	11.3	10.6
Na ₂ O	0.63	0.57	0.7	0.59	0.65	0.57	0.65	0.53	0.6	0.62	0.8	0.65	0.46	0.65	0.58
K ₂ O	0.32	0.29	0.41	0.36	0.43	0.37	0.38	0.31	0.33	0.3	0.5	0.42	0.12	0.22	0.17
P ₂ O ₅	0.41	0.44	0.41	0.39	0.45	0.43	0.51	0.48	0.39	0.34	0.36	0.39	0.12	0.48	N/A
Total	100.2		100.2		100.0		99.3		100.5		100.3		99.8	100.9	99.5

1. Data from Korotev (1997) and Lucey et al. (2006).

2. Data from Ryder and Spudis (1987).

3. Data from Taylor et al. (1973).

N/A.: Not Available

Table 2. Ar-Ar ages for centimeter-sized clasts in 60016.

Sample	Split ID	Min Age	Total Age	Plateau Age	# Steps	% Ar	Best age
,286 Clast 1							3877.5 ± 13.8
	SB16	1802.8 ± 17.6	3691.7 ± 3.3	(3870.6 ± 4.1)	4	40	
	SB17	1756.3 ± 4.5	3685.3 ± 3.3	(3891.3 ± 3.7)	6	55	
	SB18	3410.7 ± 4.0	3822.8 ± 3.2	3870.7 ± 3.5	4	53	
,290 Clast 2							3970.1 ± 12.8
	SB13	2170.2 ± 6.4	3838.8 ± 3.5	3972.5 ± 4.1	4	53	
	SB14	2576.9 ± 20.1	3881.6 ± 6.2	3980.5 ± 8.3	6	67	
	SB15	2238.4 ± 6.0	3856.0 ± 3.4	3957.3 ± 4.9	6	68	
,292 Clast 3							4000.0 ± 23.2
	SB1	2275.0 ± 125.2	3976.1 ± 4.0	(4023.2 ± 6.2)	3	24	
	SB2	1813.9 ± 23.6	3872.4 ± 7.5	(3978.8 ± 11.5)	5	35	
	SB3	1504.4 ± 10.6	3818.4 ± 4.4	(3997.7 ± 6.1)	5	34	
,305 Clast 4							3866.0 ± 3.9
	SB4	2393.7 ± 23.2	3822.2 ± 6.8	(3866.0 ± 3.9)	3	26	
,314 Clast 4							3787.2 ± 16.0
	SB8	3439.6 ± 5.3	3732.7 ± 3.4	(3803.2 ± 3.5)	6	25	
	SB9	2864.8 ± 54.3	3764.0 ± 3.3	(3771.1 ± 6.3)	4	20	
,299 Clast 5							3926.5 ± 7.3
	SB5	3702.9 ± 10.8	3941.4 ± 3.3	3926.6 ± 3.2	5	51	
	SB6	3745.0 ± 36.3	3953.2 ± 4.9	(3933.7 ± 3.2)	7	55	
	SB7	3714.62 ± 16.6	3939.5 ± 3.8	3919.2 ± 3.1	5	56	
,312 Clast 6							3906.3 ± 2.4
	SB10	2846.0 ± 47.3	3895.6 ± 3.4	3908.4 ± 3.1	4	58	
	SB11	3114.2 ± 19.4	3876.9 ± 3.6	(3906.6 ± 3.2)	9	44	
	SB12	3369.41 ± 133.3	3884.7 ± 9.6	3903.97 ± 4.5	5	60	

* Note: Under the Plateau Age column, if the apparent age plot did not produce a true plateau, we reported our best age for that split in parenthesis (#####).

Figure 1

Figure 2

Figure 2 cont.

Figure 3

Figure 4

Figure 5

Figure 6

Figure7

Figure 8