Title: Putting the ‘M’ back in maternal-fetal medicine: A five-year report card on a collaborative effort to address maternal morbidity and mortality in the U.S.

Authors:

Mary E. D’Alton MD; Vagelos College of Physicians and Surgeons, Columbia University, New York, NY

Alexander M. Friedman MD MPH; Vagelos College of Physicians and Surgeons, Columbia University, New York, New York

Peter S. Bernstein MD MPH; Albert Einstein College of Medicine, Yeshiva University, New York, New York

Haywood L. Brown MD; Morsani College of Medicine, University of South Florida, Tampa, Florida

William M. Callaghan MD MPH; Division of Reproductive Health, Centers for Disease Control and Prevention, Atlanta, Georgia

Steven L. Clark MD; Baylor College of Medicine, Baylor University, Houston, Texas

William A. Grobman MD MBA; Feinberg School of Medicine, Northwestern University, Chicago Illinois

Sarah J. Kilpatrick MD PhD; Cedars-Sinai Medical Center, Los Angeles, CA

Daniel F. O’Keeffe, MD; The Society for Maternal-Fetal Medicine, Phoenix, Arizona.

Douglas M. Montgomery MD; Kaiser Permanente Riverside Medical Center, Riverside, California

Sindhu K. Srinivas MD MSCE; Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania

George D. Wendel MD; American Board of Obstetrics and Gynecology, Dallas, Texas

Katharine D. Wenstrom MD; Women and Infants Hospital, Brown University, Providence, Rhode Island

Michael R. Foley MD; University of Arizona College of Medicine Phoenix at Banner University Medical Center Phoenix, Phoenix, Arizona.

Conflict of interests: Dr. D’Alton has received funding for ACOG II’s Safe Motherhood Initiative from Merck for Mothers. The other authors report no conflict of interest.

Corresponding author:

Mary D'Alton, MD

Division of Maternal-Fetal Fetal Medicine

Department of Obstetrics and Gynecology

Columbia University College of Physicians and Surgeons

622 West 168th Street

New York, NY 10032

Business phone: (212) 305-2377; Business fax: (212) 326-5610

Email: MD511@cumc.columbia.edu

Key words: Maternal safety, , training, certification, advanced cardiac life support, intensive care unit, simulation, clinical guidelines, bundles, levels of maternal care, postpartum hemorrhage, maternal hypertension, venous thromboembolism, research, national surveillance program, amniotic fluid embolism

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

ABSTRACT. The Centers for Disease Control and Prevention have demonstrated continuous increased risk for maternal mortality and severe morbidity with racial disparities among non-Hispanic black women an important contributing factor. More than 50,000 women experienced severe maternal morbidity in 2014 with a mortality rate of 18.0 per 100,000, higher than many other developed countries. In 2012 the first “Putting the ‘M’ back in Maternal Fetal Medicine” session was held at the Society for Maternal Fetal Medicine’s (SMFM) Annual Meeting. With the realization that rising risk for severe maternal morbidity and mortality required action, the “M in MFM” meeting identified urgent needs to: (i) enhance education and training in maternal care for maternal-fetal medicine (MFM) fellows; (ii) improve the medical care and management of pregnant women across the country; and (iii) address critical research gaps in maternal medicine. Since that first meeting a broad collaborative effort has made a number of major steps forward including the proliferation of maternal mortality review committees, advances in research, increasing educational focus on maternal critical care, and development of comprehensive clinical strategies to reduce maternal risk. Five years later, the 2017 “M in MFM” meeting served as a “report card” looking back at progress made but also looking forward to what needs to be done over the next five years given that too many mothers still experience preventable harm and adverse outcomes.

INTRODUCTION

Increasing severe maternal morbidity and mortality in the United States are well characterized. The Centers for Disease Control and Prevention have demonstrated continuous increases in maternal risk over recent decades¹⁻³ with racial disparities among non-Hispanic black women an important contributing factor.⁴⁻⁶ More than 50,000 women experienced severe maternal morbidity in 2014 with a mortality rate of 18.0 per 100,000.^{7,8} Maternal risk for a range of complications including postpartum hemorrhage, severe preeclampsia, cardiovascular and non-cardiovascular medical conditions, and peripartum hysterectomy has been stable or increased.^{1,6,9-12} In 2012 the first “Putting the ‘M’ back in Maternal Fetal Medicine” session was held at the Society for Maternal Fetal Medicine’s (SMFM) Annual Meeting. With the realization that rising risk for severe maternal morbidity and mortality^{2,13-15} required action, the “M in MFM” meeting identified urgent needs to: (i) enhance education and training in maternal care for maternal-fetal medicine (MFM) fellows; (ii) improve medical care and management of pregnant women; and (iii) address critical research gaps in maternal medicine.¹⁶ Five years later, the 2017 “M in MFM” meeting served as a “report card,” looking back at progress made and also looking forward to what needs to be done over the next five years given that too many mothers still experience preventable harm and adverse outcomes. ’

Progress on “M in MFM” initiatives has occurred within a broad collaborative effort dedicated to improving maternal care that includes the American Congress of Obstetricians and Gynecologists (ACOG), the American Board of Obstetrics and Gynecology (ABOG), SMFM, the National Partnership for Maternal Safety (NPMS) under the Council on Patient Safety in Women’s

1
2
3
4 103 Health Care, the Centers for Disease Control and Prevention (CDC), *the Eunice Kennedy Shriver*
5
6
7 104 National Institute of Child Health and Human Development (NICHD), the Alliance for Innovation on
8
9
10 105 Maternal Health, and other organizations. Leaders in these organizations participated in the
11
12 106 creation of this “report card.” In addition to reviewing progress on the original recommendations
13
14
15 107 from “Putting the ‘M’ back in maternal-fetal medicine,”¹⁶ this report card also focuses on an
16
17 108 important new development: stepwise and coordinated progress towards improving maternal
18
19
20 109 death reviews nationwide in the United States. Systematic maternal death reviews have been an
21
22 110 invaluable resource in understanding risk and improving maternal safety in the United Kingdom;¹⁵
23
24
25 111 a similar approach in United States will be critical in improving care.
26
27
28 112

31 113 **EDUCATION**

32
33
34 114 Five major educational recommendations were developed from the 2012 “M in MFM”
35
36
37 115 meeting: (i) MFM fellows should complete mandatory rotations on labor and delivery (L&D) and in
38
39
40 116 an intensive care unit (ICU), (ii) MFM fellowships should utilize simulations and case-based
41
42 117 learning to enhance education in maternal medicine, (iii) MFM should have active certification in
43
44
45 118 advanced cardiac life support (ACLS), (iv) MFM fellows should undergo a yearly in-service
46
47 119 examination similar to that administered to residents by the Council on Resident Education in
48
49
50 120 Obstetrics and Gynecology, (v) MFM fellows should receive training in leadership skills via formal
51
52 121 leadership courses as well as participation in hospital-level quality assurance committees and
53
54
55 122 quality initiative projects.
56
57

58 123 **Completed**

Mandatory L&D and ICU training – Prior to the initial “M in MFM” meeting, fellowship requirements were 12 months of clinical rotations without a requirement for L&D or ICU rotations. ABOG has subsequently modified the MFM fellowship to include 15 months of clinical rotations including 2 months of L&D/inpatient services and a 1-month ICU rotation.¹⁶ Per communication with ABOG, prior to this change, the L&D training among fellowship programs varied in structure, education, supervision, timing, and duration. In some programs, the L&D experience was unstructured and service-oriented, consisting simply of night and weekend call. All programs had to develop structured rotations with newly defined educational L&D objectives and advanced responsibilities for fellows. There has been a subsequent increase in interest in maternal care with eleven MFM physicians now certified in critical care. Many centers are pursuing the addition of a critical care fellowship as an adjunct to maternal-fetal medicine training. For the M in MFM meeting one year of MFM thesis data was reviewed: approximately 45% of MFM fellowship theses are on maternal-health topics. Per communication with ABOG, going forward data on MFM theses will be tracked.

Simulation and case-based learning – Given that maternal deaths occur rarely and many life-threatening maternal conditions are low prevalence, simulation and case-based learning represent an important means of preparing MFM providers to manage life-threatening severe morbidity scenarios.¹⁷⁻¹⁹ Obstetric simulation training may result in improved interdisciplinary teamwork and better patient outcomes with fewer medical and communication errors.²⁰ Simulation for maternal cardiac arrest has resulted in improved knowledge, confidence and competence in management of this life-threatening clinical scenario.^{21,22} The annual Critical Care in Obstetrics – An Innovative

and Integrated Model for Learning the Essentials course is held separately from SMFM's annual conference. ABOG was intimately involved in creation of this multi-day training course; ABOG worked to identify the 20 most important topics in maternal medicine and critical care obstetrics to reduce maternal morbidity and mortality. Since 2014, 1791 obstetrician-gynecologists have taken the course. The program includes online lectures, case studies, simulation sessions, as well as a 3-day onsite course with in-person simulation, virtual reality, and skill development focused on developing obstetrical critical care skills. Resident attendance at the Annual Critical Care in Obstetrics course is particularly important in maternal care education given that the majority of obstetric providers in the United States are not MFM specialists.

In-service examinations – After careful review, it was determined by ABOG that given the sample of test takers the development of in-service examinations were not psychometrically feasible for MFM fellows.

Active ACLS certification – Given that many obstetrics and gynecology residencies require ACLS training, MFM fellows often arrive at fellowship with this certification. Many programs now require maintenance of certification during fellowship.

Expanded leadership training – To address leadership gaps in MFM, over the last 5 years the Society for Maternal Fetal Medicine has sponsored the SMFM Academy for Leadership and Development. Two three-day on-site courses in Denver, Colorado with additional facilitated dialogues during the 6 interceding months focus on leadership imperatives including strength-based leadership, crucial conversations and accountability and change management. This important skill development helps focus efforts in leading change, especially as it relates to reducing maternal morbidity and mortality. Maternal care improvements require MFM-led

implementation of hospital-level care initiatives, inter-hospital collaboration, and communication with hospital and hospital-system leadership; leadership training was a gap identified by SMFM members.

CLINICAL

The original “M in MFM” meeting developed five specific recommendations for improving maternal clinical care on a national basis: (i) development of a national, stratified system for levels of maternal care with high risk patients triaged to centers with appropriate resources and expertise, (ii) creation of guidelines and bundles to prevent and manage the leading causes of severe maternal morbidity and mortality, (iii) increasing departmental and divisional support for MFM subspecialists focusing on maternal care including facilitating academic advancement for MFM inpatient directorships, (iv) improving maternal care reimbursement and billing coding for maternal care, and (v) creating models for comprehensive care for mothers at high risk secondary to acute or chronic comorbid conditions.

Completed

Levels of maternal care – The Obstetric Care Consensus on Levels of Maternal Care, developed by ACOG and SMFM, provides uniform designations for levels of maternal care, standardized definitions, quality guidelines, and nomenclature for each level of maternal care, and recommendations for equitable geographic distribution of full-service maternal care facilities and systems to improve maternal safety and outcomes across the United States. For each level of care the document makes recommendations for staffing, center capabilities, and where patients with

specific risk factors should deliver.²³ The Obstetric Care Consensus document is supported and endorsed by leading anesthesia, pediatric, and nursing societies.

In progress

Guidelines and bundles:

The National Partnership for Maternal Safety under the Council on Patient Safety in Women's Health Care, representing all major women's health care professional organizations, has published bundles on obstetric hemorrhage, severe hypertension, thromboembolism, and racial and ethnic disparities, leading causes of severe maternal morbidity and mortality.²⁴⁻²⁷ These bundles represent a selection of existing guidelines and recommendations in a form that aids implementation and consistency of practice for individual centers with the goal of improving prevention, recognition, and management of life-threatening medical and obstetric conditions. Examples of care improvement supported by these bundles include optimizing thromboembolism prophylaxis, improving recognition and management of obstetric hemorrhage, and timely treatment of severe hypertension. Additionally ACOG and SMFM have created specific management recommendations for high-risk conditions such as placenta accreta²⁸ and SMFM has focused on the role of reducing disparities in improving overall maternal outcomes.^{29,30} Facilitating statewide bundle implementation is one of the priorities of the Alliance for Innovation on Maternal Health (AIM) that has partnered with professional societies, industry, and governmental organizations to improve maternal care. The Safe Motherhood Initiative in New York State has facilitated statewide implementation of thromboembolism, obstetric hemorrhage, and hypertension bundles.³¹ Implementation of the hemorrhage bundle has led to decreased risk

for severe morbidity in California.³² Adoption of parallel recommendations for management of hypertension has led to decreased risk for eclampsia and severe morbidity.³³

Quality assessment of severe maternal morbidity

Transfusion of four or more units of blood products, admission to an intensive care unit (ICU), or both were developed at the “M in MFM” meeting as simple measures of severe maternal morbidity to prompt case reviews within all hospitals that deliver obstetrical care.³⁴ To aid in guidance of severe maternal morbidity case reviews, recommendations have been developed for a standardized interdisciplinary approach to identifying systems and professional, and facility factors with the goal of improving hospital-level safety. Included are recommendations on committee organization, the review process, medical record abstraction and assessment, review culture, data management, review timing, and review confidentiality. An accompanying structured severe maternal morbidity abstraction and assessment form to aid in morbidity reviews is available for download at: <http://safehealthcareforeverywoman.org/>.^{35,36} Uniform diagnostic criteria for amniotic fluid embolism have been developed to aid in diagnosis of this condition.³⁷ Similar to what is done by our colleagues in neonatology, severe morbidity should be tracked on large-scale, systematic bases by state and regional maternal quality care collaboratives.

Career support and academic advancement for maternal-care-focused MFM subspecialists –
Supporting maternal-care-focused MFM subspecialists is a substantial challenge, despite clear clinical needs. Although states with a high density of MFM specialists generally have lower mortality rates, several states with high MFM density have high mortality rates, suggesting that

some MFM specialists may not be providing maternal-focused care.³⁸ A significant proportion of generalist obstetricians express dissatisfaction in availability and/or the limited scope of practice of MFM specialists. Data from a recent MFM workforce survey found that only 47.5% of MFMs working full time take in-house call in hospitals providing higher acuity maternal care (Level III or IV centers).³⁹ Encouraging and supporting MFM specialists who focus on maternal care will continue to be an urgent priority given national needs.

Improving reimbursement for maternal care – Coding courses have shown great benefit in obtaining improved reimbursement for maternal care. Providers and billing staff may attend SMFM coding courses specifically for MFM coders and physicians. Departmental leadership may encourage providers and billing staff to attend these courses and billing staff to optimize coding for maternal reimbursement.

Models of comprehensive maternal care – Comprehensive maternal care centers, modeled on prenatal diagnosis/fetal referral centers, (Level IV Regional Perinatal Health Care Centers) are evolving. As an example, the Mothers Center at Columbia University serves as a quaternary referral center for maternal high-risk diagnoses such as maternal cardiac disease, placenta accreta, and other major medical, surgical and obstetric conditions. Care is coordinated with maternal-care-focused MFM providers and medical and surgical subspecialists with expertise in obstetric patients in the same clinical space; in this model the quaternary maternal center can serve as a resource to both community providers and other academic centers.

Emerging concepts – Innovation in health care delivery will play an essential role in reducing maternal risk as patient needs are often not one size fits all. For example, compared to presenting for office blood pressure checks, patient engagement of outpatient follow-up for hypertensive

diseases of pregnancy may be improved by remote patient monitoring that allows patients to monitor blood pressure at home, have their measurements digitally logged through a tablet, and then have providers automatically notified of abnormal parameters. Given that cardiac and non-cardiac medical conditions are now the leading causes of maternal death, use of innovative technological interventions to improve patient engagement and monitor high-risk conditions may represent opportunities to reduce major sources of maternal risk.⁴⁰ Aligning with initiatives to improve engagement with women in the postpartum period, remote patient monitoring along with structured “virtual visits” may represent a means of improving care for both high-risk and low-risk patients.

CRITICAL RESEARCH GAPS

The 2012 “M in MFM” meeting identified seven critical research gaps related to maternal care including: (i) the need for a standardized definition of significant maternal morbidity and “near misses,” (ii) the need for risk prediction models for severe morbidity and mortality including maternal early warning systems to avert adverse outcomes, (iii) guidance on delivery timing to optimize maternal and neonatal outcomes including management of placenta accreta, (iv) economic analyses demonstrating cost savings associated with improved maternal care, (v) research on effectiveness of maternal care training and education, (vi) research on long term outcomes after an adverse pregnancy, and, as discussed in the section below, (vii) national surveillance of maternal mortality and morbidity. Addressing these knowledge gaps is a top priority of the NICHD.

274

275 *Research supported by the NICHD*

276 The Eunice Kennedy Shriver National Institute of Child Health and Human Development

277 (NICHD) supports a broad range of maternal health research. Recent key topics in this area have

278 included advanced maternal age as it relates to maternal risk, health services and systems of care

279 in obstetrics, maternal comorbidities, depression, preeclampsia and hypertension, HIV, labor and

280 delivery complications, obstetrics pharmacology, environmental exposures, gestational diabetes,

281 obesity, and placental complications. From 2011 to 2016, maternal-focused NICHD research has

282 increased from approximately \$129 million to \$138 million annually. At the request of the NICHD,

283 the National Institutes of Health (NIH) agreed to add a maternal health category to NIH-wide

284 reporting which commenced in 2017. This new category was designed to enable better tracking of

285 “maternal health” research at the National Institutes of Health.

286 Of the seven critical research gaps in maternal health identified in 2013, NICHD has

287 supported work to address each of these areas, with examples for each highlighted in

288 **Supplementary Table 1.** In addition to specific researcher-initiated grants, NICHD supports major

289 initiatives and programs such as the National Survey of Family Growth, the Population Dynamics

290 Research Infrastructure Program, the Maternal-Fetal Medicine Units Network, the Global Network

291 for Women’s and Children’s Health, the Obstetric-Fetal Pharmacology Units Network, and the

292 Consortium on Safe Labor.^{37,41-60}

293 To simplify data-sharing with the research community, NICHD has developed and launched

294 the Data and Specimen Hub (DASH; <https://dash.nichd.nih.gov/>). This is a centralized resource for

researchers to store and access data from NICHD-funded research studies to use for secondary analyses. In addition, NICHD supports the Biospecimen Repository Access and Data Sharing (BRADS) site, a repository of epidemiologic studies and clinical trials (<https://brads.nichd.nih.gov/>).

NATIONAL SURVEILLANCE

The CDC's Pregnancy Mortality Surveillance System (PMSS) has been tracking maternal deaths in the United States since 1986. This system relies on death certificates for women thought to have died during or within one year of pregnancy as well as linked birth and/or fetal death certificates shared voluntarily from Divisions of Vital Statistics from the 50 states and the District of Columbia (<https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pmss.html>). When information on these records is viewed with a clinical eye, causes of death can be ascribed in a more nuanced fashion than when using conventional International Classification of Disease coding. However, information on death certificates is likely more accurate for certain types of deaths than others (e.g. postpartum hemorrhage versus preexisting chronic disease), and the quality of documentation may vary significantly by state (e.g. availability of records linked to birth certificates). Both under-ascertainment and false positives are a concern when surveillance is based solely on vital records.^{61,62} Furthermore, the certificate itself can rarely provide any insight into the personal or medical circumstances surrounding the death. Hence, state and regional maternal mortality review (MMR) committees are likely able to provide more particulate and accurate data based on death identification from multiple sources and these committees are well-positioned to ascertain causality based on in-depth reviews of obstetric care and circumstances of death. If surveillance and review is done for the purpose of effecting change and preventing future

1
2
3
4 317 events, it is the process information that must be understood to inform improvement. This
5
6
7 318 information can only be gleaned from in-depth review. Although currently an aspirational idea, it
8
9
10 319 is conceivable that systematic maternal death reviews on the state level across the US could begin
11
12 320 to approach the UK's Confidential Enquiries into Maternal Deaths. As of 2013 18 states and 1 city
13
14
15 321 had active MMR committees; as of today, the SMFM estimates that 36 states have MMR
16
17 322 committees with 9 states planning or starting up committees.

18
19
20 323 In supporting federal legislation to advance the national proliferation of MMR committees,
21
22
23 324 SMFM was a key partner in a coalition of national organizations including ACOG, the Association of
24
25
26 325 Maternal and Child Health Programs, the March of Dimes, the Preeclampsia Foundation and
27
28 326 others that advocated successfully for Congress to double funding for the CDC's efforts to support
29
30
31 327 MMR committees. SMFM members participated in (i) grassroots efforts to encourage Congress for
32
33 328 this support, and (ii) multiple direct education efforts with policymakers in the form of
34
35
36 329 Congressional briefings, testimony, and hearings. HR 1318, the Preventing Maternal Deaths Act, a
37
38
39 330 bill that SMFM endorsed, advocated for, hosted and cosponsored Congressional briefings for, and
40
41 331 submitted testimony to Congress on, was signed into law in December 2018. This legislation
42
43
44 332 authorizes the CDCs to provide assistance to states in setting up MMR committees and ensuring
45
46 333 standardized review and data collection processes. SMFM endorsed several state efforts to create
47
48
49 334 new or improve established MMR committees, and many MFMs serve as chairs of MMR
50
51 335 committees in their states. Additionally, the M in MFM initiative facilitated SMFM's dialogue with
52
53
54 336 the CDC to provide a role for SMFM in supporting MMR committees. In the new 2019
55
56
57 337 Congressional session, SMFM plans to support the reintroduction of other bills addressing
58
59
60
61
62
63
64
65

disparities, expanding Medicaid for postpartum women, authorizing the AIM program (which SMFM sits on the executive team for), and providing supports for perinatal quality collaboratives.

In addition to the formation of MMR committees in most states, the quality of reviews may be improving as well. Via the Alliance for Innovation on Maternal Health (AIM), approaches and “lessons learned” from mortality reviews may be broadly disseminated. The CDC is currently facilitating efforts on suggested data standards for reporting from individual MMR committees as well as providing expertise on maternal death data abstraction, informant interviews, and committee processes. The Association of Maternal Child Health Programs in partnership with the CDC Foundation and the CDC Division of Reproductive Health has recently a web-based resource, Review to Action (<http://www.reviewtoaction.org/about-us>). This effort aims to provide technical assistance to new review committees, connect existing committees with new and veteran committees, and to raise awareness of the critical roles such reviews can play in eliminating preventable deaths. The ultimate goal of improving maternal mortality surveillance in the United States is for every maternal death to undergo a high quality death review.

CONCLUSION

Since the first “M in MFM” meeting five years ago, major educational, research, and clinical strides have been made to improve maternal safety by a broad collaborative effort. Critical care and labor floor rotations are now required during MFM fellowship training. Critical care teaching has been provided to over 1700 obstetrician-gynecologists and maternal fetal medicine specialists. The NICHD has supported more than one half billion dollars in maternal research. Academic,

industry, state and federal leadership developed a number of major safety initiatives to improve maternal care.

To successfully reduce maternal mortality in the United States these achievements will have to be sustained and built upon over coming years. Models of maternal care will need to be developed to manage risk from cardiovascular and non-cardiovascular medical conditions that are now the leading causes of maternal mortality. While a number of major initiatives to improve maternal safety on the hospital level have been developed – from criteria and strategies for severe morbidity review to bundles for thromboembolism, hemorrhage, and hypertension – implementation across the more than 3,000 hospitals in the United States that provide obstetric services – in short, developing national maternal safety standards – represents a formidable challenge.⁶³ Supporting maternal-fetal medicine specialists focused on maternal safety in the setting of differential and unfavorable reimbursement relative to prenatal diagnosis and ultrasound necessitates commitment from departmental and hospital leadership. Creating a true national surveillance system that is able to critically evaluate safety gaps and disseminate lessons learned and recommendations for care improvement will rely on states developing expertise in performing high quality death reviews.

REFERENCES

1. Creanga AA, Syverson C, Seed K, Callaghan WM. Pregnancy-Related Mortality in the United States, 2011-2013. *Obstet Gynecol* 2017;130:366-73.
2. Berg CJ, Callaghan WM, Syverson C, Henderson Z. Pregnancy-related mortality in the United States, 1998 to 2005. *Obstetrics and gynecology* 2010;116:1302-9.
3. Creanga AA, Berg CJ, Syverson C, Seed K, Bruce FC, Callaghan WM. Pregnancy-related mortality in the United States, 2006-2010. *Obstetrics and gynecology* 2015;125:5-12.
4. Gyamfi-Bannerman C, Srinivas SK, Wright JD, et al. Postpartum hemorrhage outcomes and race. *American journal of obstetrics and gynecology* 2018;219:185 e1- e10.

5. Howell EA, Egorova NN, Balbierz A, Zeitlin J, Hebert PL. Site of delivery contribution to black-white severe maternal morbidity disparity. *American journal of obstetrics and gynecology* 2016;215:143-52.
6. Shahul S, Tung A, Minhaj M, et al. Racial Disparities in Comorbidities, Complications, and Maternal and Fetal Outcomes in Women With Preeclampsia/eclampsia. *Hypertens Pregnancy* 2015;34:506-15.
7. Centers for Disease Control and Prevention. Severe Maternal Morbidity in the United States. Accessed February 1, 2019 <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/severematernalmorbidity.html>.
8. Centers for Disease Control and Prevention. Pregnancy Mortality Surveillance System. Accessed February 1, 2019 <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pregnancy-mortality-surveillance-system.htm>.
9. Friedman AM, Wright JD, Ananth CV, Siddiq Z, D'Alton ME, Bateman BT. Population-based risk for peripartum hysterectomy during low- and moderate-risk delivery hospitalizations. *American journal of obstetrics and gynecology* 2016;215:640 e1- e8.
10. Bateman BT, Berman MF, Riley LE, Leffert LR. The epidemiology of postpartum hemorrhage in a large, nationwide sample of deliveries. *Anesth Analg* 2010;110:1368-73.
11. Booker WA, Ananth CV, Wright JD, et al. Trends in comorbidity, acuity, and maternal risk associated with preeclampsia across obstetric volume settings. *J Matern Fetal Neonatal Med* 2018;1-8.
12. Zelop CM, Einav S, Mhyre JM, Martin S. Cardiac arrest during pregnancy: ongoing clinical conundrum. *American journal of obstetrics and gynecology* 2018;219:52-61.
13. Clark SL. Strategies for reducing maternal mortality. *Semin Perinatol* 2012;36:42-7.
14. D'Alton ME. Where is the "M" in maternal-fetal medicine? *Obstetrics and gynecology* 2010;116:1401-4.
15. Cantwell R, Clutton-Brock T, Cooper G, et al. Saving Mothers' Lives: Reviewing maternal deaths to make motherhood safer: 2006-2008. The Eighth Report of the Confidential Enquiries into Maternal Deaths in the United Kingdom. *BJOG* 2011;118 Suppl 1:1-203.
16. D'Alton ME, Bonanno CA, Berkowitz RL, et al. Putting the "M" back in maternal-fetal medicine. *American journal of obstetrics and gynecology* 2013;208:442-8.
17. Argani CH, Eichelberger M, Deering S, Satin AJ. The case for simulation as part of a comprehensive patient safety program. *American journal of obstetrics and gynecology* 2012;206:451-5.
18. Smith A, Siassakos D, Crofts J, Draycott T. Simulation: improving patient outcomes. *Semin Perinatol* 2013;37:151-6.
19. Goffman D, Colleen L, Bernstein PS. Simulation in maternal-fetal medicine: making a case for the need. *Semin Perinatol* 2013;37:140-2.
20. Sheen JJ, Lee C, Goffman D. The utility of bedside simulation for training in critical care obstetrics. *Semin Perinatol* 2018;42:59-63.
21. Adams J, Cepeda Brito JR, Baker L, et al. Management of Maternal Cardiac Arrest in the Third Trimester of Pregnancy: A Simulation-Based Pilot Study. *Crit Care Res Pract* 2016;2016:5283765.
22. Fisher N, Eisen LA, Bayya JV, et al. Improved performance of maternal-fetal medicine staff after maternal cardiac arrest simulation-based training. *Am J Obstet Gynecol* 2011;205:239 e1-5.
23. Obstetric Care Consensus No. 2: Levels of maternal care. *Obstetrics and gynecology* 2015;125:502-15.
24. Main EK, Goffman D, Scavone BM, et al. National Partnership for Maternal Safety: Consensus Bundle on Obstetric Hemorrhage. *Obstetrics and gynecology* 2015;126:155-62.
25. D'Alton ME, Friedman AM, Smiley RM, et al. National Partnership for Maternal Safety: Consensus Bundle on Venous Thromboembolism. *Obstetrics and gynecology* 2016;128:688-98.
26. Bernstein PS, Martin JN, Jr., Barton JR, et al. National Partnership for Maternal Safety: Consensus Bundle on Severe Hypertension During Pregnancy and the Postpartum Period. *Obstetrics and gynecology* 2017;130:347-57.

27. Howell EA, Brown H, Brumley J, et al. Reduction of Peripartum Racial and Ethnic Disparities: A Conceptual Framework and Maternal Safety Consensus Bundle. *Obstetrics and gynecology* 2018;131:770-82.
28. Society of Gynecologic O, American College of O, Gynecologists, et al. Placenta Accreta Spectrum. *Am J Obstet Gynecol* 2018;219:B2-B16.
29. Moroz L, Riley LE, D'Alton M, et al. SMFM Special Report: Putting the "M" back in MFM: Addressing education about disparities in maternal outcomes and care. *Am J Obstet Gynecol* 2018;218:B2-B8.
30. Jain JA, Temming LA, D'Alton ME, et al. SMFM Special Report: Putting the "M" back in MFM: Reducing racial and ethnic disparities in maternal morbidity and mortality: A call to action. *Am J Obstet Gynecol* 2018;218:B9-B17.
31. Chazotte C, D'Alton ME. Maternal mortality in New York--Looking back, looking forward. *Semin Perinatol* 2016;40:132-5.
32. Main EK, Cape V, Abreo A, et al. Reduction of severe maternal morbidity from hemorrhage using a state perinatal quality collaborative. *American journal of obstetrics and gynecology* 2017;216:298 e1- e11.
33. Shields LE, Wiesner S, Klein C, Pelletreau B, Hedriana HL. Early standardized treatment of critical blood pressure elevations is associated with a reduction in eclampsia and severe maternal morbidity. *Am J Obstet Gynecol* 2017;216:415 e1- e5.
34. Callaghan WM, Grobman WA, Kilpatrick SJ, Main EK, D'Alton M. Facility-based identification of women with severe maternal morbidity: it is time to start. *Obstetrics and gynecology* 2014;123:978-81.
35. Kilpatrick SJ, Berg C, Bernstein P, et al. Standardized severe maternal morbidity review: rationale and process. *Obstetrics and gynecology* 2014;124:361-6.
36. Obstetric Care Consensus No. 5: Severe Maternal Morbidity: Screening and Review. *Obstetrics and gynecology* 2016;128:e54-60.
37. Clark SL, Romero R, Dildy GA, et al. Proposed diagnostic criteria for the case definition of amniotic fluid embolism in research studies. *American journal of obstetrics and gynecology* 2016;215:408-12.
38. Sullivan SA, Hill EG, Newman RB, Menard MK. Maternal-fetal medicine specialist density is inversely associated with maternal mortality ratios. *American journal of obstetrics and gynecology* 2005;193:1083-8.
39. Wenstrom K, Erickson K, Schulkin J. Are obstetrician-gynecologists satisfied with their maternal-fetal medicine consultants? A survey. *Am J Perinatol* 2012;29:599-608.
40. Hirshberg A, Bittle MD, VanDerTuyn M, Mahraj K, Asch DA, Rosin R, Bennett I, Srinivas SK. Rapid-Cycle Innovation Testing of Text-Based Monitoring for Management of Postpartum Hypertension. *Journal of Clinical Outcomes Management*. 2017 24 (2): 77-85.
41. MacDorman MF, Declercq E, Cabral H, Morton C. Recent Increases in the U.S. Maternal Mortality Rate: Disentangling Trends From Measurement Issues. *Obstetrics and gynecology* 2016;128:447-55.
42. Grobman WA, Bailit JL, Rice MM, et al. Racial and ethnic disparities in maternal morbidity and obstetric care. *Obstetrics and gynecology* 2015;125:1460-7.
43. Belanoff C, Declercq ER, Diop H, et al. Severe Maternal Morbidity and the Use of Assisted Reproductive Technology in Massachusetts. *Obstetrics and gynecology* 2016;127:527-34.
44. Wallace ME, Hoyert D, Williams C, Mendola P. Pregnancy-associated homicide and suicide in 37 US states with enhanced pregnancy surveillance. *American journal of obstetrics and gynecology* 2016;215:364 e1- e10.
45. Grobman WA, Bailit JL, Rice MM, et al. Frequency of and factors associated with severe maternal morbidity. *Obstetrics and gynecology* 2014;123:804-10.
46. Belfort MA, Saade GR, Thom E, et al. A Randomized Trial of Intrapartum Fetal ECG ST-Segment Analysis. *N Engl J Med* 2015;373:632-41.
47. Reddy UM, Rice MM, Grobman WA, et al. Serious maternal complications after early preterm delivery (24-33 weeks' gestation). *American journal of obstetrics and gynecology* 2015;213:538 e1-9.

48. Coviello EM, Grantz KL, Huang CC, Kelly TE, Landy HJ. Risk factors for retained placenta. *American journal of obstetrics and gynecology* 2015;213:864 e1- e11.
49. Downes KL, Hinkle SN, Sjaarda LA, Albert PS, Grantz KL. Previous prelabor or intrapartum cesarean delivery and risk of placenta previa. *American journal of obstetrics and gynecology* 2015;212:669 e1-6.
50. Boghossian NS, Yeung E, Mendola P, et al. Risk factors differ between recurrent and incident preeclampsia: a hospital-based cohort study. *Annals of epidemiology* 2014;24:871-7e3.
51. Arkema EV, Palmsten K, Sjowall C, Svenungsson E, Salmon JE, Simard JF. What to Expect When Expecting With Systemic Lupus Erythematosus (SLE): A Population-Based Study of Maternal and Fetal Outcomes in SLE and Pre-SLE. *Arthritis Care Res (Hoboken)* 2016;68:988-94.
52. McCall-Hosenfeld JS, Phiri K, Schaefer E, Zhu J, Kjerulff K. Trajectories of Depressive Symptoms Throughout the Peri- and Postpartum Period: Results from the First Baby Study. *J Womens Health (Larchmt)* 2016;25:1112-21.
53. Lannon SM, Guthrie KA, Vanderhoeven JP, Gammill HS. Uterine rupture risk after periviable cesarean delivery. *Obstetrics and gynecology* 2015;125:1095-100.
54. Frolova AI, Stout MJ, Tuuli MG, Lopez JD, Macones GA, Cahill AG. Duration of the Third Stage of Labor and Risk of Postpartum Hemorrhage. *Obstetrics and gynecology* 2016;127:951-6.
55. Kawakita T, Reddy UM, Iqbal SN, et al. Duration of Oxytocin and Rupture of the Membranes Before Diagnosing a Failed Induction of Labor. *Obstetrics and gynecology* 2016;128:373-80.
56. Grobman WA, Bailit J, Lai Y, et al. Association of the Duration of Active Pushing With Obstetric Outcomes. *Obstetrics and gynecology* 2016;127:667-73.
57. Williams K, Sassler S, Addo F, Frech A. First-birth Timing, Marital History, and Women's Health at Midlife. *J Health Soc Behav* 2015;56:514-33.
58. Cirillo PM, Cohn BA. Pregnancy complications and cardiovascular disease death: 50-year follow-up of the Child Health and Development Studies pregnancy cohort. *Circulation* 2015;132:1234-42.
59. Gunderson EP, Hurston SR, Ning X, et al. Lactation and Progression to Type 2 Diabetes Mellitus After Gestational Diabetes Mellitus: A Prospective Cohort Study. *Ann Intern Med* 2015;163:889-98.
60. Bateman BT, Franklin JM, Bykov K, et al. Persistent opioid use following cesarean delivery: patterns and predictors among opioid-naïve women. *American journal of obstetrics and gynecology* 2016;215:353 e1- e18.
61. Clark SL, Christmas JT, Frye DR, Meyers JA, Perlin JB. Maternal mortality in the United States: predictability and the impact of protocols on fatal postcesarean pulmonary embolism and hypertension-related intracranial hemorrhage. *American journal of obstetrics and gynecology* 2014;211:32 e1-9.
62. Creanga AA, Callaghan WM. Recent Increases in the U.S. Maternal Mortality Rate: Disentangling Trends From Measurement Issues. *Obstetrics and gynecology* 2017;129:206-7.
63. Simpson KR. An overview of distribution of births in United States hospitals in 2008 with implications for small volume perinatal units in rural hospitals. *J Obstet Gynecol Neonatal Nurs* 2011;40:432-9.

Supplementary Table 1. Examples of NICHD-funded research on maternal outcomes**National surveillance**

MacDorman MF, Declercq E, Cabral H, Morton C. Recent Increases in the U.S. Maternal Mortality Rate: Disentangling Trends From Measurement Issues. *Obstetrics and gynecology* 2016;128:447-55.

Grobman WA, Bailit JL, Rice MM, et al. Racial and ethnic disparities in maternal morbidity and obstetric care. *Obstetrics and gynecology* 2015;125:1460-7.

Belanoff C, Declercq ER, Diop H, et al. Severe Maternal Morbidity and the Use of Assisted Reproductive Technology in Massachusetts. *Obstetrics and gynecology* 2016;127:527-34.

Wallace ME, Hoyert D, Williams C, Mendola P. Pregnancy-associated homicide and suicide in 37 US states with enhanced pregnancy surveillance. *Am J Obstet Gynecol* 2016;215:364 e1- e10.

Severe morbidity

Clark SL, Romero R, Dildy GA, et al. Proposed diagnostic criteria for the case definition of amniotic fluid embolism in research studies. *Am J Obstet Gynecol* 2016;215:408-12.

Grobman WA, Bailit JL, Rice MM, et al. Frequency of and factors associated with severe maternal morbidity. *Obstetrics and gynecology* 2014;123:804-10.

Belfort MA, Saade GR, Thom E, et al. A Randomized Trial of Intrapartum Fetal ECG ST-Segment Analysis. *N Engl J Med* 2015;373:632-41.

Reddy UM, Rice MM, Grobman WA, et al. Serious maternal complications after early preterm delivery (24-33 weeks' gestation). *Am J Obstet Gynecol* 2015;213:538 e1-9.

Risk prediction

Coviello EM, Grantz KL, Huang CC, Kelly TE, Landy HJ. Risk factors for retained placenta. *Am J Obstet Gynecol* 2015;213:864 e1- e11.

Downes KL, Hinkle SN, Sjaarda LA, Albert PS, Grantz KL. Previous prelabor or intrapartum cesarean delivery and risk of placenta previa. *Am J Obstet Gynecol* 2015;212:669 e1-6.

Boghossian NS, Yeung E, Mendola P, et al. Risk factors differ between recurrent and incident preeclampsia: a hospital-based cohort study. *Ann Epidemiol* 2014;24:871-7e3.

Arkema EV, Palmsten K, Sjowall C, Svenungsson E, Salmon JE, Simard JF. What to Expect When Expecting With Systemic Lupus Erythematosus (SLE): A Population-Based Study of Maternal and Fetal Outcomes in SLE and Pre-SLE. *Arthritis Care Res (Hoboken)* 2016;68:988-94.

McCall-Hosenfeld JS, Phiri K, Schaefer E, Zhu J, Kjerulff K. Trajectories of Depressive Symptoms Throughout the Peri- and Postpartum Period: Results from the First Baby Study. *J Womens Health (Larchmt)* 2016;25:1112-21.

Lannon SM, Guthrie KA, Vanderhoeven JP, Gammill HS. Uterine rupture risk after periviable cesarean delivery. *Obstetrics and gynecology* 2015;125:1095-100.

Optimal delivery timing

Frolova AI, Stout MJ, Tuuli MG, Lopez JD, Macones GA, Cahill AG. Duration of the Third Stage of Labor and Risk of Postpartum Hemorrhage. *Obstetrics and gynecology* 2016;127:951-6.

Kawakita T, Reddy UM, Iqbal SN, et al. Duration of Oxytocin and Rupture of the Membranes Before Diagnosing a Failed Induction of Labor. *Obstetrics and gynecology* 2016;128:373-80.

Grobman WA, Bailit J, Lai Y, et al. Association of the Duration of Active Pushing With Obstetric Outcomes. *Obstetrics and gynecology* 2016;127:667-73.

Long term outcomes

Williams K, Sassler S, Addo F, Frech A. First-birth Timing, Marital History, and Women's Health at Midlife. *J Health Soc Behav* 2015;56:514-33.

Cirillo PM, Cohn BA. Pregnancy complications and cardiovascular disease death: 50-year follow-up of the Child Health and Development Studies pregnancy cohort. *Circulation* 2015;132:1234-42.

Gunderson EP, Hurston SR, Ning X, et al. Lactation and Progression to Type 2 Diabetes Mellitus After Gestational Diabetes Mellitus: A Prospective Cohort Study. *Ann Intern Med* 2015;163:889-98.

Bateman BT, Franklin JM, Bykov K, et al. Persistent opioid use following cesarean delivery: patterns and predictors among opioid-naïve women. *Am J Obstet Gynecol* 2016;215:353 e1- e18.

NICHD, Eunice Kennedy Shriver National Institute of Child Health and Human Development.