Association of *UCP1*, *UCP2* and *UCP3* gene polymorphisms with cardiovascular disease risk factors in European adolescents: the HELENA study

Cite this article as: Jose M. Pascual-Gamarra, Diego Salazar-Tortosa, Idoia Labayen, Azahara I. Rupérez, Catherine Leclercq, Ascension Marcos, Sonia Gómez, Luis A. Moreno, Aline Meirhaeghe, Manuel J Castillo and Jonatan R Ruiz, Association of *UCP1*, *UCP2* and *UCP3* gene polymorphisms with cardiovascular disease risk factors in European adolescents: the HELENA study, *Pediatric Research* doi:10.1038/s41390-019-0735-7

This Author Accepted Manuscript is a PDF file of an unedited peer-reviewed manuscript that has been accepted for publication but has not been copy edited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. <https://www.nature.com/authors/policies/license.html#AAMtermsV1>

Association of *UCP1*, *UCP2* and *UCP3* gene polymorphisms with cardiovascular disease risk factors in European adolescents: the HELENA study

Jose M. Pascual- Gamarra^{1*}, Diego Salazar-Tortosa^{2*}, Idoia Labayen³, Azahara I. Rupérez⁴, Catherine Leclercq⁵, Ascension Marcos⁶, Sonia Gómez⁶, Luis A. Moreno⁴, Aline Meirhaeghe⁷, Manuel J Castillo⁸, Jonatan R Ruiz^{1,9}

Manuscript type: Clinical Research Article

Author contributorship statement: "IL, AIR, CL, AM, SG, LAM, AM MCG, DST, JMPG and JRR designed the study; DST performed all analyses; JM, DST and JRR wrote the initial draft and all co-authors significantly contributed to the final version".

*Equally contributed

¹PROFITH “PROmoting FITness and Health through physical activity” research group, Sport and Health University Research Institute (iMUDS), Department of Physiology, Faculty of Medicine, University of Granada, Granada, Spain.

²Department of Ecology, Faculty of Sciences, University of Granada, Spain.

³Department of Health Sciences, Public University of Navarra, Pamplona, Spain

⁴Department of Health and Human Performance, School of Health Sciences, University of Zaragoza, Zaragoza, Spain. GENUUD "Growth, Exercise, Nutrition and Development" Research Group, Zaragoza, Spain

⁵CREA (Council for Agricultural Research and Economics) - Research Center for Food and Nutrition, Rome, Italy.

⁶Immunonutrition Group, Institute of Food Science, Technology and Nutrition (ICTAN), Spanish National Research Council (CSIC), Madrid, Spain; Centro de Investigación Biomédica en Red de Fisiopatología de la Obesidad y Nutrición (CIBEROBn), Spain

⁷Inserm, Institut Pasteur de Lille, Univ. Lille, UMR1167-RID-AGE-Risk factors and molecular determinants of aging-related diseases, Lille, France

⁸PROFITH “PROmotingFITness and Healththroughphysicalactivity” research group. Department of Physical Education and Sport, Faculty of Sport Sciences, University of Granada, Spain.

⁹Dep. of Biosciences and Nutrition at NOVUM, Karolinska Institutet, Huddinge, Sweden

Corresponding author: Jose Miguel Pascual Gamarra

Address: Department of Physiology, Faculty of Medicine, University of Granada, Avda. de la Investigación 11, 18016, Granada, Spain

Phone number: +34 655305348

Email address: jmpasc90@gmail.com

Funding/Support: The HELENA Study is supported by contract FOOD-CT-2005-007034 from the European Community Sixth RTD Framework Programme. The study was supported by the Spanish Ministry of Education (FPU16/02127, FPU13/03410), by the University of Granada *Plan Propio de Investigación 2016* (Excellence actions: Unit of Excellence on Exercise and Health [UCEES]), by the *Junta de Andalucía, Consejería de Conocimiento, Investigación y Universidades*, by the European Regional Development Fund (ERDF), ref. SOMM17/6107/UGR and by

Redes Temáticas de Investigación Cooperativa RETIC grant Red SAMID
RD16/0022.

ABSTRACT

Introduction: Cardiovascular diseases (CVDs) are responsible of 31% of all deaths worldwide. Genetic predisposition to CVDs in adolescents remains largely unknown.

Objective: To examine the association of *UCP1*, *UCP2* and *UCP3* gene polymorphisms with CVDs risk factors in European adolescents.

Study design: A cross-sectional study that involves 1.057 European adolescents (12-18 years old) from the HELENA Study. A total of 18 polymorphisms of *UCP1*, *UCP2* and *UCP3* genes were genotyped. We measured serum total cholesterol, HDL, ApoA1, ApoB, leptin, triglycerides, glucose, insulin, blood pressure, and calculated HOMA (homeostatic model assessment) and a CVD Risk Score.

Results: The G allele of *UCP2* rs2735572 and T allele of *UCP2* rs17132534 were associated with higher diastolic blood pressure ($P=0.001$; $FDR=0.009$ and $P=8e-04$; $FDR=0.009$, respectively). We observed that the AATAG haplotype of *UCP1* was associated with higher serum ApoB/ApoA1 ($P= 0.008$; $FDR = 0.031$) and ApoB levels ($P= 0.008$; $FDR = 0.031$). Moreover, the ACC haplotype of *UCP3* was associated with a higher CVD risk score ($P= 0.0036$; $FDR = 0.01$).

Conclusions: Two *UCP2* polymorphisms and haplotypes of *UCP1* and *UCP3* were associated with CVD risk factors. These findings suggest that *UCPs* may have a role in the development of CVD already in adolescents.

Introduction

Cardiovascular diseases (CVDs) are the main cause of premature death and chronic disability worldwide (1). CVD events occur most frequently during or after the fifth decade of life, however, there is evidence indicating that the precursors of CVD have its origin in the first decades of life (2). Therefore, prevention is fundamental to reduce the incidence of these pathologies, especially in young people.

CVDs are a result of complex harmful interactions between environmental and genetic risk factors. Environmental factors include unhealthy diet, tobacco use or physical inactivity (3). However, efforts have been insufficient to decrease the prevalence and new pathogenic dimensions come into play. Recent studies have described the association between some single nucleotide polymorphisms (SNPs) with myocardial infarction (4) and other cardiovascular complications (5). For example, uncoupling protein genes (UCPs) have been associated with risk factors of cardiovascular disease such as prediabetes and type 2 diabetes mellitus (T2DM) (6), overweight and obesity (7,8), plasma levels of cholesterol (9) or hypertension (HT) (10), mainly in adults over 50 years of age.

The most studied UCP genes are: i) *UCP1*, which main function is heat production through non-shivering thermogenesis in brown adipose tissue (BAT) (11); ii) *UCP2*, in addition to may have a regulating role in thermogenesis of BAT, seems to be involved in the control of reactive oxygen species (ROS) production (12,13), the modulation of insulin secretion (14) and the regulation of mitochondrial fatty acid oxidation (15), and iii) *UCP3*, which role has been related to the coupling regulation of mitochondrial respiration in skeletal muscle mitochondria (16) and fatty acids oxidation (17), and is a mediator of thermogenesis (18). The playing role of uncoupling proteins in human physiology makes UCPs ideal targets against

cardiovascular-associated pathologies. Indeed, several SNPs of *UCP2* (rs660339, rs659366) (19,20) and *UCP3* (rs2075577, rs3781907, rs1800006, rs1800849) (19,21–23) have been associated with T2DM, overweight/obesity, serum total and LDL-cholesterol and others cardiovascular risk markers. Nevertheless, poor evidence of CVD risk factors, especially in youth has been described.

Data obtained within the Healthy Lifestyle in Europe by Nutrition in Adolescence Cross-Sectional Study (HELENA-CSS) study provide an excellent opportunity to study the association of *UCP1*, *UCP2* and *UCP3* SNPs with CVD risk factors in European adolescents. The HELENA study was designed to provide reliable data on nutrition and health-related variables in a relatively large sample of European adolescents from 9 different countries and includes information on 18 SNPs of *UCP1*, *UCP2* and *UCP3* genes as well as a number of CVD risk factors. To our knowledge, UCPs polymorphisms have not been identified in GWAS of body weight or body composition in adults or other age groups

The aim of this study was therefore to examine the association of 18 *UCP1*, *UCP2* and *UCP3* SNPs (see Table 1) with CVD risk factors in European adolescents.

Material and methods

Participants

The HELENA-CSS attempted to report the lifestyle and nutritional status of European adolescents. A total of 3865 participants (12-18 year old) of nine European countries (Greece, Germany, Belgium, France, Hungary, Italy, Sweden, Austria and Spain) were selected to be part of this study. They were randomly selected from public and private schools in each city between October 2006 and December 2007. We collected blood samples of one-third of these participants (N=1155) with the consequent genetic analysis and clinical biochemistry assays. Finally, 1057 (552 girls) adolescents with UCPs SNPs and CVD risk factors data were included in this study. Adolescents and corresponding parents/guardians were fully informed about aims and methods of the study such as inclusion criteria (24,25), and signed an informed written consent. Ethical guidelines of the Declaration of Helsinki 1964 (revision of Edinburgh 2000), Good Clinical Practice, and legislation about clinical research in humans in each of the participating countries were respected by the study. Human research committees of each center involved approved the protocol (26).

Assessment of Cardiovascular Risk Factors

A total of 30 ml of blood samples were drawn after a 10-h overnight fast at school between 0830 and 0900 hours following a standardized blood collection protocol. Serum/plasma was centrifuged directly at the schools at 3500 rpm (for 15 min, at room temperature). After centrifugation they were stored and transported (4-7°C) to the central laboratory (Bonn, Germany) where they were deposited at -80°C as explained in detailed elsewhere (27). Serum concentrations of cardiovascular risk factors were measured in centralized laboratories.

The CVD risk factors analysed included serum total cholesterol, high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL), ApoA1, ApoB, leptin, triglycerides and glucose, which were measured on the Dimension RxL clinical chemistry system (Dade Behring, Schwalbach, Germany) with enzymatic methods. Insulin was measured by a solid-phase two-site chemiluminescent immunometric assay with an Immulite 2000 analyzer (DPC Biermann GmbH, Bad Nauheim, Germany). Homeostasis model assessment (HOMA) was calculated ($[\text{glycaemia} \times \text{insulin}] / 22.5$) as resistance to insulin indicator, along with the Quantitative Insulin Sensitivity Check Index (QUICKI), which was calculated as $1 / [\log(\text{insulin}) + \log(\text{glycaemia})]$. Blood pressure was measured with an automatic oscillometric device (OMRON M6). Adolescents quietly sat for 5 min before the measurements, conducted on the right arm in an extended position. Two measures of diastolic and systolic blood pressure (DBP and SBP, respectively) were taken 5 min apart, and the mean of both values (in mmHg) was used in analyses.

We computed a CVD risk score with the mean of the standardized value $[(\text{value} - \text{mean}) / \text{standard deviation}]$ of the following variables: Total cholesterol/HDL, triglycerides, HOMA, systolic blood pressure, and triceps and subscapular skinfolds (28). The characteristics of the study sample, including CVD risk factors, are shown in Table 2.

Genotyping

Blood for DNA extraction was collected in EDTA K3 tubes, stored at the Analytical Laboratory at the University of Bonn, and then sent to the Genomic Analysis Laboratory at the Institut Pasteur de Lille (Lille, France). DNA was extracted from

white blood cells with the Puregene kit (QIAGEN, Courtaboeuf, France) and stored at 20°C. Samples were genotyped by an Illumina System (Illumina, Inc, San Diego, California) and the software used was GoldenGate (Inc, San Francisco, California). High rate of genotyping success was performed ($\geq 97,8\%$) and each polymorphism respected the Hardy-Weinberg equilibrium ($P > 0.2$ in all cases; Table 1). Several polymorphisms of the same genes showed linkage disequilibrium between them (Figures 1 and 2).

Statistical analysis

Deviations from Hardy-Weinberg equilibrium (HWE) were determined by means of an exact test and considering a p value of 0.05 as a threshold. Associations between genetic markers and CVD risk factors were assessed through linear models. Five inheritance models (dominant, recessive, log-additive, codominant and overdominant) were used for all analyses, except in those where rs2071416, rs2735572 and rs17132534 SNPs were involved. These polymorphisms were analysed using only a dominant model due to the low number of minor homozygotes (minor allele frequency < 0.1 ; Table 1). Previous findings shown the association between non-additive models with UCPs, which indicates the interest of perform this five models and compare the additive models with non-additives ones (29). Adjustment variables were body mass index (calculated as weight in kilograms divided by height in meters squared), age, gender and center. For each SNP, P values were computed using the likelihood ratio test (LRT) between a model with the polymorphism and a null model without it. These analyses were performed with the “SNPassoc” R package (30). We considered the associations between all SNPs and each phenotype under a given heritage model such as the family test, i.e. the number of tests were equal to the

number of SNPs analysed for a given phenotype. We selected the significant genotype-phenotype associations to perform haplotype analysis, i.e. only SNPs and phenotypes significant associated were considered for next analyses. Given the exploratory nature of these analyses and the reduced number of independent tests (markers are in linkage disequilibrium), the Bonferroni correction could be too conservative (31). Instead of this method, we performed an exploratory selection of associations using an approach that controls the expected proportion of false positives (False Discovery Rate [FDR]) (32). Therefore, associations with $FDR < 0.1$ were used in haplotype analyses.

Linkage disequilibrium between polymorphisms and haplotype block structures were evaluated with Haploview 4.2 (<http://www.broad.mit.edu/mpg/haploview>) and the R package haplo.stats (33). First, haplotype blocks were generated by the algorithm of four-gamete rules (34) using Haploview. For each block, we tested if the observed frequencies of haplotypes were deviated from those expected under linkage equilibrium using “haplo.em” {haplo.stats} ($P = 0$ in all blocks). Finally, we assessed the association between haplotypes and phenotypes by means of a permutation procedure performed with haplo.score {haplo.stats}. Only additive and dominant models were considered given the low frequency of some haplotypes. For those significant associations we performed regressions between haplotypes and phenotypes with the purpose of testing significant differences between haplotype levels. These regressions were performed with haplo.glm {haplo.stats}. Again, the FDR was calculated from the P values for differences between the reference haplotype (the most frequent) and other haplotypes.

Results

Association between UCP polymorphisms and CVD Risk Factors

Two *UCP2* SNPs were individually associated with CVD risk factors after multiple-comparison corrections with a FDR<0.05 threshold. Associations with FDR<0.1 were also selected as a screening method to further haplotype analyses (Figure 3). All individual comparisons between genotypes and CVD risk factors are presented in supplementary file. We observed that the G minor allele of rs2735572 SNP and the T minor allele of rs17132534 SNP were associated with higher DBP under a dominant inheritance model (beta coefficient=0.04, P=0.001; FDR=0.009 and beta coefficient=0.04, P=8e-04; FDR=0.009 respectively; beta coefficients obtained from models with the response variable log transformed; Figure 3).

Association between UCP polymorphism haplotypes and CVD risk factors

UCP1 block contains the rs12502572, rs11932232, rs6822807, rs6818140, and rs2071416 SNPs (Figure 1). The AATAG haplotype of *UCP1* was significantly associated with a higher ApoB/ApoA1 ratio than the reference GATAT haplotype (global P=0.046; difference between groups =0.06; 95CI = 0.02 - 0.10; P= 0.008; FDR = 0.031; under additive model; differences between groups obtained from models with the response variable log transformed, respectively). The AATAG haplotype was also associated with higher ApoB levels than the reference GATAT haplotype (global P=0.045; difference between groups =0.06; 95CI = 0.02 - 0.10; P= 0.008; FDR = 0.031; under additive model). Also an association between the *UCP3* block 2 (rs7930460, rs2075577 and rs2734828, Figure 2) and the risk score was observed. The ACC haplotype was associated with a higher risk score compared with

the ATC haplotype (global $P=0.008$; difference between groups= 0.06; 95CI = 0.02 - 0.10; $P= 0.0036$; FDR = 0.01; under dominant model).

Discussion

We observed that the *UCP2* G and T alleles of the rs2735572 and rs17132534 SNPs were associated with higher diastolic blood pressure in European adolescents. Moreover, we found that the *UCP1* AATAG haplotype was associated with higher serum ApoB/ApoA1 and ApoB levels. Finally, the *UCP3* ACC haplotype was associated with a higher CVD risk score. Taken together, these findings suggest that UCPs may have an important role in cardiovascular health already in the first decades of life.

To our knowledge, this is the first study investigating the association between UCPs and CVD risk factors in European adolescents. Several studies with smaller sample sizes (6,21,35,36) reported an association of single polymorphisms of *UCP3* SNPs or combined haplotypes of *UCP2/UCP3* with several CVD risk factors such as TC and LDL levels, insulin or HOMA in adults. In contrast, we observed no association of these UCPs SNPs with CVD risk factors in European adolescents. These discordances may be due to the fact that they are population dependent, with possibly different allele frequencies and penetrance in these populations. Also differences in age, inter-country differences in lifestyle behaviours and sample sizes are important and could lead to differences across studies.

A plausible mechanism to partially explain the observed associations is that polymorphisms or haplotypes of UCPs could alter UCP functions and predispose to cardiovascular risk or an increased Risk Score (due to it is made up of cardiovascular risk factors). This dysfunction may explain the phenotypes observed with CVD risk

through, i) dysfunction of the process of oxidation of fatty acids, leading to altered serum lipid levels as TC, LDL, HDL or TG (9,15), ii) in relation to ROS regulation mediated by UCP2; Pierelli *et al.* (37) showed that knockout mice deletion of the *UCP2* gene contributes to atherosclerosis lesion development and a significantly shorter lifespan. Several studies in humans and cultured cells suggested that excessive ROS production is involved in the atherosclerotic plaque formation and progression (38,39). Therefore, evidence suggests that decreasing ROS production is a remarkable target to prevent the atherosclerotic process. *UCP2* negatively regulates intracellular ROS production (12,13) making it a potential therapeutic target for the treatment of vascular diseases, iii) UCPs have been also related to blood pressure control. Dhamrait *et al.* (40) described the role of UCPs in the regulation of angiotensin-converting enzyme (ACE). This enzyme plays a pivotal component of the endocrine renin–angiotensin system (RAS), also playing a key role in the regulation of the human circulation. ACE favours the rise of angiotensin II and aldosterone, leading to salt and water retention by the kidney and to constriction of small blood vessels in the arterial tree. Taken together, these actions serve to elevate blood pressure. Moreover, Dhamrait *et al.* (40) showed that some *UCP3* and *UCP2* SNPs were associated with higher age-adjusted ACE activity, which could contribute to a hypertension status, which is also consistent with our findings. Cardiovascular diseases are pathologies with a long-term latency period and modifiable cardiovascular risk factors, which makes prevention fundamental, especially in these populations with genetic predisposition.

A limitation of our study is its cross-sectional nature. Our results should be considered carefully and studies with larger sample size could help to further confirm this possible genetic predisposition.

In conclusion, we observed an association between the *UCP2* rs2735572 and rs17132534 SNPs with higher diastolic blood pressure in adolescents from nine European countries. We also observed a haplotype association of *UCP1* and *UCP3* with higher blood apolipoproteins levels and risk score, respectively. These findings suggest that *UCPs* may have an important role in the development of CVD predisposition already in European adolescents.

References

1. Roth GA, Johnson C, Abajobir A, Abd-Allah F, Abera SF, Abyu G, et al. Global, Regional, and National Burden of Cardiovascular Diseases for 10 Causes, 1990 to 2015. *J Am Coll Cardiol*. 2017;70(1):1–25.
2. Shah S. Primary prevention of Cardiovascular Disease. *World Heal Organ*. 2011;364:937.
3. World Health Organization. Global status report on noncommunicable diseases 2010. World Health [Internet]. 2010;176. Available from: http://whqlibdoc.who.int/publications/2011/9789240686458_eng.pdf
4. Nasibullin TR, Timasheva YR, Sadikova RI, Tuktarova IA, Erdman V V., Nikolaeva IE, et al. Genotype/allelic combinations as potential predictors of myocardial infarction. *Mol Biol Rep*. 2016;43(1):11–6.
5. Fiatal S, Ádány R. Application of Single-Nucleotide Polymorphism-Related Risk Estimates in Identification of Increased Genetic Susceptibility to Cardiovascular Diseases: A Literature Review. *Front Public Heal* [Internet]. 2018;5(January). Available from: <http://journal.frontiersin.org/article/10.3389/fpubh.2017.00358/full>
6. Hsu Y-H, Niu T, Song Y, Tinker L, Kuller LH, Liu S. Genetic variants in the UCP2-UCP3 gene cluster and risk of diabetes in the Women's Health Initiative Observational Study. *Diabetes*. 2008;57(4):1101–7.
7. Pascual-Gamarra JM, Salazar-Tortosa D, Martinez-Tellez B, Labayen I, Rupérez AI, Censi L, et al. Association between UCP1 , UCP2 , and UCP3 gene polymorphisms with markers of adiposity in European adolescents: The HELENA study. *Pediatr Obes* [Internet]. 2019 Jan 18;(December 2018):e12504. Available from: <http://doi.wiley.com/10.1111/ijpo.12504>

8. Brondani LA, Assmann TS, de Souza BM, Bouças AP, Canani LH, Crispim D. Meta-analysis reveals the association of common variants in the uncoupling protein (UCP) 1-3 genes with body mass index variability. PLoS One [Internet]. 2014;9(5):e96411. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24804925>
9. Kotani K, Fujiwara S, Tsuzaki K, Sano Y, Nagai N, Yamada T, et al. The Association Between the Uncoupling Protein-1 Gene A-3826G Polymorphism and High-density Lipoprotein Cholesterol in A General Japanese Population: A Consideration of the Obesity Status. J Clin Med Res [Internet]. 2011;3(6):319–24. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3279477&tool=pmcentrez&rendertype=abstract>
10. Kotani K, Sakane N, Saiga K, Tsuzaki K, Shimohiro H, Tabata M, et al. The uncoupling protein-1 gene -3826A/G polymorphism and hypertension in Japanese subjects. Clin Chem Lab Med. 2007;45(9):1186–9.
11. Golozoubova V, Cannon B, Nedergaard J. UCP1 is essential for adaptive adrenergic nonshivering thermogenesis. Am J Physiol Endocrinol Metab. 2006;291(2):E350–7.
12. Moukdar F, Robidoux J, Lyght O, Pi J, Daniel KW, Collins S. Reduced antioxidant capacity and diet-induced atherosclerosis in uncoupling protein-2-deficient mice. J Lipid Res [Internet]. 2009;50(1):59–70. Available from: <http://www.jlr.org/lookup/doi/10.1194/jlr.M800273-JLR200>
13. Arsenijevic D, Onuma H, Pecqueur C, Raimbault S, Manning BS, Miroux B, et al. Disruption of the uncoupling protein-2 gene in mice reveals a role in immunity and reactive oxygen species production. Nat Genet [Internet].

- 2000;26(4):435–9. Available from:
<http://www.nature.com/doi/10.1038/82565%5Cnpapers3://publication/doi/10.1038/82565>
14. Zhang CY, Baffy G, Perret P, Krauss S, Peroni O, Grujic D, et al. Uncoupling protein-2 negatively regulates insulin secretion and is a major link between obesity, beta cell dysfunction, and type 2 diabetes. *Cell* [Internet]. 2001;105(6):745–55. Available from:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11440717
 15. Pecqueur C, Alves-Guerra C, Ricquier D, Bouillaud F. UCP2, a metabolic sensor coupling glucose oxidation to mitochondrial metabolism? *IUBMB Life*. 2009;61(7):762–7.
 16. Vidal-Puig AJ, Grujic D, Zhang C-Y, Hagen T, Boss O, Ido Y, et al. Energy Metabolism in Uncoupling Protein 3 Gene Knockout Mice. *J Biol Chem* [Internet]. 2000;275(21):16258–66. Available from:
<http://www.jbc.org/lookup/doi/10.1074/jbc.M910179199>
 17. Bezaire V, Spriet LL, Campbell S, Sabet N, Gerrits M, Bonen A, et al. Constitutive UCP3 overexpression at physiological levels increases mouse skeletal muscle capacity for fatty acid transport and oxidation. *FASEB J* [Internet]. 2005;19(8):977–9. Available from:
<http://eutils.ncbi.nlm.nih.gov/entrez/eutils/efetch.fcgi?dbfrom=pubmed&id=15814607&retmode=ref&cmd=prlinks>
 18. Riley CL, Dao C, Kenaston MA, Muto L, Nowinski SM, Solmonson AD, et al. The Complementary and Divergent Roles of Uncoupling Proteins 1 and 3 in Thermoregulation. *J Physiol*. 2016;594(24):7455–7464.

19. Su M, Chen X, Chen Y, Wang C, Li S, Ying X, et al. UCP2 and UCP3 variants and gene-environment interaction associated with prediabetes and T2DM in a rural population: A case control study in China. *BMC Med Genet*. 2018;19(1):1–9.
20. Andersen G, Dalgaard LT, Justesen JM, Anthonsen S, Nielsen T, Thørmø L, et al. The frequent UCP2 –866G>A polymorphism protects against insulin resistance and is associated with obesity: a study of obesity and related metabolic traits among 17 636 Danes. *Int J Obes* [Internet]. 2013 Feb 21;37(2):175–81. Available from: <http://www.nature.com/doi/10.1038/ijo.2012.22>
21. Mottagui-Tabar S, Hoffstedt J, Brookes AJ, Jiao H, Arner P, Dahlman I. Association of ADRB1 and UCP3 gene polymorphisms with insulin sensitivity but not obesity. *Horm Res*. 2007;69(1):31–6.
22. Hsu YH, Niu T, Song Y, Tinker L, Kuller LH, Liu S. Genetic variants in the UCP2-UCP3 gene cluster and risk of diabetes in the Women’s Health Initiative Observational Study. *Diabetes* [Internet]. 2008;57(4):1101–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18223008>
23. Ochoa MC, Santos JL, Azcona C, Moreno-Aliaga MJ, Martínez-González MA, Martínez JA, et al. Association between obesity and insulin resistance with UCP2-UCP3 gene variants in Spanish children and adolescents. *Mol Genet Metab*. 2007;92(4):351–8.
24. Moreno LA, De Henauw S, González-Gross M, Kersting M, Molnár D, Gottrand F, et al. Design and implementation of the healthy lifestyle in europe by nutrition in adolescence cross-sectional study. *Int J Obes*. 2008;32:S4–11.
25. Moreno LA, González-Gross M, Kersting M, Molnár D, de Henauw S, Beghin

- L, et al. Assessing, understanding and modifying nutritional status, eating habits and physical activity in European adolescents: the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) Study. *Public Health Nutr* [Internet]. 2008 Mar;11(3):288–99. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17617932>
26. Béghin L, Castera M, Manios Y, Gilbert CC, Kersting M, De Henauw S, et al. Quality assurance of ethical issues and regulatory aspects relating to good clinical practices in the HELENA Cross-Sectional Study. *Int J Obes (Lond)*. 2008;32 Suppl 5:S12–8.
 27. Gonzalez-Gross M, Breidenassel C, Gomez-Martinez S, Ferrari M, Beghin L, Spinneker A, et al. Sampling and processing of fresh blood samples within a European multicenter nutritional study: evaluation of biomarker stability during transport and storage. *Int J Obes (Lond)*. 2008 Nov;32 Suppl 5:S66-75.
 28. Lobelo F, Pate RR, Dowda M, Liese AD, Ruiz JR. Validity of cardiorespiratory fitness criterion-referenced standards for adolescents. *Med Sci Sports Exerc*. 2009;41(6):1222–9.
 29. Krishnan M, Thompson JMD, Mitchell EA, Murphy R, McCowan LME, Shelling AN, et al. Analysis of association of gene variants with obesity traits in New Zealand European children at 6 years of age. *Mol BioSyst* [Internet]. 2017;13(8):1524–33. Available from: <http://xlink.rsc.org/?DOI=C7MB00104E>
 30. González JR, Armengol L, Solé X, Guinó E, Mercader JM, Estivill X, et al. SNPAssoc: An R package to perform whole genome association studies. *Bioinformatics*. 2007;23(5):644–5.
 31. Sham PC, Purcell SM. Statistical power and significance testing in large-scale genetic studies. *Nat Rev Genet*. 2014;15(5):335–46.

32. Qu H-QH-Q, Tien M, Polychronakos C. Statistical significance in genetic association studies. *Clin Investig Med*. 2010;33(5):E266–70.
33. Sinnwell AJ, Maintainer Jason P Sinnwell SD. Statistical Analysis of Haplotypes with Traits and Covariates when Linkage Phase is Ambiguous. 2016; Available from: <http://www.mayo.edu/research/labs/statistical-genetics-genetic-epidemiology/>
34. Wang N, Akey JM, Zhang K, Chakraborty R, Jin L. Distribution of Recombination Crossovers and the Origin of Haplotype Blocks: The Interplay of Population History, Recombination, and Mutation. *Am J Hum Genet* [Internet]. 2002;71(5):1227–34. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0002929707604182>
35. Salopuro T, Pulkkinen L, Lindström J, Kolehmainen M, Tolppanen A-M, Eriksson JG, et al. Variation in the UCP2 and UCP3 genes associates with abdominal obesity and serum lipids: The Finnish Diabetes Prevention Study. *BMC Med Genet* [Internet]. 2009;10(1):94. Available from: <http://bmcmmedgenet.biomedcentral.com/articles/10.1186/1471-2350-10-94>
36. Ochoa MC, Santos JL, Azcona C, Moreno-Aliaga MJ, Martínez-González MA, Martínez JA, et al. Association between obesity and insulin resistance with UCP2-UCP3 gene variants in Spanish children and adolescents. *Mol Genet Metab* [Internet]. 2007 Dec;92(4):351–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17870627>
37. Pierelli G, Stanzione R, Forte M, Migliarino S, Perelli M, Volpe M, et al. Uncoupling protein 2: A key player and a potential therapeutic target in vascular diseases. *Oxid Med Cell Longev*. 2017;2017.
38. Chisolm GM, Steinberg D. The oxidative modification hypothesis of

- atherogenesis: An overview. *Free Radic Biol Med.* 2000;28(12):1815–26.
39. Mallat Z, Nakamura T, Ohan J, Lesèche G, Tedgui A, Maclouf J, et al. The relationship of hydroxyeicosatetraenoic acids and F2-isoprostanes to plaque instability in human carotid atherosclerosis. *J Clin Invest.* 1999;103(3):421–7.
40. Dhamrait SS, Maubaret C, Pedersen-Bjergaard U, Brull DJ, Gohlke P, Payne JR, et al. Mitochondrial uncoupling proteins regulate angiotensin-converting enzyme expression: crosstalk between cellular and endocrine metabolic regulators suggested by RNA interference and genetic studies. *BioEssays.* 2016;38:S107–18.

Figure Legend

- **Figure 1.** Haploview result belonging to block 1 of *UCP1* polymorphisms, which contains (rs12502572, rs11932232, rs6822807, rs6818140 y rs2071416), according to genotyping data of this study. Boxes number referred to linkage disequilibrium (D') between SNPs, boxes with no number means 100% linkage ($D' = 1$). Colour legend: i) Bright red = high D' ; White = low D' (see Haploview documentation for further details; <http://www.broad.mit.edu/mpg/haploview>).
- **Figure 2.** Blocks 1, 2 and 3 of *UCP2* and *UCP3* polymorphisms, which contains rs2735572, rs660339, rs17132534 and rs659366 SNPs for block 1 (*UCP2*); rs7930460, rs2075577 and rs2734828 SNPs for block 2 (*UCP3*); rs1800006 and rs1800849 SNPs for block 3 (*UCP3*), according to genotyping data of this study. Boxes number referred to linkage disequilibrium (D') between SNPs, boxes with no number means 100% linkage ($D' = 1$). Colour legend: i) Bright red = high D' ; White = low D' (see Haploview documentation for further details; <http://www.broad.mit.edu/mpg/haploview>).
- **Figure 3.** Significant associations between phenotypes and UPC SNPs ($FDR < 0.1$). For each association, the phenotypes and markers implicated are shown, along with the inheritance model, P value and FDR. Values are adjusted for body mass index, center, sex, and age.
- **Figure 3 (continuation).** Significant associations between phenotypes and SNPs ($FDR < 0.1$). For each association, the phenotypes and markers implicated are shown, along with the inheritance model, P value and FDR. Values are adjusted for body mass index, center, sex, and age.

Table 1: Frequency of the major and minor alleles is shown, along with the corresponding frequency of genotypes. In addition, sample size and results of exact test to assess deviations from Hardy-Weinberg equilibrium (HWE) are presented.

	Major allele	Minor allele	Major allele freq.	Minor allele freq.	Major homozygous freq.	Heterozygous freq.	Minor homozygous freq.	Sample size	pHWE
<i>UCP1</i>									
rs2071415	T	C	0.85	0.15	0.73	0.25	0.03	1057	0.39
rs7688743	G	A	0.81	0.19	0.65	0.32	0.04	1057	0.92
rs6818140	A	G	0.81	0.19	0.65	0.32	0.03	1057	0.32
rs6822807	T	C	0.73	0.27	0.53	0.4	0.07	1056	0.59
rs11932232	A	G	0.81	0.19	0.65	0.32	0.03	1057	0.32
rs12502572	G	A	0.64	0.36	0.42	0.45	0.13	1057	0.74
rs6536991	T	C	0.73	0.27	0.53	0.4	0.07	1057	1
rs2071416	T	G	0.92	0.08	0.84	0.15	0.01	1057	0.31
<i>UCP2</i>									
rs2735572	G	A	0.94	0.06	0.88	0.12	0	1057	0.58
rs17132534	T	C	0.94	0.06	0.88	0.12	0	1057	0.79
rs660339	C	T	0.61	0.39	0.38	0.47	0.15	1034	0.65
rs659366	C	T	0.64	0.36	0.42	0.46	0.13	1057	0.84
<i>UCP3</i>									
rs7930460	A	G	0.79	0.21	0.62	0.33	0.05	1057	0.78
rs2075577	T	C	0.53	0.47	0.27	0.52	0.21	1057	0.22
rs2734828	C	T	0.75	0.25	0.56	0.37	0.06	1057	0.81
rs3781907	T	C	0.74	0.26	0.54	0.39	0.07	1057	0.75

Table 2: Mean $\pm$ SD of cardiovascular risk factors.

Phenotype	All (n=1057)	Male (n=505)	Female (n=552)
Age (years)	14.71 $\pm$ 1.22	14.74 $\pm$ 1.25	14.68 $\pm$ 1.2
Weight (kg)	58.72 $\pm$ 12.67	61.86 $\pm$ 14.29	55.85 $\pm$ 10.17
Height (cm)	165.46 $\pm$ 9.34	169.5 $\pm$ 9.91	161.76 $\pm$ 6.98
BMI (kg / m ²)	21.34 $\pm$ 3.67	21.39 $\pm$ 3.99	21.3 $\pm$ 3.37
Cholesterol (mg/dL)	160.74 $\pm$ 27.69	154.03 $\pm$ 26.13	166.88 $\pm$ 27.68

HDL (mg/dL)	55.26 ± 10.67	53.17 ± 10.12	57.17 ± 10.81
LDL (mg/dL)	94.49 ± 25.09	90.78 ± 24.32	97.89 ± 25.33
Triglycerides (mg/dL)	69 ± 35.09	64.13 ± 31.65	73.46 ± 37.45
LDL/HDL	1.78 ± 0.63	1.78 ± 0.65	1.78 ± 0.6
Cholesterol/HDL	2.99 ± 0.66	2.98 ± 0.69	2.99 ± 0.63
Triglycerides/HDL	1.33 ± 0.88	1.29 ± 0.83	1.37 ± 0.92
ApoA1(mg/dL)	1.5 ± 0.22	1.46 ± 0.21	1.55 ± 0.23
ApoB (mg/dL)	0.65 ± 0.16	0.63 ± 0.15	0.68 ± 0.16
ApoB/ApoA1	0.44 ± 0.13	0.44 ± 0.13	0.45 ± 0.13
apoB/LDL	0.27 ± 0.03	0.27 ± 0.03	0.27 ± 0.03
Leptin (ng/ml)	19.61 ± 22.19	9.55 ± 14.21	28.36 ± 24.11
insulin (micro IU/mL)	10.31 ± 7.79	10.16 ± 8.82	10.46 ± 6.7
HOMA	2.35 ± 1.96	2.36 ± 2.24	2.34 ± 1.65
QUICKI	0.35 ± 0.03	0.35 ± 0.03	0.35 ± 0.03
SBP (mm Hg)	120.03 ± 13.3	124.16 ± 13.93	116.29 ± 11.5
DBP (mm Hg)	68.03 ± 8.84	67.52 ± 8.91	68.49 ± 8.76
CVD Risk score	-0.01 ± 0.61	-0.03 ± 0.66	0.02 ± 0.56

BMI: body mass index (calculated as weight in kilograms divided by height in meters squared) HDL: high density lipoprotein; LDL: low density lipoprotein; Apo: apolipoprotein; HOMA: homeostatic model assessment; QUICKI: quantitative insulin sensitivity check index; SBP: systolic blood pressure; DBP: diastolic blood pressure

•