Highlights

Rangeland Ecology & Management, January 2011

Rotational Grazing Systems and Livestock Grazing Behavior in Shrub-Dominated Semi-Arid and Arid Rangelands

Derek W. Bailey and Joel R. Brown

Rotational grazing systems are often implemented to alleviate undesirable selective grazing by livestock, but research generally shows no advantage in forage production and animal performance for these systems. Literature discussing grazing systems and grazing behavior was evaluated. Higher stock densities combined with higher stocking rates can increase livestock use of less-preferred areas, but overall distribution patterns of intensive-rotational and extensive grazing systems are often comparable at similar stocking rates and distances from water. Adjustments to animal numbers and practices that improve grazing distribution are more likely to be effective for ensuring rangeland health than fencing and rotational grazing systems.

Threshold Distinctions Between Equilibrium and Nonequilibrium Pastoral Systems Along a Continuous Climatic Gradient

Tomoo Okayasu, Toshiya Okuro, Undarmaa Jamsran, and Kazuhiko Takeuchi

On rangelands with relatively stable rainfall and plant production, carrying capacities can be calculated to determine sustainable animal numbers. In contrast, in areas with widely fluctuating plant biomass availability herders must adapt by moving to regions with better growth conditions. The relationship between plant biomass and rainfall in Mongolian grasslands has a threshold boundary that corresponds closely with the prevailing grazing management regime. It is important to identify and monitor the threshold boundary between land use systems so that managers can encourage appropriate grazing regimes in response to climatic change. Our study provides a simple, low-cost tool to evaluate ecosystems in this context.

Grazing Deferment Effects on Forage Diet Quality and Ewe Performance Following Summer Rangeland Fire

Richard C. Waterman and Lance T. Vermeire

Complete rest or grazing deferment is a general recommendation to encourage vegetative recovery following fire in the western United States. Objectives of this research were to determine grazing deferment effects on ewe diet quality and body weight change following summer fire on semiarid rangeland. Ewe average daily gain did not differ between spring and early summer grazing periods and gains during both of these periods were greater than the negligible body weight gains of the late summer grazing period. Results indicate that deferment until early summer may be preferable so that stocking rates can be more accurately determined while still optimizing animal performance.

Growing-Season Disturbance in Tallgrass Prairie: Evaluating Fire and Grazing on *Schizachyrium scoparium*

Ryan F. Limb, Samuel D. Fuhlendorf, David M. Engle, and Jay D. Kerby

Historically, fire occurred throughout the year in grassland ecosystems, but current fire prescriptions are generally limited to the dormant season because of concerns for potential damage to fire-sensitive herbaceous plants. We investigated both dormant and growing-season fire combined with grazing on little bluestem abundance and the influence of growing-season fire and clipping on survival and production of individual little bluestem plants in a controlled greenhouse experiment. Plant survival was decreased by fire and grazing in the seedling stage, but these effects were absent when plants matured. Growing-season fire did increase both herbaceous and root production and should be considered part of an overall land management strategy.

Plant Community and Soil Environment Response to Summer Fire in the Northern Great Plains

Lance T. Vermeire, Jessica L. Crowder, and David B. Wester

Summer is the natural fire season of most wildfires in the western United States. We evaluated summer fire effects on soil temperature and moisture, aboveground and root biomass, and functional group composition on Northern Great Plains rangelands dominated by semiarid cool season (C_3) species. Summer fire reduced litter, made soils 1% drier, and shifted species composition in favor of native cool season perennial grasses, but had no effect on above- or belowground productivity. Biomass was more responsive to precipitation than to fire. The fire-induced changes in species composition suggest that the exclusion of fire may be a greater disturbance than the occurrence of summer fire.

Biomass Not Linked to Perennial Grass Mortality Following Severe Wildfire in the Southern High Plains

Sandra Rideout-Hanzak, David B. Wester, Carlton M. Britton, and Heather Whitlaw

The East Amarillo Complex wildfires burned over 360,000 ha in the northern panhandle of Texas during March 2006. We investigated the effects of these wildfires on perennial grass mortality and herbaceous production. Perennial grass mortality was higher in burned areas than nonburned areas and this persisted over the course of the 3-year study; however, production was minimally affected by the wildfires. These responses illustrate the resilience of short- and mixed-grass ecosystems to severe wildfire.

Estimating Grass Yield on Blue Grama Range From Seasonal Rainfall and Soil Moisture Measurements

L. Allen Torell, Kirk C. McDaniel, and Victor Koren

Soil water availability effectively predicts herbaceous rangeland production because it considers the accumulated, recent, and past history of rainfall events. Predictive models incorporating either probe-recorded or simulated soil moisture measurements significantly improved the predictive power of regression equations relating environmental conditions to end-of-season herbaceous production. Potential exists to better predict forage conditions based on forecast information using soil water data instead of the traditional input of seasonal rainfall totals.

Long-Term Effects of Weed Control With Picloram Along a Gradient of Spotted Knapweed Invasion

Yvette K. Ortega and Dean E. Pearson

Broadleaf herbicides are commonly used in rangelands to suppress exotic weeds, but few studies have evaluated

whether these treatments succeed in restoring native communities. We conducted a 6-year experiment in western Montana to measure effects of a common herbicide treatment on grassland plants. The treatment effectively suppressed the target invader and allowed some recovery of native grasses while also depressing native wildflowers and promoting increases in a secondary invader. Our results also show that treatment outcomes can depend on initial levels of weed invasion, suggesting that selective vs. broadcast applications can be used to maximize benefits and reduce negative outcomes.

Plant Age and Growing Season Nutritional Content Relationships of Three *Artemisia tridentata* Subspecies

Barry L. Perryman, Teshome Shenkoru, Leroy B. Bruce, and Hussein S. Hussein

The effect of plant age on chemical compositions and rumen fermentation characteristics were determined for basin, mountain, and Wyoming subspecies of big sagebrush. In vitro dry matter and organic matter disappearance and ammonia nitrogen and volatile fatty acid content varied with both sagebrush age and subspecies. Age effects had minimal effect on nutritional content; therefore, we question the contention that older sagebrush plants offer less nutritional value than younger plants, at least during the growing season. Differences among sagebrush subspecies are presented and can be used to design supplementation strategies for domestic animals on diets high in big sagebrush.

Performance of Sheep and Goats Fed *Arctostaphylos canescens* With and Without Polyethylene Glycol Supplementation

Nelmy Narvaez, Arie Brosh, Miguel Mellado, and Wolfgang Pittroff

Small ruminants may provide ecologically and economically viable control of manzanita in California's chaparral and coniferous forest plantations. However, manzanita's low nutritional quality and high condensed tannin content limit its potential nutritive value. We determined the effects of polyethylene glycol supplementation on intake and digestibility of manzanita in goats and sheep. The tannins present in manzanita affect the utilization of the supplementary protein provided to goats and sheep. Adding polyethylene glycol to the drinking water at 0.15% body weight for sheep and goats is an effective way of neutralizing the deleterious effects of tannins in this shrub.

LiDAR-Based Classification of Sagebrush Community Types

Temuulen Tsagaan Sankey and Pamela Bond

LiDAR active remote sensing collects detailed, three-dimensional vegetation structure information. This study

aimed to determine if LiDAR-derived vegetation heights could be used to classify different community types in a sagebrush steppe including herbaceous, low sagebrush, big sagebrush, bitterbrush, and tree-dominated communities. Field-measured vegetation heights were strongly correlated with LiDAR-derived vegetation heights. LiDAR-derived vegetation heights varied significantly among the five community types suggesting that LiDAR data might be used to separately classify the communities. The resultant vegetation classifications could be used for land cover classification, wildfire fuels and wildlife habitat modeling, and change detection.

Field Test of Digital Photography Biomass Estimation Technique in Tallgrass Prairie

Sherry A. Leis and Lloyd W. Morrison

The digital photographic technique presented by Limb and associates in 2007 for estimating biomass was tested in two national parks with differing management history. The technique was effective at one park, but not at the other. Our results indicate the technique may not be useful for grasslands with relatively high biomass or digital image

obstruction greater than 50%; large amounts of litter may also potentially reduce the accuracy of the technique. Use of the technique requires that decision rules be developed in both the field and laboratory to reduce potential sources of error and obtain representative results.

Long-Term Vegetation Trends on Grazed and Ungrazed Chihuahuan Desert Rangelands

Francisco Molinar, Joe Navarro, Jerry Holechek, Dee Galt, and Milt Thomas

The long-term effects of managed grazing vs. grazing exclusion on vegetation composition of desert rangelands are not well known. We evaluated vegetation composition and ecological condition over a 38-year period on grazed and livestock-excluded plots dominated by black grama in the Chihuahuan Desert of southwestern New Mexico. Vegetation composition and rangeland ecological scores were similar in both treatments at the beginning and end of the study. Managed livestock grazing and exclusion had the same long-term effects on plant frequency and ecological condition. These data indicated that managed livestock grazing is sustainable on Chihuahuan Desert rangelands.