By Jeff Mosley

Browsing the Literature

This section reviews new publications available about the art and science of rangeland management. Personal copies of these publications can be obtained by contacting the respective publishers or senior authors (addresses shown in parentheses). Suggestions are welcomed and encouraged for items to include in future issues of *Browsing the Literature*. Contact Jeff Mosley, jmosley@montana.edu.

Animal Ecology

Contrasting effects of wolves and human hunters on elk behavioral responses to predation risk. K. M. Proffitt, J. L. Grigg, K. L. Hamlin, and R. A. Garrott. 2009. *Journal of Wildlife Management* 73:345–356. (Dept of Ecology, Montana State Univ, Bozeman, MT 59717, USA). Elk behavioral responses were similar to predation risk from wolves or humans (e.g., changed grouping patterns, increased movement rates), but elk responses to human-predation risk were stronger than responses to wolf-predation risk.

Habitat selection by nesting and brood-rearing sharp-tailed grouse. A. D. Goddard, R. D. Dawson, and M. P. Gillingham. 2009. *Canadian Journal of Zoology* 87:326–336. (Ministry of Environment, 400-10003-110th Ave, Fort St. John, BC V1J 6M7, Canada). In northeastern British Columbia, nesting females selected shrub-steppe habitats, greater shrub and grass cover, taller vegetation, and greater residual vegetation compared with random sites. Brood-rearing females selected agricultural habitats during the late brood-rearing period.

Landscape nutritional patterns and cattle distribution in rangeland pastures. D. C. Ganskopp and D. W. Bohnert. 2009. *Applied Animal Behaviour Science* 116:110–119. (USDA-ARS, Eastern Oregon Agricultural Research Center, 67826-A Hwy 205, Burns, OR 97720, USA). Cattle in eastern Oregon preferred to graze where forage nutritive quality exceeded threshold levels, but cattle did not prefer rare sites with exceptionally high forage nutritive quality.

Grazing Management

Consequences of prescribed fire and grazing on grassland ant communities. E. C. Underwood and C. E. Christian. 2009. *Environmental Entomology* 38:325–332. (Dept of Environmental Science and Policy, Univ of California, Davis, CA 95616, USA). Livestock grazing had no effect on ant abundance in grasslands of northern California. Prescribed fire increased some species of ants immediately post-burn, whereas other species of ants were in greater abundance 1 year after fire.

Hydrology/Riparian

An ecological risk assessment of the acute and chronic toxicity of the herbicide picloram to the threatened bull trout (*Salvelinus confluentus*) and the rainbow trout (*Oncorhynchus mykiss*). J. F. Fairchild, K. P. Weltz, L. C. Sappington, A. L. Allert, K. J.

Nelson, and J. Valle. 2009. *Archives of Environmental Contamination and Toxicology* 56:761–769. (US Geological Survey, 4200 New Haven Rd, Columbia, MO 65201, USA). “. . . Picloram, if properly applied according to the manufacturer’s label, poses little risk to the threatened bull trout or rainbow trout in northwestern rangeland environments on either an acute or a chronic basis.”

An ecological risk assessment of the exposure and effects of 2,4-D acid to rainbow trout (*Onchorhynchus mykiss*). J. F. Fairchild, K. P. Feltz, A. L. Allert, L. C. Sappington, K. J. Nelson, and J. A. Valle. 2009. *Archives of Environmental Contamination and Toxicology* 56:754–760. (US Geological Survey, 4200 New Haven Rd, Columbia, MO 65201, USA). “. . . Use of 2,4-D acid for invasive weed control in aquatic and terrestrial habitats poses no substantial risk to growth or survival of rainbow trout or other salmonids, including the threatened bull trout (*Salvelinus confluentus*).”

Competitive abilities of *Tamarix aphylla* in southern Nevada. W. E. Hayes, L. R. Walker, and E. A. Powell. 2009. *Plant Ecology* 202:159–167. (School of Life Sciences, Univ of Nevada, Las Vegas, NV 89154, USA). Athel tamarisk outcompeted the native Goodding’s willow, but Athel tamarisk was slightly less competitive than saltcedar (*T. ramosissima*).

Differential responses of postmetamorphic amphibians to cattle grazing in wetlands. E. C. Burton, M. J. Gray, A. C. Schmutzer, and D. L. Miller. 2009. *Journal of Wildlife Management* 73:269–277. (Dept of Forestry, Wildlife, and Fisheries, Univ of Tennessee, Knoxville, TN 37996, USA). In Tennessee wetlands, excluding cattle benefitted green frogs, but American toads benefitted from cattle grazing.

Impact of water troughs on cattle use of riparian zones in the Georgia Piedmont in the United States. D. H. Franklin, M. L. Cabrera, H. L. Byers, M. K. Matthews, J. G. Andrae, D. E. Radcliffe, M. A. McCann, H. A. Kuykendall, C. S. Hoveland, and V. H. Calvert. 2009. *Journal of Animal Science* 87:2151–2159. (USDA-ARS, Campbell Natural Resource Conservation Center, 1420 Experiment Station Rd, Watkinsville, GA 30677, USA). Water troughs placed away from unfenced streams reduced the amount of time that cattle spent in a riparian zone, provided the ambient temperature and relative humidity were not too high.

Seasonal changes in depth of water uptake for encroaching trees *Juniperus virginiana* and *Pinus ponderosa* and two dominant C-4 grasses in a semiarid grassland. K. D. Eggemeyer, T. Awada, F. E. Harvey, D. A. Wedin, X. Zhou, and C. W. Zanner. 2009. *Tree Physiology* 29: 157–169. (T. Awada, School of Natural Resources, Univ

of Nebraska, Lincoln, NE 68583, USA). In the Nebraska sandhills, switchgrass and little bluestem extracted most of their water from the upper soil profile. Eastern redcedar and ponderosa pine trees also extracted most of their water from shallow soil water when it was available, but the trees used deeper soil water when conditions became drier as the summer progressed.

Sources of water used by natural mesquite vegetation in a semi-arid region of northern Mexico. J. P. Brunel. 2009. *Hydrological Sciences Journal* 54:375–381. (IRD, UMR LISAH, F-34060 Montpellier 01, France). “. . . Mesquite shrubs that have access to both groundwater and surface water use surface water when available, potentially competing with grass for surface water and outcompeting grass during successive periods of drought when only groundwater is available.”

Plant Ecology

Climate change and plant invasions: restoration opportunities ahead? B. A. Bradley, M. Oppenheimer, and D. S. Wilcove. 2009. *Global Change Biology* 15:1511–1521. (Woodrow Wilson School of Public and International Affairs, Princeton Univ, Princeton, NJ 08544, USA). Authors predict that climate change in the western United States will contract the range of leafy spurge, expand the range of yellow starthistle and salt cedar, and shift the range of cheatgrass and spotted knapweed, causing both expansion and contraction.

Plant population and community responses to removal of dominant species in the shortgrass steppe. S. M. Munson and W. K. Lauenroth. 2009. *Journal of Vegetation Science* 20:224–232. (Dept of Forest, Rangeland, and Watershed Stewardship, Colorado State Univ, Fort Collins, CO 80523, USA). Removal of blue grama, the dominant plant species, did not increase plant cover of subdominant plants, nor did it increase plant diversity.

Quantifying successional rates in western aspen woodlands: current conditions, future predictions. E. K. Strand, L. A. Vierling, S. C. Bunting, and P. E. Gessler. 2009. *Forest Ecology and Management* 257:1705–1715. (Dept of Rangeland Ecology and Management, Univ of Idaho, Moscow, ID 83844, USA). Fire return intervals of 50–70 years are needed to sustain aspen stands on uplands where conifers can encroach. Current fire regimes in southwestern Idaho will cause upland aspen stands to be lost within 80–200 years.

Hawkweed identification. P. Rice and J. Halpop. 2009. Montana State University Extension Bulletin EB187. (Montana State Univ Extension Communications, 115 Culbertson Hall, Montana State Univ, Bozeman, MT 59717, USA; 12 p.) Helps readers identify seven different

species (three noxious and four native) of hawkweeds in Montana. Includes color photos of full-grown plants, leaf, root, and rosette characteristics.

Rehabilitation/Restoration

Carbon addition interacts with water availability to reduce invasive forb establishment in a semi-arid grassland. D. M. Blumenthal. 2009. *Biological Invasions* 11:1281–1290. (USDA-ARS, 1701 Center Ave, Fort Collins, CO 80526, USA). Adding carbon to the soil helped soil microbes reduce the amount of plant available nitrogen which, in turn, suppressed the noxious weeds diffuse knapweed, dalmation toadflax, and baby's breath.

Dynamic responses of a British Columbian forest-grassland interface to prescribed burning. K. Ducherer, Y. G. Bai, D. Thompson, and K. Broersma. 2009. *Western North American Naturalist* 69:75–87. (Y. Bai, Dept of Plant Science, Univ of Saskatchewan, Saskatoon, SK S7N 5A8, Canada). Prescribed fire on grassland effectively removed encroaching conifers, reduced shrubs, increased forbs, and increased aboveground total biomass.

Effects of community-level grassland management on the non-target rare annual *Agalinis auriculata*. P. Vitt, K. Havens, B. E. Kendall, and T. M. Knight. 2009. *Biological Conservation* 142:798–805. (Dept of Plant Science and Conservation, Chicago Botanical Garden, 1000 Lake Cook Rd, Glencoe, IL 60022, USA). Reducing the grazing intensity of white-tailed deer and brush removal are both necessary for the rare annual plant earleaf false foxglove to persist in tallgrass prairie.

Non-target effects of broadleaf herbicide on a native perennial forb: a demographic framework for assessing and minimizing impacts. E. E. Crone, M. Marler, and D. E. Pearson. 2009. *Journal of Applied Ecology* 46:673–682. (Wildlife Biology Program, Univ of Montana, Missoula, MT 59812, USA). Picloram herbicide application to suppress spotted knapweed benefits the native forb arrowleaf balsamroot, provided that picloram is not applied more frequently than once every 5 years.

Pyric herbivory: rewilding landscapes through the recoupling of fire and grazing. S. D. Fuhlendorf, D. M. Engle, J. Kerby, and R. Hamilton. 2009. *Conservation Biology* 23:588–598. (Dept of Natural Resource Ecology and Management, Oklahoma State Univ, Stillwater, OK 74078,

USA). Interactions between fire and ungulate grazing promote heterogeneity and biological diversity on the landscape, thereby enabling grassland ecosystems to function properly.

Soil and runoff response to dairy manure application on New Mexico rangeland. V. E. Cabrera, L. J. Stavast, T. T. Baker, M. K. Wood, D. S. Cram, R. P. Flynn, and A. L. Ulery. 2009. *Agriculture Ecosystems and Environment* 131:255–262. (Dept of Dairy Science, Univ of Wisconsin, Madison, WI 53706, USA). Light rates of dairy manure application enhanced soil properties, whereas heavy rates degraded rangeland soils. Moderate rates were not evaluated.

Successful biological control of diffuse knapweed, *Centaurea diffusa*, in British Columbia, Canada. J. H. Myers, C. Jackson, H. Quinn, S. R. White, and J. S. Cory. 2009. *Biological Control* 50:66–72. (Dept of Zoology, Univ of British Columbia, Vancouver, BC V6T 1Z4, Canada). The weevil *Larinus minutus* effectively suppressed diffuse knapweed.

Socioeconomics

A critical analysis of ecosystem services as a tool in conservation projects: the possible perils, the promises, and the partnerships. R. L. Goldman and H. Tallis. 2009. *Annals of the New York Academy of Sciences, Year in Ecology and Conservation Biology* 1162:63–78. (Nature Conservancy, 4245 N Fairfax Dr, Suite 100, Arlington, VA 22203, USA). Seven case studies highlight opportunities for using ecosystem service payments to accomplish natural resource conservation on working landscapes.

The American West's longest large mammal migration: clarifying and securing the common interest. D. N. Cherney and S. G. Clark. 2009. *Policy Sciences* 42:95–111. (Center for Science and Technology Policy Research, Univ of Colorado, Campus Box 488, Boulder, CO 80309, USA). Concludes that locally-led, collaborative group processes are the most likely ways to resolve conflicts surrounding a 170-mile migration of pronghorns that crosses multiple landowners in western Wyoming.

Jeff Mosley is Professor of Range Science and Extension Range Management Specialist, Dept of Animal and Range Sciences, Montana State University, Bozeman, MT 59717, USA.