Highlights

Rangeland Ecology & Management, September 2010

Redistribution of Runoff Among Vegetation Patch Types: On Ecohydrological Optimality of Herbaceous Capture of Runon

Anna M. Urgeghe, David D. Breshears, Scott N. Martens, and Peter C. Beeson

Vegetation growth depends not only on the amount of rainfall, but also on the amount of water redistributed laterally by surface runoff–runon. Redistribution of runoff across bare patches probably increases water available to plants and should translate into patches with increased plant productivity. This tradeoff between source areas for generating runoff and sink areas for capturing runon represents a central tenet of dryland ecohydrology. In our simulation on a semiarid piñon–juniper woodland hill slope, total amount of runon for all herbaceous patches was greatest when the amount of bare cover was intermediate. Rangeland management could potentially be improved by describing optimal relationships among the total area and distribution of bare and herbaceous patch arrangement.

Improving Potential Geographic Distribution Models for Invasive Plants by Remote Sensing

E. Raymond Hunt, Jr., John H. Gillham, and Craig S. T. Daughtry

Potential distribution models use geographic data to predict favorable locations of invasive plants; however, it is difficult to test these models with ground data. We compared areas predicted to have leafy spurge with a hyperspectral image classified for presence of the distinctive yellow-green flower bracts. By examining each geographic variable with the classification, the predictive ability of the potential distribution model was significantly improved for leafy spurge. It is generally more cost-effective to deal with the initial stages of plant invasion detected by ground monitoring; improved potential distribution models can prioritize areas that need to be monitored.

Characterizing Western Juniper Expansion via a Fusion of Landsat 5 Thematic Mapper and Lidar Data

Temuulen Tsagaan Sankey, Nancy Glenn, Sara Ehinger, Alex Boehm, and Stuart Hardegree

Juniper encroachment into rangelands is one of the most prominent changes occurring in western North America. We fused Landsat 5 Thematic Mapper (TM)– and Light Detection and Ranging (lidar)–based juniper classifications to evaluate juniper encroachment patterns in southwestern Idaho. Lidar applications for characterizing juniper encroachment attributes at finer scales were also explored. The fusion-based juniper classification model performed well (83% overall accuracy). A comparison of the resulting juniper presence/absence map to a 1965 vegetation cover map indicated 85% juniper expansion. Landsat 5 TM and lidar applications highlight areas where intensive juniper management might be prioritized.

Assessing Transportation Infrastructure Impacts on Rangelands: Test of a Standard Rangeland Assessment Protocol

Michael C. Duniway, Jeffrey E. Herrick, David A. Pyke, and David Toledo P.

Linear disturbances associated with on- and off-road vehicle use on rangelands has increased dramatically throughout the world in recent decades. However, few standardized methods exist for assessing impacts of transportation-related land-use activities on soils and vegetation. We tested the sensitivity of a qualitative assessment protocol, Interpreting Indicators of Rangeland Health (IIRH), to impacts of roads, trails, and pipelines by surveying plots at three distances from these disturbances. Results from this study indicate that IIRH is well suited for and can be consistently applied for detecting areas adversely impacted by transportation infrastructure and that it has the potential to provide information on cumulative impacts.

Comparison of Two Sampling Methods for Quantifying Changes in Vegetation Composition Under Rangeland Development

Etienne Laliberté, David A. Norton, Jason M. Tylianakis, and David Scott

Rapid vegetation sampling methods based on visual estimation are useful for monitoring changes in rangeland vegetation composition. In this paper, we compare two sampling methods in their ability to detect changes in vegetation composition following rangeland development: 1) species percent cover estimates within subplots (a more time-consuming approach), and 2) visual ranking of relative biomass of the ten most abundant species across the whole plot and the ratio between two of them (a more rapid approach). Our results support the use of the visual ranking method as a rapid yet powerful method for monitoring changes in vegetation composition under rangeland development.

Simulation of Sandsage–Bluestem Forage Growth Under Varying Stocking Rates

S. G. K. Adiku, G. H. Dunn, L. R. Ahuja, S. Gunter, J. Bradford, L. Garcia, and A. A. Andales

The long-term effects of high stocking rates on rangeland condition have been debated for years, but with a general acceptance that high stocking rates reduce ecological condition. We observed that the decline in forage growth under high stocking rates can be attributed to increased soil compaction, and we established a relationship between forage growth and the bulk density. By incorporating this relationship into a forage growth model, observed losses in forage production can be predicted. The ability to predict soil compaction effects on forage growth will help determine appropriate stocking rates for sustainable rangeland production.

Lack of Native Vegetation Recovery Following Biological Control of Leafy Spurge

Jack L. Butler and Stefanie D. Wacker

Although considerable effort is dedicated to evaluating various strategies to suppress invasive plants, detailed information on the responses of the native vegetation to these strategies are limited. This 9-year study investigated vegetation responses following significant reductions in leafy spurge dominance using flea beetles in southeastern Montana. Although total vegetative cover increased substantially following successful suppression, no change in composition or structure of the native plant community was detected for the duration of the study. Failure of the native community to recover may be attributed to a variety of interacting scenarios, some of which may be ameliorated by treating infestations as soon as possible.

Revegetation of Medusahead-Invaded Sagebrush Steppe

Kirk W. Davies

Revegetation of sagebrush steppe invaded with medusahead is needed to increase livestock forage production and to improve wildlife habitat. This study evaluated herbicide (imazapic) application, prescribed burning, and their combinations at controlling medusahead and promoting establishment of seeded bunchgrasses. In this study, seeded bunchgrasses successfully revegetated rangeland infested with medusahead when medusahead had been controlled with prescribed fire followed by fall application of imazapic. The results of this study question the long-term effectiveness of using imazapic in revegetation efforts of medusahead-infested sagebrush steppe without initially burning the infestation. This study provides useful prescriptions for successfully revegetating sagebrush steppe invaded with medusahead.

Grazing Management and Microclimate Effects on Cattle Distribution Relative to a Cool-Season Pasture Stream

M. M. Haan, J. R. Russell, J. D. Davis, and D. G. Morrical

Pollution of pasture streams may be decreased by controlling distribution of grazing cattle. Based on visual observations and GPS collar information, cows under continuous stocking with unrestricted stream access spent more time in the stream or in the streamside zone than cows under continuous stocking with restricted stream access or under rotational stocking. However, the availability of off-stream water had no effect on the amount of time cattle spent in or near the stream. Use of rotational stocking or stabilized stream access and crossing sites allows use of stream water by grazing cattle while controlling pollution.

Spring Grazing Impacts on the Vegetation of Reed Canarygrass-Invaded Wetlands

Heidi L. Hillhouse, Susan J. Tunnell, and James Stubbendieck

Grazing has been used to control reed canarygrass invasion in wetlands in the Rainwater Basin region of Nebraska, but it has been unclear whether current grazing techniques were providing the desired results. We collected data on four publicly owned wetlands to determine whether reed canarygrass abundance was decreased by grazing and to see how other aspects of the vegetation were affected. We found that grazing did not decrease reed canarygrass, but it did have beneficial impacts on other habitat characteristics. By decreasing litter buildup and increasing bare ground, graz-

ing produced conditions that can promote the germination and growth of desirable annual wetland plants.

Chaining and Burning Modifies Vegetation Structure, Fuel, and Post-Disturbance Sprouting Capacity

Carl R. Gosper, Suzanne M. Prober, and Colin J. Yates

Dislodging vegetation using a chain and then burning it after a period of curing is used to manipulate fuels in native vegetation in the global biodiversity hotspot of southwest Western Australia. We investigated the ecological outcomes of combining these two disturbances with burning alone. Chained and burnt strips had reduced standing dead vegetation and produced fewer stems in sprouting *Eucalyptus* species than did solely burnt vegetation. Balancing the fire management benefits of chaining and burning with the ecological consequences is a significant challenge facing land managers in this fire-prone landscape.

Vegetation Composition and Nutritional Quality of Forage for Gazelles in Eastern Mongolia

Kirk A. Olson, Martyn G. Murray, and Todd K. Fuller

Mongolian gazelles face numerous threats to the integrity of their rangeland habitat. Although conservation plans are being developed for these rangelands, there is little knowl-

edge of their vegetation's nutritional qualities. We studied vegetation composition, forage quality, and trace-element content to understand how forage characteristics could influence gazelle distribution across the steppe. Vegetation is generally mineral deficient; however, certain species contain high levels of important nutrients that could meet the nutritional demands of the migratory gazelles. Understanding how key forage species are distributed is important so that gazelle habitat conservation can take into consideration regions of high forage value.

Weather Variables Affecting Oklahoma Wildfires

Angela M. Reid, Samuel D. Fuhlendorf, and John R. Weir

Wildfires can be a destructive force to human life and property, and it is important that we constantly improve our ability to predict situations with a high danger of fire. We used weather and wildfire data from Oklahoma to look at the relationship between different weather variables and wildfire size, as well as describe the timing of wildfires. Minimum relative humidity, maximum and average wind speeds, and precipitation were the most important factors relating to wildfire size. Additionally, the largest wildfires generally occurred in December and January. This information is important to improve modeling capabilities as well as weather condition recommendations for prescribed fires.