

Control of Citrus Nematode with Cadusafos¹

Michael A. McClure and Mark E. Schmitt

Abstract

Granular (Rugby 10G) and liquid (Rugby 100ME) formulations of Cadusafos were evaluated for the control of Tylenchulus semipenetrans on mature lemon trees in a commercial citrus orchard at Yuma, Arizona. Three applications of cadusafos, with two months between applications, at the rate of 2 g a.i./m² reduced nematode populations to undetectable levels and increased the yield and rate of fruit maturity of 'Rosenberger' lemons. Yields were increased 12,587 kg per hectare with Rugby 100ME and 8,392 kg per hectare with Rugby 10G. Nematode populations were suppressed for at least 12 months after the last application.

Introduction

The citrus nematode, *Tylenchulus semipenetrans*, is one of the most debilitating pests of citrus worldwide (2). It occurs throughout Arizona where up to 65% of the bearing trees are infested (1). Until 1977, 1,2-dibromo-3-chloropropane (DBCP) was used to maintain nematode populations at acceptable levels. DBCP was low in phytotoxicity, highly nematicidal, and easily applied in irrigation water. Fruit size and total yield were increased for up to 3 years following a single treatment (4).

In 1977 the registration of DBCP was canceled due to environmental and human health concerns, leaving Arizona's citrus growers with few alternatives for nematode control. Currently, the only nematicides registered for use on citrus in Arizona are aldicarb and fenamiphos. Fenamiphos has not proven efficacious on desert-grown citrus that is flood irrigated (5) and aldicarb while effective, has not gained favor in Arizona due to its high mammalian toxicity, mobility in soil water, and requirements for mechanical incorporation into the soil. Discovery of aldicarb in groundwater in California and Florida has restricted its use in those states.

Cadusafos (S,S-di-sec-butyl O-ethyl phosphorodithioate), an organophosphate pesticide (FMC Corp., Philadelphia, PA) has been registered in the Republic of South Africa for citrus nematode control since 1992. The efficacy of Cadusafos has been thoroughly documented by le Roux (3) who found that the life cycle of *T. semipenetrans* could be broken by three applications of 2.0 g a.i./m² at two-month intervals during the growing season, essentially "eliminating" the nematode from the orchard." In addition to its nematicidal properties, Cadusafos has several characteristics that make it a promising nematicide for citrus nematode control. According to the manufacturer, Cadusafos is not absorbed by the plant and residues of the material are not found in fruit of treated trees. Limited water solubility and low to moderate soil mobility reduce the threat of ground water contamination. Half-life in silty clay loam and sandy soils is estimated to be about 45 days. It is formulated as an emulsifiable concentrate (10% w/v), or granule (10% w/w) under the trade names Rugby 100ME and Rugby 10G respectively.

¹Supported by a grant from the Arizona Citrus Research Council. This is the final report for project 94-16 'Citrus Nematode control with Cadusafos' The authors thank C.V. Spencer, Yuma, Arizona for providing the test sites, harvest labor, and for sharing his vast knowledge of citriculture.

Our purpose was to evaluate Cadusafos for the control of citrus nematode on desert-grown citrus. Emulsifiable liquid and granular formulations were tested for efficacy and improvement of citrus yield and quality.

Materials and Methods

Two plots were established on the property of C.V. Spencer, Inc., a commercial citrus grower in Yuma, Arizona. Both plots, naturally infested with *T. semipenetrans*, contained 27 to 30-year-old "Rosenberger" lemon trees, *Citrus limon* (L.) Raf., on rough lemon root stock at a density of 119 trees per hectare. One plot, irrigated by a low-pressure microjet system was used to evaluate the emulsifiable formulation of Cadusafos (Rugby 100ME) and the other, flood irrigated, was used for the granular formulation (Rugby 10G). The plots were located approximately 3 miles apart on the Yuma mesa. The soil of this area is a deep Superstition sand consisting of sandy, mixed, hyperthermic Typic Calciorthids.

Both plots received three applications of Cadusafos at two-month intervals on May 18, July 14, and September 13, 1995. The Rugby 100ME plot consisted of 15 treated trees and 15 untreated control trees, in 10 rows of 3 trees each. Treated rows alternated with untreated rows. Rugby 100ME was applied by pressure-injection through the microjet system at the rate of 2.0 g a.i. per m². Each tree received approximately 36 g a.i. of Rugby 100ME in 75 liters of water followed by 300 liters of water from the microjets. Thereafter, the orchard was irrigated daily with 300 liters of water per tree from June through October and, as needed, from November to May.

The Rugby 10G plot consisted of a single block of 24 trees, 12 treated and 12 controls, in a completely randomized design. Rugby 10G was distributed evenly with a hand-held fertilizer spreader, from the drip line to the trunk of the tree, at the rate of 2.0 g a.i. per m². Each tree received about 36 g a.i., depending upon the area treated. The granules were then hand-raked into the top 2.5 cm of soil and the orchard was flooded with approximately 200 cm of water. Irrigation by flooding continued, every 14 days from June through October, and at irregular intervals, as needed, from November to May.

Pretreatment and post-treatment nematode population densities were estimated by counting the number of juveniles and males extracted from feeder roots. Roots were taken from the top 30 cm of soil from the north, east, south, and west sides of each tree, 30 to 60 cm inside the drip line just prior to application of Cadusafos and at 2 months (November 17, 1994) and 12 months (September 21, 1995) after the third application. Roots were shaken gently to remove loosely adhering soil and the four subsamples from individual trees were combined, cut into 2 cm pieces, and a 5 g portion was incubated on Baermann funnels under an intermittent mist of distilled water for 72 hours. Juveniles and males that migrated from the roots, or hatched from egg masses on the roots, were collected and counted.

Rate of fruit maturity was estimated by measuring fruit diameter at weekly intervals for the 9 weeks prior to harvest on October 11, 1994. Fifty lemons, selected at random from each tree, were measured and their mean diameter calculated. Five of these, each with a diameter within the standard error of the mean of the parent set, were tagged and measured thereafter at weekly intervals. Yields were not measured in 1994.

At harvest, twelve months after the third treatment with Cadusafos, all of the lemons from each tree were removed in a single picking and weighed. Fifty lemons from each tree were randomly selected and graded according to industry standards (1st grade, 2nd grade, and culls) by Associated Citrus Packers, Inc., Yuma, AZ. All numerical data were analyzed by SigmaStat Scientific Software (Jandel Corp., San Rafael, CA).

Results and Discussion

Populations of *T. semipenetrans* were reduced ($P < 0.001$) with both Rugby 100ME and Rugby 10G (Fig. 1). Two months after the third application of these materials, nematodes were detected on roots of only two of the 15 trees treated with Rugby 100ME and only three of 12 trees treated with Rugby 10G. Twelve months after the third application, nematodes were detected on roots of six trees treated with Rugby 100 ME and on five trees treated with Rugby 10G. On the treated trees with detectable nematode populations, the numbers of nematodes ranged

from 10 to 22 percent of the controls. Natural seasonal fluctuations in nematode populations, unrelated to the treatments, were evident in both plots.

Rugby 100ME increased ($P=0.007$) fruit size during the nine weeks prior to harvest (Fig. 2). Fruit size was not affected by treatment with Rugby 10G (data not shown). Based on 118 trees per hectare, yield was increased ($P=0.005$) by 12,587 kg per hectare following three applications of Rugby 100ME and by 8,392 kg per hectare ($P=0.019$) with Rugby 10G, a 45% and 29% increase, respectively (Fig 3). Fruit quality, based on the percentage of 1st and 2nd grades was not affected by either treatment (data not shown).

Cadusafos is one of the most promising nematicides for control of citrus nematode since DBCP was withdrawn from the market in 1977. Not only were nematode populations controlled but, on many trees, they were reduced to a level below the limit of detection. Nematode suppression was achieved for at least 12 months, suggesting that repeated applications would be necessary only in alternate years or longer intervals. le Roux (3) estimated that, in South Africa, it would take at least four years for populations to reach levels requiring additional treatment.

Because the test plots used in this trial were geographically separated and employed different irrigation systems, it was not possible to compare the efficacy of Rugby 100ME and Rugby 10G. Both formulations reduced nematode populations, but yield responses and fruit maturity rates were greater in the plot treated with Rugby 100ME. This may have been due, in part, to the higher initial populations of nematodes in the Rugby 10G plot. According to le Roux (3), nutritional imbalances, that occur in citrus trees infected for many years with nematodes, may not be rectified immediately when the nematodes are eliminated. Initial populations of citrus nematodes in the Rugby 10G plot were four times those in the Rugby 100ME plot and so it is reasonable to infer that damage to these trees was correspondingly greater and that the trees would take longer to recover.

Relative amounts of 1st and 2nd grade lemons harvested from the test plots did not change as a result of Cadusafos treatment. However, because total yield was substantially increased, the absolute yield of higher grades increased proportionally. Thus, greater economic returns could be expected from treatment with Cadusafos than would be realized from yield increases alone.

Currently, Cadusafos is not registered for use on any crop in the United States. Should registration be achieved, our trials show that Cadusafos could be an alternative to DBCP or currently registered nematicides. Cadusafos may also prove to be a management tool for "earliness" when date of fruit maturity is an important factor in marketing strategy.

Literature Cited

1. Baines, R. C. and R. H. Small. 1969. Efficacy of some new compounds for control of the citrus nematode, *Tylenchulus semipenetrans* on established trees. Proceedings of the 1st International Citrus Symposium, Riverside, CA, 1968. 2:973-977.
2. Duncan, L. W., and E. Cohn. 1990. Nematode parasites of citrus. Pp. 321-346 in M. Luc, R.A. Sikora, and J. Bridge, eds. Plant parasitic nematodes in subtropical and tropical agriculture. Wallingford, UK: CAB International.
3. le Roux, H. F. 1995. Control of the citrus nematode in South Africa. Ph.D. dissertation, Department of Microbiology and Plant Pathology, University of Pretoria, South Africa.
4. Reynolds, H. W., O'Bannon, J. H. and E. L. Nigh. 1974. The citrus nematode and its control in the southwest. Technical Bulletin No. 1478. Agricultural Research Service, United States Department of Agriculture, Hyattsville, MD.
5. Van Gundy, S. D., Garabadian, S. and E. L. Nigh. 1981. Alternatives to DBCP for citrus nematode control. Proceedings of the International Society for Citriculture 1:387-390.

