

## RESEARCH REPORT

### THE DEVELOPMENT AND STATE OF DENDROCHRONOLOGY IN THE USSR

STEPAN G. SHIYATOV

Institute of Plant and Animal Ecology  
Ural Division of the USSR Academy of Sciences  
8 Marta Street, 202, Sverdlovsk, 620008, USSR

#### ABSTRACT

The first dendrochronological investigation in the USSR was carried out at the end of the past century. Systematic study of tree rings for the purpose of dating different events and reconstruction of natural conditions began in 1950-1960's. Tree-ring analysis is most intensively used in the studies of forest ecosystem dynamics, timing and frequency assessment of catastrophic phenomena, reconstruction of radiocarbon content in the Earth atmosphere, and dating of historical wood. Much attention is given to the development of long-term prognoses of tree growth and forest environments.

The first dendrochronological investigation in the USSR, which has not been recognized so far, was carried out by Professor Shvedov in 1882 on two trees of black locust (*Robinia pseudoacacia* L.) growing in the streets of Odessa (Shvedov 1892). He discovered that there is a close relation between ring widths and precipitation of the hydrological year (from September to August) and that the narrowest wood layers form in very dry years, which are regularly repeated in three and nine years. On the basis of this periodicity, he predicted a drought for 1891. This prognosis proved to be accurate and Shvedov subsequently published the results of the investigations. He demonstrated the possibility of "dendrometric method" for reconstruction of climatic conditions of the past. Shvedov should be rightfully considered as one of the founders of dendrochronology.

Unfortunately these investigations, which began so successfully, were not continued over a long period of time. Only in the 1930's dendrochronological papers began to be published. These articles were basically methodological (Zaozersky 1934; Tolsky 1936) and only the work by Kostin (1940) was devoted to the reconstruction of droughts in the European part of the Soviet Union. At the same time abstracts of the research of American dendrochronologists (A. Douglass, W. Glock) were published (Jashnov 1925; Krishtofovitsh 1934; Chrgian 1938) and these were valued very highly.

Systematic investigations in the field of dendrochronology began only from the 1950's. Rudakov (1951, 1958, 1961) popularized the dendroclimatological method. In particular, he made popular the moving average method for determination of the age trend and the calculation of growth indices. Some investigations were carried out to discover the effects of climatic factors on tree-ring growth: in the Latvian SSR (Zwiedris and Sacinieks 1960; Zwiedries and Matuzanis 1962), in the Karelian Neck (Dmitrieva 1959), in the Eastern part of the European territory of the USSR (Liseev 1962; Rudakov 1958, 1961), and in the Lithuanian SSR (Bitvinskas 1965). Interesting work on qualitative reconstruction of humidity in the Middle Asia highlands during the last millenium on the basis of the ring-widths study of Turkestan juniper (*Juniperus turkestanica* Kom.) was carried out by Gursky et al. (1953). Tree rings were extensively used by Galazy (1954) for reconstruction of climatic conditions

and the upper timberline dynamics in the Zabaikalie and for estimation of the dates of high water levels in Lake Baikal (Galazy 1955, 1967). Kostin (1960, 1963) has studied ring-width variability for the purpose of reconstruction of severe droughts in the last 200-300 years in the steppe zone. Shiyatov (1965) has made a qualitative reconstruction of the thermal conditions in summer months in the Polar Urals.

At the end of the 1950's dendrochronological methods began to be used for dating historical and archaeological monuments and relics. Zamotorin (1959, 1963) carried out relative dating of the Altai-Sajan Barrows. In 1959 the first Dendrochronological Laboratory at the Institute of Archaeology of the USSR Academy of Sciences (Moscow) under the leadership of Professor Kolchin was organized (Kolchin 1963). The Laboratory was engaged in absolute dating of medieval buildings, churches, and paved wood roadways in the northwestern part of the Soviet Union, in Novgorod in particular.

From 1968 until the present, there has been a period of broad application of dendrochronology in the study of variability of various natural and anthropogenic processes and dating of various events. That year the First All-Union Conference on Dendrochronology and Dendroclimatology (in Vilnius) was held and the Dendroclimatochronological Group at the Institute of Botany of the Lithuanian Academy of Sciences (in Kaunas) was organized. In 1976 this Group was reorganized into the Dendroclimatochronological Laboratory and at present this Laboratory is the largest scientific subdivision in the Soviet Union that specializes in dendrochronology (Bitvinskas 1978).

In the Soviet Union during the last two decades, tree-ring analysis has been most intensively used for the study of forest ecosystem dynamics. First of all this is related to the fact that the majority of Soviet dendrochronologists are foresters and they work in Forestry and Ecological Institutes.

The work of Soviet dendrochronologists is based on a form of forest ecosystem dynamics called the cyclic form. Within the cold and temperate zones, annual fluctuations and long-term changes in climatic conditions significantly affect the composition and structure of forest ecosystems. Changes of forest environments caused by climatic variations (moisture and heat supply, droughts, fires, floods, snow avalanches, and other catastrophic phenomena) are the main reasons of the cyclic forest dynamics. The duration and amplitude of processes which are not strictly periodic, but change to some extent, allowing the estimate of the oscillation parameters by statistical methods are called cyclic. Not only short-term fluctuations (daily, seasonal, annual, intrasecular), but long-term and essential ones (secular and oversecular) should be attributed to cyclic forest dynamics, including forest vegetation succession into woodless vegetation and back.

The cyclic processes can be observed in the changes of almost all components of forest ecosystems. Most of all they are expressed in tree growth. There are various length cycles in the dynamics of separate components of forest ecosystems (polycyclicality). But there are usually only a few dominant cycles, which bring about the highest contribution in the variability of the process.

The presence of certain cycles are characteristic of various forest ecosystem components and processes. For example, seed crop dynamics are characterized by short-term cycles, less than 10 years. Forest successions are determined by long-term climatic cycles (secular and oversecular). The cyclic processes are expressed primarily in regions which unfavorable for tree growth (arid and cold regions and sites). The greatest contribution to the study of forest dynamics was made by the following research: Komin (1963, 1970); Bitvinskas (1964, 1974); Kolishchuk (1966, 1979); Shiyatov (1972, 1975, 1986); Gortinsky (1968); Malokvasov (1974); Olenin (1976, 1977); Polyushkin (1979); Mazepa (1986); and Dyrenkov et al. (1987).

Frequently dendrochronology is used for estimation of the effectiveness of various forest measures (melioration, fertilization and so on). The comparison of the growth of trees which are or are not subject to the effects of the studied factors are used most of all (Bitvinskas 1965, 1974; Buzikin 1978; Evdokimov 1979; Pshenichnikova 1987).

In the coniferous boreal and broad-leaf forests of the Soviet Union severe insect outbreaks are often observed. As the result of the full defoliation large massives of forests are dried out. Therefore an important task is estimation of the timing and intensity of insect outbreaks. For reconstruction of outbreaks such indicators as growth of host and non-host species and trees, the width and index of summer wood, dimension and number of cells in the annual rings, and histograms of indices for each calendar year are used (Litvinenko 1972; Vaganov et al. 1972; Isaev and Kiselev 1987; Kucherov 1987).

Several investigations regarding the influence of seed crops on the annual growth of trees have been carried out. The dark-needled trees (*Picea*, *Abies*) grow at a reduced rate of about 50 percent during the harvest year in comparison with non-seedbearing trees. Reduction of the summer wood percentage in annual rings and a sharp decrease of apical growth were observed (Danilov 1953; Kolishchuk et al. 1975; Voronin 1986).

The influence on the tree growth of such anthropogenic factors as pollution and recreation are studied intensively at present. More frequently comparison of tree growth, which is influenced or not by the examined factor is conducted (Lairand et al. 1979; Lovelius 1979; Laletin 1987; Yuknees 1987; Sabirov 1987). In studies of forest decline, climatic response models have not yet been used.

For estimating pollution effects on forest ecosystems, a determination of chemical elements, especially heavy metals in the annual wood layers, has been determined (Chetverikov 1986; Adamenko et al. 1987). It is interesting that the ratio of potassium in the annual wood layers was determined (Chetverikov 1986).

Soviet dendrochronologists also pay much attention to reconstruction of the timing and frequency of such catastrophic phenomena as snow avalanches, mud slides, land slides, fires, and windthrows (Melechov 1948; Karpenko and Medvedev 1963; Turmanina 1971, 1979; Zabelin 1979; Gorchakovskiy and Shiyatov 1985; Nesvetailo 1986; Shiyatov and Uljanov 1987). For dating these phenomena the following indicators are used: evidence of mechanical damages on trees, reaction wood, new vertical shoots, and appearance and determination of when the trees sprouted and died. The greatest contribution to this research was made by scientific workers of the Problem Laboratory of the Geographical Faculty of the Moscow State University (Turmanina 1971, 1972, 1979; Akifieva and Turmanina 1970; Lukjanova and Mjagkov 1979).

Presently intensive investigations dating past forest fires are being carried out in Siberian forests (Furyaev 1987; Valentic and Ivanova 1987; Evdokimenko and Koptsev 1987).

Although dendroclimatic reconstructions are being carried out by many researcher (Glebov and Pogodina 1972; Bitvinskas 1974; Muchamedshin 1974; Bitvinskas and Kairaitis 1975; Lovelius 1979; Borshova 1981; Shiyatov 1986; Adamenko 1986), only simple linear and non-linear regressive models are being used. There are no spatial reconstructions because the network of dendrochronological stations is not yet fully developed in many regions of the USSR.

Dating of historical and archaeological wood is being conducted mainly in the European territory of the Soviet Union (Kolchin and Chernich 1977; Kolishchuk et al. 1984; Brukstus 1986). In the eastern and southern regions, such dating has been performed rarely (Shiyatov 1980; Komin 1980). This is because of the absence of specialists and laboratories in these regions.

In the Soviet Union, research on the problem "Astrophysical Phenomena and Radiocarbon" have been carried out for the past 20 years. The purpose of these researches is the annual reconstruction of the radiocarbon content in the Earth atmosphere on the basis of estimation of its contents in annual wood layers (Dergachev and Kocharov 1981). The Dendroclimatochronological Laboratory (Kaunas), the Ioffe Physico-Technical Institute (Leningrad) and many Radiocarbon Laboratories participate in this research (Bitvinskas 1981). The annual reconstruction of radiocarbon contents for the last 200-300 years has been calculated. The close relation of radiocarbon content in the annual layers of trees growing in various regions of the country has been demonstrated. The relation between solar activity, especially in 11-, 22- and 80-year cycles and radiocarbon content in wood has been suggested (Bitvinskas 1984).

Dendrochronological methods are used for determining the time and place of criminal actions, especially regarding the illegal cutting and selling of timbers (Rosanov 1965, 1968).

Soviet dendrochronologists give much attention to the development of the long-term prognoses of tree growth and forest environments. Such prognoses are of a great importance, especially in regions of insufficient moisture and warmth. Economic effects of prognoses may lead redistribution and more effective utilization of investment, as well as due to the increase of forest productivity and forest protective functions. The polyharmonic models are most frequently used (Komin 1972; Polyushkin 1979; Shiyatov 1986; Mazepa 1986; Kairiukstis and Dubinskaite 1986). They are based on establishment, approximation, and extrapolation of the most important cyclic components in dendrochronological series. To determine the necessary parameters of cycles (length, amplitude and phase), Mazepa (1986) used data of spectral density and narrow-band filtration. The approximation of the cycles was conducted by sinusoids. Usually from 11 to 20 cycles are used in approximation and extrapolation of each chronology. The correlation coefficient between the original and approximated chronologies ranges from 0.5 to 0.8 (Shiyatov and Mazepa 1986).

For prognoses of growth indices, predicted data of solar activity (Bitvinskas 1984) and calculated data of distribution of the maxima atmospheric tides of the Moon and of the Sun (Javorsky 1975) are also used.

In many regions of the Soviet Union, seasonal growth of trees and factors which determine the duration and rate of growth are studied. The microscopic method is used most frequently in studying seasonal growth (Kairiukstis and Yuodvalkis 1970; Lobzhanidze 1975; Kishenko 1978; Goryachev 1987). Presently the most intensive research of seasonal and cell growth are carried out at the Biophysical Institute of the Siberian Division of the USSR Academy of Sciences (Krasnoyarsk). At this Institute, a special device "Ring Structure Measurer" which semiautomatically registers the number and dimension of cells in the annual rings is constructed (Vaganov et al. 1985).

Up to the present, about 370 tree-ring chronologies have been published in the form of indices. From these chronologies, only 115 are from the eastern and southern regions of the USSR (including the Urals). Most of the chronologies have been obtained from coniferous species (*Larix*, *Pinus*, *Picea*, *Abies*), while only about 60 chronologies are from broadleaf-bearing species (mainly from *Quercus*). Very few chronologies have been developed in Siberia, the Far East, Middle Asia, or the Caucasus. The longest published chronologies are the following: for *Juniperus turkestanica* Kom., Middle Asia, 1224 years (Kolchin and Chernich 1977), for *Pinus sylvestris* L., Novgorod, 1200 years (Kolchin and Chernich 1977), for *Larix sibirica* Ldb., Polar Urals, 1010 years (Shiyatov 1986), for *Larix sibirica* Ldb., North of Western Siberia, 867 years (Shiyatov 1975), and for *Larix sibirica* Ldb., Altai Mountains, 677 years (M. F. Adamenko 1978).

Dendrochronological investigations in the USSR are coordinated by the Commission for Dendroclimatology of the USSR Academy of Sciences. This Commission sponsors all the All-Union Conferences on problems of dendrochronology and dendroclimatology (1968 - Vilnius, 1972 - Kaunas, 1978 - Archangelsk, 1983 - Irkutsk). On the initiative of the Commission in 1980 the Dendrochronological Bank of the Soviet Union (DBSU) at the Dendroclimatochronological Laboratory and the Lithuanian Forest Research Institute (Kaunas) was organized. With the Commission assistance, three volumes of "Dendroclimatological Scales of the Soviet Union" (1978, 1981, 1984) and the bibliographic reference "Dendroclimatochronology, 1900-1970" (Vilnius 1978) were published.

### REFERENCES

- Adamenko, M. F.  
 1978 Growth dynamics of Larch as indicator of thermal regime of summer seasons in the Gorny Altai. In Regional Geographical Investigations in the Western Siberia. Novosibirsk: Nauka, pp. 20-25 (in Russian).  
 1986 Reconstruction of thermal regime dynamics of summer seasons during XIV-XX centuries in the Gorny Altai. In Dendrochronology and Dendroclimatology. Novosibirsk: Nauka, pp. 110-114 (in Russian).
- Adamenko, V. N., E. L. Juravljova and S. A. Sinjakov  
 1987 Elemental composition of the season increment as a characteristic of natural and anthropogeneous factors. In Dendrochronological Methods in Forest Science and Ecological Forecasting. Thesis of the International Workshop. Irkutsk, pp. 126-128 (in Russian, English).
- Akifieva, K. V. and V. I. Turmanina  
 1970 Phytoindication of avalanche fall frequency in the Elbrus region. In Materials of the IV Conference on the Snow Cover, Snow Avalanches and Glaciers Study. Leningrad: Gidrometeoizdat, pp. 74-81 (in Russian).
- Bitvinskas, T. T.  
 1964 Growth stand dynamics and possibilities of their prognosis (in the Lithuanian SSR conditions). Reports of the Timirjazev Agricultural Academy, No. 99:497-503 (in Russian).  
 1965 On the question of the use of dendroclimatological methods in forestry. Reports of the Timirjazev Agricultural Academy, No. 115:201-207 (in Russian).  
 1974 Dendroclimatological investigations. Leningrad: Gidrometeoizdat, 170 p. (in Russian).  
 1978 Scientific results of the Dendroclimatochronological Laboratory of the Institute of Botany of the Lithuanian Academy of Sciences. In Environments and Radial Tree Growth. Kaunas, pp. 3-9 (in Russian).  
 1981 Dendroclimatochronology of the Northwestern part of the European Sector of the USSR (from the investigations of the dendroclimatochronological Laboratory of the Institute of Botany of the Academy of Sciences, Lithuanian SSR). Report to the International Symposium on "The Influence of Changes in Solar Activity on Climate", May 24-30, 1981, Kaunas, 26 p. (in Russian).  
 1984 Formulation of the fundamentals of dendroclimatological researches in Lithuanian SSR. In Temporal and Spatial Changes and Annual Tree Rings. Kaunas, pp. 4-48 (in Russian).
- Bitvinskas, T. T. and I. I. Kairaitis  
 1975 Radial growth dynamics of oak stands in the Lithuanian SSR and its relations with environments, climate and solar activity. In Biocological Backgrounds of Dendrochronology. Vilnius-Leningrad, pp. 69-74 (in Russian).
- Borshova, N. M.  
 1981 Effects of climatic factors on radial tree growth of *Picea schrenkiana* in the Zailiysky Alatau. In Biocological Investigations in the Tien-Shan Spruce Forests. Frunze, pp. 157-167 (in Russian).
- Buzikin, A. I.  
 1978 On the evaluation of effectiveness of forest management upon the analysis of the annual ring-widths. Thesis on Reports on the 3rd All-Union Conference on Dendroclimatology. Archangelsk, pp. 158-159 (in Russian).
- Brukstus, V. I.  
 1986 Experience of dendrochronological scale construction on pine in the Lithuanian SSR. In Dendrochronology and Dendroclimatology. Novosibirsk: Nauka, pp. 187-191 (in Russian).
- Chetverikov, A. F.  
 1986 Chemical composition of annual wood layers and environments. In Dendrochronology and Dendroclimatology. Novosibirsk: Nauka, pp. 126-130 (in Russian).

- Chrgian, A.  
1938 Waldo Glock. Principles and methods of tree-ring analysis, Carnegie Institution, Washington, 1937. *Meteorology and Hydrology*, No. 7:165-166 (in Russian).
- Danilov, D. N.  
1953 The effect of seed crops on annual layers structure of spruce (*Picea exelsa* Link.) *Botanical Journal*, Vol. 38, No. 3:367-377 (in Russian).
- Dergachev, V. A. and G. E. Kocharov  
1981 Regularity manifestation of some natural processes in the past by radiocarbon concentration study in the tree wood layers. In *Spatial Climatic Changes and Annual Tree Rings*. Kaunas, pp. 27-39 (in Russian).
- Dmitrieva, E. V.  
1959 Analysis of climatic influences on tree growth at different sites in the Karelian Neck. *Botanical Journal*, Vol. 44, No. 2:162-176 (in Russian).
- Dyrenkov, S. A., S. N. Sawitskaya and I. G. Sokolova  
1987 Dendrochronological analysis of secondary successions. In *Dendrochronological Methods in Forest Science and Ecological Forecasting*. Thesis of the International Workshop. Irkutsk, pp. 229-233 (in Russian, English).
- Evdokimov, V. N.  
1979 Tree growth as a function of drainage intensity at the horsetail-bogmoss spruce stand in the Komi Autonomous Republic. *Proceedings of Institute of Higher Education. Forestry Journal*, No. 2:30-32 (in Russian).
- Evdokimenko, M. D. and S. G. Koptsev  
1987 The peculiarity of the annual increment of the forest in the extremal conditions (an example of the lightpine forest of the Baikal region). In *Dendrochronological Methods in Forest Science and Ecological Forecasting*. Thesis of the International Workshop. Irkutsk, pp. 289-294 (in Russian, English).
- Furyaev, V. V.  
1987 Dendrochronological indication and aerial photos and satellite imagery processing in long-term fire effects prediction. In *Dendrochronological Methods in Forest Science and Ecological Forecasting*. Thesis of the International Workshop. Irkutsk, pp. 237-242 (in Russian, English).
- Galazy, G. I.  
1954 Altitudinal limit of arboreal vegetation in the Eastern Siberia Mountains and its dynamics. *Proceedings of Botanical Institute of the USSR Academy of Sciences, Series III (Geobotany)*, No. 9:210-329 (in Russian).  
1955 Botanical method of determination of the high historical water levels and some other phenomena in the Baikal Lake. *Proceedings of the USSR Academy of Sciences*, Vol. 103, No. 5:811-813 (in Russian).  
1967 Tree growth dynamics on the Baikal shores in connection with cyclic water level oscillations in the Baikal Lake. In *Geobotanical Investigations in the Baikal Lake*, Moscow: Nauka, pp. 44-301 (in Russian).
- Glebov, F. Z. and A. I. Pogodina  
1972 Stand growth of some bog forests at the Tomsk station in connection with hydrothermal conditions. In *Dendroclimatochronology and Radiocarbon*. Kaunas, pp. 120-126 (in Russian).
- Gorchakovskiy, P. L. and S. G. Shiyatov  
1985 Phytoindication of environments and natural processes in the highlands. Moscow: Nauka, 208 p. (in Russian).
- Gortinsky, G. B.  
1968 Experience of annual productivity dynamics analysis of the spruce stands in the southern taiga biogeocenoses. In *Experimental Investigations of the Taiga Biogeocenoses*. Leningrad: Nauka, pp. 33-49 (in Russian).
- Goryachev, V. M.  
1987 Ecological factors affected on seasonal radial growth dynamics in dark coniferous forests in the Middle Urals. In *Dendrochronological Methods in Forest Science and Ecological Forecasting*. Thesis of the International Workshop. Irkutsk, pp. 194-199 (in Russian, English).
- Gursky, A. V., I. V. Kanevskaya and L. F. Ostapovitsh  
1953 Main results of plant introduction in the Pamir Botanical Garden. *Proceedings of the Pamir Botanical Garden of the Institute of Botany of the Tajik Academy of Sciences*, Vol. 16:1-95 (in Russian).
- Isaev, A. S. and V. V. Kiselev  
1987 *Monochamus urussovi* Fisch. outbreaks investigation: dendroclimatological contribution. In *Dendrochronological Methods in Forest Science and Ecological Forecasting*. Thesis of the International Workshop. Irkutsk, pp. 249-252 (in Russian, English).
- Jashnov, L.  
1925 Climatic cycles and tree growth. *Collection of Scientific Papers of the Kazan Agriculture and Forestry Institute*, No. 1:58-59 (in Russian).

- Javorsky, N. Ch.  
1975 Astro-dynamic-statistic forecasting of climatic conditions in which it is possible the formation of droughts and overwettings. In Questions of Agronomy. Frunze, pp. 118-125 (in Russian).
- Kairiukstis, L. and J. Dubinskaite  
1986 The use of the rhythmic radial tree growth oscillations for prediction of climatic condition variability. In Dendrochronology and Dendroclimatology. Novosibirsk: Nauka, pp. 161-174 (in Russian).
- Kairiukstis, L. A. and A. I. Yudovalkis  
1970 Peculiarities of seasonal tree growth in the light of dendrochronological and dendroclimatology investigations. Lesovedenie, No. 3:29-34 (in Russian).
- Karpenko, A. S. and Ju. O. Medvedev  
1963 Exposure and mapping of fires of various remoteness for aims of the taiga vegetative dynamics study. In Geobotanical Mapping. Moscow - Leningrad: Nauka, pp. 25-30 (in Russian).
- Kishchenko, I. T.  
1978 The cambial activity of pine in habitats of the Karelian region. Thesis on Reports on the 3rd All-Union Conference on Dendroclimatology. Archangelsk, p. 87 (in Russian).
- Kolchin, B. A.  
1963 Dendrochronology of Novgorod. In Materials to Investigations on the USSR Archaeology, No. 117, Vol. 3, pp. 5-103 (in Russian).
- Kolchin, B. A. and N. V. Chernich  
1977 Dendrochronology of the Eastern Europe. Moscow: Nauka, 127 p. (in Russian).
- Kolishchuk, V. G.  
1966 Growth dynamics of mountain pine (*Pinus mughus* Scop.) in connection with solar activity. Proceedings of the USSR Academy of Sciences, Vol. 167, No. 3:710-713 (in Russian).  
1979 Dendrochronological and dendroecological investigations in the Ukrainian Carpathians highlands. In Ecology and Biology of Highland Plants. Novosibirsk: Nauka, pp. 11-17 (in Russian).
- Kolishchuk, V. G., V. V. Mazjak and S. A. Chmara  
1984 The use of archaeological and buried wood for long-term scale construction. In Materials of the 36th Scientific-Technical Meeting of the Lvov Forestry Institute, Lvov, February 1984, Forestry Section. Lvov, pp. 28-31 (Manuscript deposited at CBSTI-forestry, February 26, 1985, No. 3521ch-85 Dep.) (in Russian).
- Kolishchuk, V. G., O. N. Petriv and R. I. Polovnikov  
1975 Spruce growth, its structure, dynamics and conditionality. In Bioecological Principles of Dendrochronology. Vilnius-Leningrad, pp. 31-34 (in Russian).
- Komin, G. E.  
1963 Influence of cyclic fluctuations of the climate on growth and age structure of original bog forests. Proceedings of the Siberian Division of the USSR Academy of Sciences, No. 3, Series of Biological Medical Sciences, No. 12:16-24 (in Russian).  
1970 Cyclicity of trees and stands growth dynamics of pine in the taiga zone of Western Siberia. Proceedings of the Siberian Division of the USSR Academy of Sciences, No. 15, Series of Biological Sciences, No. 3:36-44 (in Russian).  
1972 Cyclicity in the tree growth dynamics. In Problems of vegetative objects examinations. Moscow, pp. 54-64 (in Russian).  
1980 Dendrochronology of Kazim town. In The Historical-Architectural Museum Under the Open Sky. Principles and Methods of Organization. Novosibirsk: Nauka, pp. 121-126 (in Russian).
- Kostin, S. I.  
1940 Repetitiveness of droughts in the Voronez region (on the data of annual growth of European ash). Proceedings of Voronez Agricultural Institute, Vol. 19, No. 1:91-96 (in Russian).  
1960 Climatic changes in the Central forest-steppe region of the Russian Plain. Scientific Notes of the Voronez Forestry Institute, Vol. 21:62-70 (in Russian).  
1963 Repetitiveness of dry and moist periods in the central part of forest-steppe of the Russian Plain. Scientific Notes of the Voronez Forestry Institute, Vol. 29, No. 1:91-101 (in Russian).
- Kristofovitsh, A. T.  
1934 Tree rings of wood as the basis of historical and prehistorical chronologies. Nature, No. 6: p. 77 (in Russian).
- Kuchеров, S. E.  
1987 *Lymantria dispar* outbreaks effect on radial growth of *Quercus robur* in the South Urals. In Dendrochronological Methods in Forest Science and Ecological Forecasting. Thesis of the International Workshop. Irkutsk, pp. 285-288 (in Russian, English).
- Lairand, N. I., N. V. Lovelius and A. A. Yatsenko-Kchmelevsky  
1979 Effects of anthropogenic factors on the growth of Scotch pine (*Pinus sylvestris*) in the Bratsk region. Botanical Journal, Vol. 64, No. 8:1187-1195 (in Russian).

- Laletin, A. P.  
1987 The influence of recreational loads on the radial increment of *Pinus sylvestris* L. subsp. *kulundensis* Suk. In *Dendrochronological Methods in Forest Science and Ecological Forecasting*. Thesis of the International Workshop. Irkutsk, pp. 218-221 (in Russian, English).
- Liseev, A. S.  
1962 About the quantitative characteristic of the influence of climatic factors on tree growth. In *Reforestation and forest cultures*. Vol. 42. Moscow: Selchosizdat, pp. 193-203 (in Russian).
- Litvinenko, V. I.  
1972 Dendroclimatic investigations as the basis for quantitative estimation of struggle measure effectivity with Siberian bombyx. In *Dendroclimatochronology and Radiocarbon*. Kaunas, pp. 205-209 (in Russian).
- Lobzhanidze, E. D.  
1975 The effect of light schedule change on the formation and structure of annual wood rings in even-aged stands. In *Bioecological Principles of Dendrochronology*. Vilnius-Leningrad, pp. 27-30 (in Russian).
- Lovelius, N. V.  
1979 Tree growth variability. Dendroindication of natural processes and anthropogenic impacts. Leningrad: Nauka, 230 p. (in Russian).
- Lukjanova, L. M. and S. M. Mjagkov  
1979 Quantitative estimation of avalanche activity in the Khibin Mountains during the last 2000 years. In *Rhythms of Glacial Processes*. Moscow: Moscow State University Publishing House, pp. 87-112 (in Russian).
- Malokvasov, D. S.  
1974 About the cyclicity in radial tree growth in Korean pine. In *The Increase of Forest Productivity in the Far East*. Chabarovsk, pp. 34-35 (in Russian).
- Mazepa, V. S.  
1986 The use of spectral presentation and linear filtering of the stationary sequences in analysis of cyclicity of dendrochronological series. In *Dendrochronology and Dendroclimatology*. Novosibirsk: Nauka, pp. 49-68 (in Russian).
- Melechov, I. S.  
1948 Effects of fires on the forest. Moscow-Leningrad: Goslestechizdat, 125 p. (in Russian).
- Muchamedshin, K. D.  
1974 Growth dynamics of juniper in the Tian-Shan highlands during the last millenium. In *Proceedings of the 5th All-Union Meeting on the Problem "Astrophysical Phenomena and Radiocarbon"* (Tbilisi, October 4-6, 1973). Tbilisi, pp. 149-161 (in Russian).
- Nesvetailo, V. D.  
1986 Dendrochronological dating of "telegraphic forest" in the area of Tungus meteorite fall. In *Dendrochronology and Dendroclimatology*. Novosibirsk: Nauka, pp. 191-194 (in Russian).
- Olenin, S. M.  
1976 Radial growth of pine in bog pine stands in connection with the secular solar cycles. *Lesovedenie*, No. 2:35-42 (in Russian).  
1977 Radial growth dynamics of pine stands in the middle taiga subzone of the western foot of the Ural Mountains. *Soviet Journal of Ecology*, No. 6:72-75. (in Russian).
- Polyushkin, Yu. V.  
1979 Variability of radial stand growth as a source of information for ecosystem dynamics prediction. In *Problems of Prognostic Investigations of Natural Phenomena*. Novosibirsk: Nauka, pp. 154-167 (in Russian).
- Pshenichnikova, L. S.  
1987 Dendrometrical changes in stand productivity when affected by silvicultural operations. In *Dendrochronological Methods in Forest Science and Ecological Forecasting*. Thesis of the International Workshop. Irkutsk, pp. 295-298 (in Russian, English).
- Rosanov, M. I.  
1965 Dendrochronological method of wood identification. In *Criminalistic and criminal expertise*, No. 103:285-290 (in Russian).  
1968 The subject of criminal dendrochronology. In *Materials of the All-Union Conference on Dendrochronology and Dendroclimatology*. Vilnius, pp. 23-39 (in Russian).
- Rudakov, V. E.  
1951 The method of investigation of the effects of climatic changes on annual ring width in trees. *Proceedings of the Armenian Academy of Sciences*, Vol. 13, No. 3:75-79 (in Russian).  
1958 About the method of investigation of the effects of climatic changes on annual tree ring width. *Botanical Journal*, Vol. 43, No. 12:1708-1712 (in Russian).  
1961 Tree growth and climatic elements. *Proceedings of the USSR Academy of Sciences*, Vol. 137, No. 4:984-985 (in Russian).



- Saborov, R. N.  
1987 Evaluation of the influence of pollutants on forest phytocenoses using the dendrochronological method. *Dendrochronological Methods in Forest Science and Ecological Forecasting. Thesis of the International Workshop. Irkutsk*, pp. 212-217 (in Russian, English).
- Shiyatov, S. G.  
1965 Age structure and formation of low-density larch stands at the upper timberline in the Sob river basin (Polar Urals). *Proceedings of the Institute of Biology of the Ural Branch of the USSR Academy of Sciences*, No. 42:81-86 (in Russian).  
1972 Dendrochronological study of *Picea obovata* in the Lower Taz river. In *Dendroclimatochronology and Radiocarbon*. Kaunas, pp. 76-81 (in Russian).  
1975 The oversecular cycle of indices of *Larix sibirica* at the northern treeline. In *Bioecological Principles of Dendrochronology*. Vilnius-Leningrad, pp. 47-53 (in Russian).  
1980 Dating of wooden buildings of Mangazeia by the dendrochronological method. In Belov, M. I., O. V. Ovsjannikov and V. F. Starkov "Mangazeia. Mangazeia sea way. Part 1". Leningrad: *Gidrometeoizdat*, pp. 93-107, Appendices XXXI and XXXII (in Russian).  
1986 Dendrochronology of the upper treeline in the Urals. Moscow: Nauka, 136 p. (in Russian).
- Shiyatov, S. G. and V. S. Mazepa  
1986 Natural fluctuations of climate in the eastern regions of the USSR based on tree-ring series. In *Collaborative Papers. Regional Resource Management. Vol. 1. Laxenburg, Austria*, pp. 47-73 (in Russian).
- Shiyatov, S. G. and A. V. Uljanov  
1987 Experience in windthrow dating by Dendrochronological methods. In *Dendrochronological Methods in Forest Science and Ecological Forecasting. Thesis of the International Workshop. Irkutsk*, pp. 279-284 (in Russian, English).
- Shvedov, F.  
1892 Tree as a drought chronicle. *Meteorological Vestnik*, No. 5:163-178 (in Russian).
- Tolsky, A. P.  
1936 On the exposure of climatic changes on the basis of tree growth analysis. *Proceedings on Agricultural Meteorology*, No. 24:117-123 (in Russian).
- Turmanina, V. I.  
1971 The use of some morpho-anatomical peculiarities of trees for slope processes indication. In *Theoretical Questions of Phytoindication*. Leningrad: Nauka, pp. 151-154 (in Russian).  
1972 Peculiarities of tree growth near the Elbrus Mountain. In *Dendroclimatochronology and Radiocarbon*. Kaunas, pp. 37-41 (in Russian).  
1979 Dendrochronology of avalanches in the upper reaches of the Baksan Valley. In *Rhythms of Glacial Processes*. Moscow: Moscow State University Publishing House, pp. 128-134 (in Russian).
- Vaganov, E. A., A. S. Isaev and Ju. P. Kondakov  
1972 Dendrochronological analysis of coniferous trees growth damaged by forest insects. In *Dendroclimatochronology and Radiocarbon*. Kaunas, pp. 192-194 (in Russian).
- Vaganov, E. A., A. V. Shashkin, I. V. Sviderskaya and L. G. Vysotskaya  
1985 Histometric analysis of woody plants growth. Novosibirsk: Nauka, 100 p. (in Russian).
- Valentic, E. N. and G. A. Ivanova  
1987 Seasons of extreme fire and their prediction in forests in Siberia. In *Dendrochronological Methods in Forest Science and Ecological Forecasting. Thesis of the International Workshop. Irkutsk*, pp. 259-262 (in Russian, English).
- Voronin, V. I.  
1986 Relation between seed crops and annual growth of Siberian fir in the South Pre-Baikalie (Khamar-Daban Mountain Range). In *Dendrochronology and Dendroclimatology*. Novosibirsk: Nauka, pp. 76-79 (in Russian).
- Yuknees, R. A.  
1987 Anthropogenic changes estimation in trees and stand growth. In *Dendrochronological Methods in Forest Science and Ecological Forecasting. Thesis of the International Workshop. Irkutsk*, pp. 168-173 (in Russian, English).
- Zabelin, O. F.  
1979 The changes in forest stands after fires. In *Forest Fires in Yakutija and its Influence on Forest Nature*. Novosibirsk: Nauka, pp. 87-115 (in Russian).
- Zamotorin, I. M.  
1959 Relative chronology of the Pazirik Borrows. *Soviet Archaeology*, No. 1:21-30 (in Russian).  
1963 About possibility of relative dating of territorially remote constructions by the tree-ring analysis in the Gorny Altai. *Soviet Archaeology*, No. 2:131-138 (in Russian).
- Zaozersky, S. G.  
1934 Towards the methods of retrospective revealing of climatic conditions by the tree growth study. In *Questions of Irrigation*. No. 1, Samarkand, pp. 75-86 (in Russian).

Zviedris, A. and J. Matuzanis

1962 Regularities of ring-width tree variations in the Latvian SSR. Proceedings of the Latvian Academy of Sciences, No. 8:117-128 (in Russian).

Zviedris, A. and R. Sacinieks

1960 About effect of climatic factors on ring-width layers of spruce. Proceedings of the Latvian Academy of Sciences, No. 3:177-184 (in Russian).