

EXPERIMENTAL FOREST WATERSHED STUDIES CONTRIBUTION TO THE EFFECT OF DISTURBANCES ON WATER QUALITY

Daniel G. Neary¹

The most sustainable and best quality fresh water sources in the world originate in forested watersheds (Dissmeyer 2000, Brooks et al. 2003, Barten and Ernst 2004). The biological, chemical, and physical characteristics of forest soils are particularly well suited to delivering high quality water to streams, and moderating the climatic extremes which affect stream hydrology and water quality (Neary et al. 2009). Forest soils are usually characterized by high porosities, low bulk density, and high saturated hydraulic conductivities and infiltration rates (Neary 2011). Consequently, surface runoff is rare in forest environments, and most rainfall moves to streams by subsurface flow pathways where nutrient uptake, cycling, and contaminant sorption processes are rapid. Because of the dominance of subsurface flow processes, peak flows are moderated and baseflows with high water quality are prolonged (Vertessey 1999, Ice and Stednick 2004).

In much of the world, municipalities ultimately rely on forested watersheds to provide adequate quantities of high quality water for continually growing demand (Lee 1980). This is particularly true in semi-arid regions where water supplies are limited, water quality is affected by high mineral content, and human populations are large or growing rapidly. Forest soils provide the ideal conditions for creating high quality water supplies (Neary et al. 2009). Paired catchment research has provided the scientific basis for understanding disturbance effects on water quality and led to the development of Best Management Practices (BMPs) for sustaining water quality (Ice 2004).

Socio-Economic Importance Of Water Quality

The U.S. Geological Survey's NAWQA studies comparing water quality across land uses have demonstrated that forest lands provide the highest quality supplies (Rosen and Lapham 2008). Much of urbanized America depends on forested catchments to provide stable supplies of high quality water. The Colorado River, that supplies water to major metropolitan areas in Arizona, California, and Nevada, derives the bulk of its flow from forested mountain headwater streams. The quality of water is so high from some forested catchments that

cities like Portland, Oregon, utilize unfiltered river water from forested watersheds to supply their residents and businesses (Harr and Fredriksen 1988). New York City has long relied upon forests in the Catskill Mountains to provide one of the highest quality metropolitan drinking water supplies in the world (Pires 2004).

In the USA, over 3400 towns and cities depend on National Forest catchments for their public water supplies (Ryan and Glasser 2000). An additional 3000 administrative sites such as campgrounds, picnic areas, and historical sites rely on the same or similar sources. It has been estimated that 25% of the people in the USA, predominantly in western regions where the bulk of the National Forest lands are located, rely on streams and groundwater emanating from National Forests for their public water supplies. Since 70% of the forest area in the USA is outside the National Forest System, particularly in the eastern USA, a conservative estimate is that 50–75% of the USA's population relies on forest lands to produce adequate supplies of good quality water.

In Canada, the percentage of cities utilizing water from forested catchments is higher due to that country's vast forest area (Bakker 2007). Germany has established Water Conservation Districts (*Wasserschutzgebieten*) for the protection of municipal water supplies (Napier 2000). Australia is one of the driest continents on Earth with less than 1% of the world's freshwater resources. Thus, water supply has always been a major issue for the socio-economic fabric of the country (Pigram 2006). The major cities of Brisbane, Sydney, Canberra, Melbourne, Adelaide, Hobart, and Perth rely on water flow from mostly forested catchments (Foran and Poldy 2002).

Paired Catchment Science

The early 20th Century was unique in that it had the beginnings of paired catchment research in several parts of the world. The Sperbelgraben and Rappengraben experimental catchments were established in 1903 near Emmental, Switzerland (Penman 1963). This was followed by establishment of the Ota watershed study in Japan in 1908 and the Wagon Wheel Gap study in Colorado, USA, in 1910 (Steen 1976, Neary 2000). Paired catchment experiments have been reviewed by a

¹ U.S. Forest Service, Rocky Mountain Research Station, 2500 South Pine Knoll Drive, Flagstaff, AZ 86001; dneary@fs.fed.us

number of authors (Bosch and Hewlett 1982, Binkley and Brown 1993, Sahin and Hall 1996, Stednick 1996, Neary 2002, Andréassian 2004, Brown et al. 2005). Most of these reviews have dealt with the topic of water yield. However, many of the paired catchment experiments initially designed for water yield research were expanded to include water quality.

This paper is based on a publication produced for a conference on at a meeting of the International Association of Hydrological Sciences in Melbourne, Australia, in 2011 (Neary 2012). It provides a historical perspective of some of the many accomplishments of water quality research over the past century made possible by using the paired catchment methodology. It also examines current research efforts, and makes recommendations about future research directions for water quality science.

PAST CATCHMENT WATER QUALITY RESEARCH

Water quality is a relative concept that is based on measurable physical, chemical, and biological characteristics in relation to specific uses such as human consumption, crop irrigation, livestock watering, fisheries maintenance, and recreational usage (Scatena 2000). As water moves through a catchment, it interacts with ecosystem components and transports dissolved gases, cations and anions, organic compounds, trace metals, sediment particles, and microscopic organisms. The principle parameters of concern in water quality are bacteria and other microorganism contents, suspended sediment, organic matter, dissolved oxygen, nitrogen (N) compounds, anions and cations, trace metals, salts, and pesticides. Temperature, color, and turbidity are physical parameters that round out the concept of water quality. These contaminants and physical conditions then define water quality based on the uses of water mentioned above (Gleick 1993). Understanding of the movement of water through forest catchments provided the scientific foundation for evaluating water quality in both natural and disturbed forests (Swanson et al. 2000). In the USA, there was early interest in water quality research at locations like the Coweeta Hydrologic Laboratory in the 1930s and 1940s, but it did not become well established until the middle of the 20th century when water pollution legislation such as the Federal Water Pollution Act of 1948 was enacted (Douglass and Swank 1975, Swank et al. 2001). By the time the Clean Water Act was enacted in the USA (1972) there was considerable interest in water quality. This aspect of catchment science became fully integrated into hydrologic research.

Natural Processes

Although the initial focus of early catchment research was water yield, the adoption of the paired catchment approach set the stage for examining physical, chemical, and biological processes that controlled nutrient cycling and other water quality related functions of forest catchments (Bormann and Likens 1967). The untreated half of catchment study pairs provided the opportunity to study natural processes that controlled water quality. However, the disturbances to these processes produced by practices such as harvesting, site preparation, road construction, fire, fertilization, herbicide use and insect outbreaks provided the real insight into natural catchment processes that affect water quality.

Disturbance Effects

Most of the forest catchment water quality studies reported in the literature deal with tree harvesting and post-harvest site preparation since much of the early interest in paired catchment science related to vegetation management to increase water yield. In addition, these practices were considered to produce the most disruptions to ecological processes and therefore the most influence on water quality. Since forest fertilization has been a basic feature of intensive forest management throughout the world, the impact of fertilizers on water quality has been an issue easily addressed by paired catchment research (Binkley et al. 1999). Paired catchments provided a sound basis for acid deposition research in the 1980s and 1990s (Likens et al. 1996), and continue to support scientific endeavors on climate change in the 21st Century (Bouraouii et al. 2004).

A number of water quality parameters are affected by disturbances, but only nutrients, sediments, and temperature will be discussed in the limited space available for this paper. Other papers present a much more detailed discussion of these topics (Binkley and Brown 1993, Swanson et al. 2000, Neary 2002, Ice and Stednick 2004).

Harvesting and Site Preparation – Nutrients: Neary (2002) summarized a number of paired catchment studies looking at N losses in streamflow after harvesting and site preparation (Tables 1 and 2). Nitrate nitrogen (NO₃-N) dynamics are considered to be very susceptible to disturbance and NO₃-N concentration is a commonly accepted indicator of catchment health and water quality throughout the world since low levels (10 mg L⁻¹) can affect infant health (Neary 2002). For the most part, large increases in NO₃-N levels in streams draining harvested catchments have not been observed (Tables

Table 1. Paired catchment comparison of the effects of forest harvesting on mean NO₃-N concentrations in streamflow in North America the year after cutting (Adapted from Neary 2002).

Forest Type	Location	NO ₃ -N		Reference
		Uncut	Cut	
		mg L ⁻¹	mg L ⁻¹	
Lodgepole Pine	Alberta, Canada	0.2	0.7	Singh & Kalra 1975
Spruce, Fir	British Columbia, Canada	0.1	0.2	Hetherington 1976
Spruce, Fir	British Columbia, Canada	<0.1	0.5	Feller & Kimmons 1984
Spruce, Fir	Colorado, USA	<0.1	<0.1	Stottlemeyer 1992
Slash pine	Florida, USA	<0.1	0.3	Riekerk et al. 1980
Loblolly Pine	Georgia, USA	0.1	0.1	Hewlett & Doss 1984
Mixed Conifer	Idaho, USA	0.2	0.2	Snyder et al. 1975
Aspen, Birch, Spruce	Minnesota, USA	0.1	0.2	Verry 1972
Mixed Conifer	Montana, USA	0.1	0.2	Bateridge 1974
Northern Hardwoods	New Brunswick, Canada	0.1	0.6	Krause 1982
Northern Hardwoods	New Hampshire, USA	0.3	11.9	Pierce et al. 1970
Mixed Hardwoods	North Carolina, USA	<0.1	0.1	Swank et al. 2001
Spruce, Fir, Pine	Nova Scotia, Canada	<0.1	0.3	Vaidya et al. 2008
Douglas-fir	Oregon, USA	<0.1	0.2	Fredrickson et al. 1975
Mixed Conifers	Oregon, USA	<0.1	0.2	Fredrickson et al. 1975
Oak-Maple	Pennsylvania, USA	0.1	5.0	Corbett et al. 1975
Spruce, Fir, Pine, Birch	Quebec, Canada	<0.1	<0.1	Carignan et al. 2000
Loblolly Pine	South Carolina, USA	<0.1	<0.1	Van Lear et al. 1985
Mixed Hardwoods	West Virginia, USA	0.1	0.5	Aubertin & Patric 1974

Table 2. Paired catchment comparison of the effects of forest harvesting on mean NO₃-N concentrations in streamflow in Europe, Africa, Asia, and the South Pacific the year after cutting (Adapted from Neary 2002).

Forest Type	Location	NO ₃ -N		Reference
		Uncut	Cut	
		mg L ⁻¹	mg L ⁻¹	
Native Beech-Podocarp	Chile	<0.1	<0.1	Oyarzun et al. 2007
Spruce, Fir, Peat	Finland	<0.1	0.1	Ahtiainen & Huttunen 1999
Spruce, Fir, Beech	Germany	0.7	1.0	Bäumler & Zech 1999
Native Hardwoods [#]	Japan	0.7	1.6	Ohrui & Mitchell (1998)
Radiata Pine	New Zealand	<0.1	0.5	Graynoth 1979
Beech-Podocarp	New Zealand	<0.1	<0.1	O'Loughlin et al. 1980
Radiata Pine	New Zealand	<0.1	0.2	O'Loughlin 1994
Evergreen Forest/Scrub	South Africa	<0.1	0.1	Scott & Lesch 1996
Pine, Spruce, Hardwood	Sweden	0.1	0.2	Rosen 1996
Spruce, Moor	United Kingdom	0.2	0.3	Neal et al. 1992
Eucalyptus spp.	Victoria, Australia	<0.1	<0.1	Hopmans & Bren 2007

[#] 4 years after cutting

1 and 2). Certainly there is no general indication that the World Health Organization (2006) water quality standard (10 mg L⁻¹ NO₃-N) is commonly breached by post-harvesting NO₃-N concentration increases. The largest increase reported (Table 1, Pierce et al. 1970) was measured in an experiment where herbicides were used to suppress vegetation regrowth. Other causes of increased NO₃-N losses in forested catchments

have been documented where severe fire occurred (DeBano et al. 1998), nitrogenous fertilizers were used during regeneration (Neary and Hornbeck 1994), or N saturation of ecosystems has reached a critical level due to atmospheric deposition (Aber et al. 1989). Paired catchments have been instrumental in demonstrating that, except in rare instances of delayed vegetation regrowth (e.g. Pierce et al. 1970) or forest ecosystems impacted by

Table 3. Effects of harvesting and related disturbances on sediment outputs from paired catchments (Adapted from Neary 2002, Diaz-Chavez 2011).

Forest Type	Location	Treatment	Sediment Increase	Reference
			Mg ha ⁻¹ yr ⁻¹	
Mixed Conifers	Arizona, USA	Clearcut	0.003	Heede 1987
Mixed Conifers	Arizona, USA	Cut, Road	0.081	Heede 1987
Loblolly Pine	Arkansas, USA	Clearcut	0.225	Beasley & Granillo 1988
Slash Pine	Florida, USA	Clearcut, Bed	0.033	Riekerk et al. 1980
Tropical Forest	Malaysia	Clearcut, Skid	1.200	Malmer 1990
Northern Hardwoods	New Hampshire, USA	Clearcut	0.323	Hornbeck et al. 1987
Beech, Podocarps	New Zealand	Clearcut	0.182	O'Loughlin et al. 1980
Beech, Podocarps	New Zealand	Clearcut, Skid	3.003	O'Loughlin et al. 1980
Douglas-fir	Oregon, USA	Clearcut	0.510	Beschta 1978
Loblolly Pine	South Carolina, USA	Clearcut	0.131	Van Lear et al. 1985
Native Forest	Peru	Clearcut	0.421	Plamondon et al. 1991
Spruce, Douglas-fir	United Kingdom	Clearcut	0.327	Leeks & Roberts 1987
Eucalyptus spp.	Victoria, Australia	Clearcut	0.026	Grayson et al. 1993

atmospheric deposition (Bäumler and Zech 1999), forest harvesting does not significantly raise stream NO₃-N or other nutrient concentrations for long periods of time.

Harvesting and Site Preparation – Sediment:

After forest harvesting, forest catchments produce sediments yields that are highly variable depending on factors such as soils, climate, topography, ground cover, road networks, and catchment condition (Rosen 1984). Although sediment yields increase after harvesting due to the physical disturbance of soil, they are usually transient due to vegetation re-growth (Neary 2002). There is a large body of literature that reported using paired catchments to assess the effects of harvesting and site preparation on the sediment component of water quality (Binkley and Brown 1993, Neary 2002). The largest increases documented in the literature have been associated with post-harvest mechanical site preparation (Beasley 1979), slope instability (O'Loughlin and Pearce 1976), road construction and maintenance (e.g. O'Loughlin et al. 1980, Swanson et al. 1986, Heede 1987), highly erosive soils (Beasley and Granillo 1988), and steep terrain (Beschta 1978) (Table 3).

Sediment movement to and within stream systems is a constant environmental concern in managed forest catchments, but it also occurs naturally without management. Herein rests the importance of paired catchment analyses. Catchments can vary greatly in their natural suspended and bedload sediment characteristics (Trimble and Crosson 2000). Both natural and

anthropomorphic erosion material can be re-entrained after initial deposition in ephemeral or perennial stream channels, and move downstream with streamflow for long time periods and distances. The cumulative effects of erosion and sedimentation that occurred centuries ago from agriculture or forestry can present forest managers with many challenges (Terrene Institute 1993). Sediment is an important water quality parameter since it can harm aquatic organisms and habitats, and render water unacceptable for drinking water supply or recreation purposes (Table 3). However, adequate BMPs can significantly limit increases in sediment delivery to streams (Grayson et al. 1993, Neary et al. 2011).

Harvesting and Site Preparation – Temperature:

Forest vegetation shades stream channels from solar radiation, thereby producing stream temperatures that are cooler and less variable than for unshaded sites. Increases in temperature that result from forest harvesting affect physical, chemical, and biological processes. Thus, temperature is a critical water quality characteristic of many streams and aquatic habitats. Temperature controls the survival of certain flora and fauna in the water that are sensitive to water temperature. The removal of streambank vegetation by burning can cause water temperature to rise, causing thermal pollution to occur, which in turn can increase biological activity in a stream (DeBano et al. 1998, Brooks et al. 2003). Increases in biological activity place a greater demand on the dissolved oxygen content of the water, one of the more

Table 4 Paired catchment studies of the effects of forest harvesting on stream temperature (Adapted from Binkley and Brown 1993, Binkley et al. 1999, Moore et al. 2005).

Location	Temperature			Time	Reference
	Control	Cut	Change		
	°C	°C	°C		
Clear Cut, No Buffer					
Brit. Columbia, Canada	17.9	21.8	3.9	Mean Annual	Feller 1981b
New Hampshire, USA	16.0	20.0	4.0	Mean Daily 30 Days AUG	Likens et al. 1970
New Zealand	12.4	15.0	2.6	Mean Annual	Quinn & Stroud 2002
North Carolina, USA	18.3	21.7	3.4	Mean Daily 30 Days AUG	Swift & Messer 1971
Oregon, USA	13.3	15.6	2.3	1 Day - JUL	Brown et al. 1971
Oregon, USA	20.6	28.3	7.7	1 Day - JUL	Brown et al. 1971
Oregon, USA	14.4	22.8	8.4	Mean Daily 30 Days AUG	Levno & Rothacher 1969
Oregon, USA	12.2	22.2	10.0	Mean Daily 30 Days AUG	Brown & Krygier 1970
Pennsylvania, USA	17.8	25.0	7.2	Mean Daily 30 Days AUG	Rishel et al. 1982
Clear Cut, With Buffer					
Georgia, USA	21.1	25.0	3.9	Mean Daily 30 Days AUG	Hewlett & Fortson 1982
New Zealand	16.8	20.0	3.2	Mean 10 Minute OCT-JAN	Boothroyd et al. 2004
Oregon, USA	14.4	15.0	0.6	1 Day - July	Brown et al. 1971
Oregon, USA	16.7	18.3	1.6	1 Day - July	Brown et al. 1971
West Virginia, USA	14.4	16.1	1.7	Mean Weekly – Growing Season	Aubertin & Patric 1974
Partial Cut With Buffer					
Pennsylvania, USA	19.4	20.6	1.2	Mean Daily 30 Days AUG	Rishel et al. 1982
Oregon, USA	12.0	15.0	3.0	Mean Daily 21 Days AUG	Harr & Fredriksen 1988
Oregon, USA	12.5	14.4	2.0	Mean Monthly Maximum	Harris 1977
Tasmania, Australia	8.2	9.0	0.8	3 Year Mean	Ringrose et al. 2001

important water quality characteristics from a biological perspective.

In the USA there are no established national standards for the temperature of drinking water (Dissmeyer 2000). However, under the Clean Water Act, States are required to develop water quality standards to protect beneficial uses such as fish habitat and water quality restoration. The U.S. Environmental Protection Agency provides oversight and approval of these State standards. One of the problems with these standards is identifying natural temperature patterns caused by vegetation, geology,

geomorphology, climate, season, and natural disturbance history. Also, increases in stream water temperatures can have important and often detrimental effects on stream eutrophication. Acceleration of stream eutrophication can adversely affect water quality by adversely affecting the color, taste, and smell of drinking water. Severe wildfires can function like streamside timber clearcuts in raising the temperature of streams due to direct heating of the water surface (Neary et al. 2005, Table 4).

Forest Fertilizers: Forest fertilization is another management disturbance that has the potential to

Table 5. Paired catchment comparison of the effects of forest fertilization on maximum NO₃-N concentrations in streamflow (Adapted from Binkley and Brown 1993, Binkley et al. 1999).

Location	Temperature			Time	Reference
	Control	Cut	Change		
	°C	°C	°C		
Clear Cut, No Buffer					
Brit. Columbia, Canada	17.9	21.8	3.9	Mean Annual	Feller 1981b
New Hampshire, USA	16.0	20.0	4.0	Mean Daily 30 Days AUG	Likens et al. 1970
New Zealand	12.4	15.0	2.6	Mean Annual	Quinn & Stroud 2002
North Carolina, USA	18.3	21.7	3.4	Mean Daily 30 Days AUG	Swift & Messer 1971
Oregon, USA	13.3	15.6	2.3	1 Day - JUL	Brown et al. 1971
Oregon, USA	20.6	28.3	7.7	1 Day - JUL	Brown et al. 1971
Oregon, USA	14.4	22.8	8.4	Mean Daily 30 Days AUG	Levno & Rothacher 1969
Oregon, USA	12.2	22.2	10.0	Mean Daily 30 Days AUG	Brown & Krygier 1970
Pennsylvania, USA	17.8	25.0	7.2	Mean Daily 30 Days AUG	Rishel et al. 1982
Clear Cut, With Buffer					
Georgia, USA	21.1	25.0	3.9	Mean Daily 30 Days AUG	Hewlett & Fortson 1982
New Zealand	16.8	20.0	3.2	Mean 10 Minute OCT- JAN	Boothroyd et al. 2004
Oregon, USA	14.4	15.0	0.6	1 Day - July	Brown et al. 1971
Oregon, USA	16.7	18.3	1.6	1 Day - July	Brown et al. 1971
West Virginia, USA	14.4	16.1	1.7	Mean Weekly – Growing Season	Aubertin & Patric 1974
Partial Cut With Buffer					
Pennsylvania, USA	19.4	20.6	1.2	Mean Daily 30 Days AUG	Rishel et al. 1982
Oregon, USA	12.0	15.0	3.0	Mean Daily 21 Days AUG	Harr & Fredriksen 1988
Oregon, USA	12.5	14.4	2.0	Mean Monthly Maximum	Harris 1977
Tasmania, Australia	8.2	9.0	0.8	3 Year Mean	Ringrose et al. 2001

affect stream water quality because of the additions of N, phosphorus (P), cations, etc. to forest catchments (Binkley et al. 1999, Neary 2002). Streams originating in agricultural areas have about 9 times the load of N and P than forested catchments so the water quality of forested areas is highly valued. The growth of tree plantations in high production silviculture regions of the world is often limited by soil nutrient availability (Fox et al. 2007). Hence, fertilization is a common silvicultural practice in these high-intensity production forests. Fertilizer applications are rarely incorporated in stand management

in slower growing forests due to economic limitations. Nitrogen and P fertilizers are the most frequently used but, in some locations, cations and micronutrients are applied to deal with local deficiencies. Here again, paired catchments have been invaluable in understanding the water quality implications of this management practice (Table 5). Higher stream concentrations are usually associated with higher fertilizer application rates (e.g. >200 kg-N ha⁻¹) (Smith et al. 1994) or aerial applications that fly over or near monitored streams (Grip 1982, Hetherington 1985, Helvey et al. 1989, Göethe et al.

Table 6. Paired catchment comparisons of sediment yield the first year after prescribed fires and wildfire (Adapted from Neary et al. 2005, Shakesby and Doerr 2006, and Smith et al. 2011)

Vegetation	Location	Fire Type	Sediment Yield		Reference	
			Control	Post-Fire		
			Mg ha ⁻¹ yr ⁻¹	Mg ha ⁻¹ yr ⁻¹		
Ponderosa Pine	Arizona, USA	WF Low ¹	0.003 ^C	0.080	Campbell et al. 1977	
Ponderosa Pine	Arizona, USA	WF Mod ¹	0.003 ^C	0.300	Campbell et al. 1977	
Ponderosa Pine	Arizona, USA	WF High ¹	0.003 ^C	1.254	Campbell et al. 1977	
Chaparral	Arizona, USA	Wildfire	0.175 ^C	204.000	Glendening et al. 1961	
Chaparral	Arizona, USA	Wildfire	0.096 ^C	28.694	Pase & Ingebo 1965	
Shortleaf Pine	Arkansas, USA	Prescribed	0.036 ^C	0.237	Miller et al. 1988	
Mixed Conifer	B.C. Canada	Wildfire	0.005 ^S	0.008 ^S	Petticrew et al. 2006	
Chaparral	California, USA	Wildfire	0.880 ^C	146.000	Krammes 1960	
Ponderosa Pine	California, USA	Prescribed	<0.00 ^C	<0.001	Biswell & Schultz 1965	
Chaparral	California, USA	Wildfire	0.043 ^C	28.605	Wells et al. 1981	
Ponderosa Pine	Colorado, USA	Wildfire	0.280 ^P	68.000	Moody & Martin 2001	
Oak spp.	Mississippi, USA	Prescribed	0.470 ^C	1.142	Ursic 1970	
Hardwoods	Oklahoma, USA	Prescribed	0.022 ^C	0.246	Daniel et al. 1943	
Fynbos Scrub	South Africa	Prescribed	0.013 ^C	0.420	Scott 1993	
Fynbos	South Africa	Wildfire	0.013 ^C	7.800	Scott 1993	
Loblolly Pine	South Carolina, USA	Prescribed	0.027 ^C	0.042	Van Lear et al. 1985	
Loblolly Pine	Texas, USA	Prescribed	0.112 ^C	0.806	Pope et al. 1946	
Juniper 0%	Texas, USA	Prescribed	0.025 ^C	0.029	Wright et al. 1976	
Slope						
Juniper 15-20%	Texas, USA	Prescribed	0.076 ^C	1.874	Wright et al. 1976	
Juniper 43-54%	Texas, USA	Prescribed	0.013 ^C	8.443	Wright et al. 1976	
Eucalyptus spp.	Victoria, Australia	Wildfire	0.230 ^P	2.960	Lane et al. 2006	
Radiata Pine	Victoria, Australia	Wildfire	0.744 ^P	12.300	Smith et al. 2012	
Eucalyptus spp.	Victoria, Australia	Wildfire	0.057 ^P	0.110	Smith et al. 2012	
Mixed Conifer	Washington, USA	Wildfire	0.008 ^P	0.120	Helvey 1980	

^P Pre-fire data; ^C Control catchment data; ¹ Fire Severity: Low Moderate, High; ^S Seasonal Suspended Sediment

1993). Nitrogen saturation of soils from atmospheric deposition (Aber et al. 1989) can predispose forest stands to leak highly mobile NO₃-N if it is not utilized by vegetation (Pierce et al. 1970). Paired catchments provide investigators the ability to sort out fertilizer water quality effects from those produced by other processes (e.g. herbicide suppression of vegetation regrowth, N saturation of soils, naturally high N soils, inputs from agricultural areas, etc.).

Roads: Best Management Practices for roads are most effective on minimizing sediment impacts to water quality when properly planned and implemented prior to, during, and after harvesting (Neary et al. 2011). Most of these guidelines relate to designing, constructing, and maintaining major access roads, logging roads, skid trails, and landings. Permanent roads and associated temporary

roads are the primary sources for 90% of the sediment generated by harvesting (Swift 1988). The underlying principles of road BMP guidelines are to minimize disturbances in streamside zones, reduce the erosive power of runoff on bare road surfaces, and to maintain the normally high infiltration capacity of forest soils (Neary et al. 2011). Lane and Sheridan (2002) clearly pointed out the role of stream crossings in routing sediment into streams that Swift (1988) alluded to. They employed an above and below approach rather than a paired catchment method. In Lane and Sheridan's (2002) study of unsealed road stream crossings in Victoria, Australia, suspended sediments increased 3.5 times below crossings.

Fire: A major disturbance to catchment hydrology, geomorphology, and water quality in fire-prone regions like the western USA, the Mediterranean Basin, and

Australia is wildfire (Shakesby and Doerr 2006). The random nature of wildfires and their characteristic severities rarely gives researchers the opportunity to use paired catchment techniques to assess impacts on water quality. Prescribed fires are much more amenable to paired catchment comparisons because they are easier to manage. However, even the best managed paired catchment study of prescribed fire can produce surprises (Gottfried et al. 2012). Wildfire impacts on water quality evaluations reported in the literature have been a mixture of paired catchment methods and before-fire and after-fire approaches using the same catchment (DeBano et al. 1998, Neary et al. 2005, Smith et al. 2011).

Post-fire sediment yields can vary widely from <0.001 to over $204 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ depending on the type of fire (prescribed or wildfire), fire severity, topography, fuel type, and climate (Table 6). The highest soil erosion values usually involve intense rainfall on steep terrain (Glendening et al. 1961, Moody & Martin 2001, Neary et al. 2012). Wright et al. (1976) demonstrated the effect of slope with his study in juniper stands in Texas. As slope increased from zero to the 43-54 % range, the annual prescribed fire sediment losses rose from about 0.029 to $8.443 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ compared to a range of 0.013 to $0.025 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ in unburned paired catchments.

Usually post-fire maximum $\text{NO}_3\text{-N}$ levels are in the 0.1 to 0.6 mg L^{-1} range since wildfires volatilize most of the N in the fuels they consume and prescribed fires are usually low-level disturbances (Neary et al. 2005). One of the few and the most striking response of water quality streamflow to fire was observed in southern California, where N loadings from atmospheric deposition are relatively high, and the frequent wildfires in the chaparral shrublands are characterized by high fire severity (Riggan et al. 1994). Severe burning of a catchment in the Mediterranean-type chaparral resulted in a maximum $\text{NO}_3\text{-N}$ level of 15.3 mg L^{-1} in streamflow compared to 2.5 mg L^{-1} peak in streamflow from an unburned control watershed. The maximum concentration for a moderately burned catchment was 9.5 mg L^{-1} . These results represent an "unusual response" because the catchments studied were subject to a chronic atmospheric deposition of air pollutants from the Los Angeles basin that are among the highest recorded in the USA.

Pesticides: Another water quality parameter of considerable concern that has been amenable to study with the paired catchment approach is herbicide and insecticide residue environmental fate. Michael and Neary (1993) discussed this topic in considerable detail. A study by Neary et al. (1983) that utilized four

1.0 ha chemically-treated catchments plus an untreated control was adopted as a template for required herbicide registration studies in the USA by the U.S. Environmental Protection Agency. Since that study, paired watersheds have been an integral part of forestry pesticide environmental fate studies the past three decades (Neary et al. 1993). Virtually all monitoring protocols now require use of untreated control watersheds. Any future research on newly developed forestry pesticides must incorporate paired-watershed methodology. Despite the frequent criticisms of pesticides like herbicides, they should be kept as tools that can achieve vegetation or other pest management goals and maintenance of water quality (Neary and Michael 1996).

Additional research conducted by Michael and Neary (1993) and Neary et al. (1993) expanded on the work of Norris (1970). However, it incorporated newer, rapidly degrading pesticides. They enhanced earlier findings regarding the importance of forest soils in protecting water quality. Neary et al. (1993) and Neary and Michael (1996) concluded that the risks to water quality posed by modern silvicultural chemicals is very low due to the low toxicity of the chemicals, infrequent use over the rotations of conventional forest stands, the lack of bioaccumulation by these pesticides, and the function of forest soil organic matter and microorganisms in adsorbing and decomposing pesticide residues. If forest pesticides are not applied directly to water, their tendency to migrate into streams is limited by forest soil biological and chemical processes. Although herbicides, especially water soluble ones like picloram and hexazinone, have been measured to move through forest soils, they do so in small non-toxic amounts because of the biological and chemical actions of organic matter in forest soils (Neary et al., 1985).

PRESENT CATCHMENT WATER QUALITY RESEARCH

The present level of paired catchment water quality research is quite varied world-wide depending on funding levels and personal commitments and dedication of individual scientists. As an example, Neary et al. (2012) mentioned 180 gauged catchments that are currently active in the U.S. Forest Service's Experimental Forests and Ranges network. However not all of these catchments are engaged in providing data for water quality research. Some topics researched in the past have been suspended because the forestry practices are not being used or decisions were made that sufficient information exists on the topic. These include clearcutting, fertilizer use, road construction, and pesticide application.

Present water quality science is more aligned with the objectives of long-term programs like the Long Term Ecological Research (LTER) program that has international sites, and the Europe-based International Cooperative Program on Assessment and Monitoring of Air Pollution (ICP2). A series of new ICP2 sites added to the network in the USA are based on U.S. Forest Service Experimental Forest and Range sites that have actively-gauged paired catchments. Paired catchments also can be used for climate change research. Although they do not provide the control vs treated comparisons in a classical sense, they can be used for replications within an ecosystem and comparisons between sites/ecosystems.

Fire impacts on water quality are currently another major area of concern because of adverse effects on municipal water supplies (Neary et al. 2005, Smith et al. 2011). Prescribed fire effects on water quality are being studied at sites that have paired catchments (Gottfried et al. 2012). Wildfires rarely present the opportunity to use a paired catchment approach to study water quality effects because of the random nature of these events and their unpredictability.

FUTURE WATER QUALITY RESEARCH USING PAIRED CATCHMENTS

The direction of future water quality research using paired catchments will depend greatly on governmental support for the science. This type of water quality research is expensive and it requires the commitment to the long-term that only government entities can afford. Water quantity and quality are going to be increasingly important topics as nations come to grips with water security problems (Vose et al. 2011). Human populations are increasing most in regions where the abundance of good quality water is being affected by climate change. The importance of long-term studies will loom large since these studies are good indicators of climate change and its effects (Archer and Predick 2008). Specific topics that require further water quality investigation include wildfire, fire retardant use, atmospheric deposition, trace organic chemicals, oil development, large-scale mining, inter-basin water diversions, and bioenergy.

Water is now an area of keen interest in bioenergy development because of its potential footprint on water supplies and its effects on water quality. Some recent publications have addressed the latter issue (Diaz-Chavez et al. 2011a, 2011b). Currently, BMPs offer the best solution to achieving the goal of energy production with biofuels that minimizes the impact on water quality. In some instances, sound research using the paired

catchment approach will be needed to convince regulatory authorities that bioenergy feedstock production can co-exist with water quality goals and standards.

Over the span of the 20th Century, the perception of what constitutes watershed management and hydrologic science has grown considerably. At the beginning of the century, it was mostly concerned with the development and maintenance of water supplies. Water quality was a big issue then and it still is. At the beginning of the 21st Century, it is probably best defined as a comprehensive understanding of the components of watersheds and their physical, chemical, and ecological interactions to produce high quality water in sufficient supply to meet human demands (Reimold 1998). This definition also reflects thinking on the discipline at the end of the 20th Century that watershed management and hydrologic science incorporates the holistic approach to a watershed as an ecosystem, and not just manipulation of physical processes. The goal of watershed management is to assess the effects of current and future land uses on soil and water resources, determine the potential social and ecological impacts, and provide solutions to watershed problems.

As Rango (1995) pointed out, the increase in the world's human population (now at 7 billion) will cause the demand, scarcity, price, and need for high quality water to expand on a global scale into the foreseeable future. He forecast that, in this era of "Global Hydrology" for hydrologic science and watershed management, worldwide emphasis will be placed on large area assessments using modeling, remote sensing, watershed management expertise, and the best hydrologic science. The technological tools and paired catchment infrastructure are in place. The key to the future success of these endeavors lies in watershed management professionals using their expertise and understanding of paired catchment science to develop positive outcomes for human populations of all countries.

SUMMARY AND CONCLUSIONS

This paper provides an overview of the role of paired catchment science in the understanding of water quality in forest ecosystems. Many cities throughout the world rely on forested catchments for the source of their water supplies. This resource is constantly being stressed due to burgeoning demand for high quality water. The current level of knowledge of how forest management has affected water quality has been dependent on science conducted on a large array of catchment studies in mainly Europe, North America, Asia and the South

Pacific. These studies have been expensive to establish in the past, but their value to future water resources science is invaluable. They need to be supported to address important science topics and water management concerns of the 21st century.

REFERENCES

- Aber, J.D.; Nadelhoffer, K.J.; Steudler, P.; Melillo, J.M. 1989. Nitrogen saturation in northern forest ecosystems. *Bioscience* 39: 378-386.
- ACTEW. 2011. Water networks. <http://www.actewagl.com.au/water/networks/default.aspx> (accessed: 04-20-2011)
- Adams, M.A.; Attiwill, P.M. 1991. Nutrient balance in forests of northern Tasmania. 2. Alteration of nutrient availability and soil water chemistry as a result of logging, slash-burning and fertilizer application. *Forest Ecol. Manage.*, 44: 115-131.
- Ahtiainen, M.; Huttunen, P. (1999) Long-term effects of forestry managements on water quality and loading in brooks. *Boreal Environ. Res.* 4: 101-114.
- Andréassian, V. 2004. Water and forests: from historical controversy to scientific debate. *J. Hydro.* 291: 1-27.
- Archer, S.R.; Predick, K. I. 2008. Climate change and ecosystems of the southwestern United States. *Rangelands* 30: 23-28.
- Aubertin, G.M.; Patric, J.H. 1974. Water quality water after clearcutting a small watershed in West Virginia. *J. Environ. Qual.* 3: 243-249.
- Bakker, K.J. 2007. *Eau Canada: The future of Canada's water*. University of British Columbia Press, Vancouver, BC, Canada, 417 p.
- Barten, P.K.; Ernst, C.E. 2004. Land conservation and watershed management for source protection. *J. Amer. Water Works Assoc.* 96: 121-135.
- Bateridge, T.E. 1974. Effects of clearcutting on water discharge and nutrient loss. Bitterroot National Forest, Montana. M.S. Thesis. Office of Water Resources Research, University of Montana, Missoula, MT. 68 p.
- Bäumler, R.; Zech, W. 1999. Effects of forest thinning on the streamwater chemistry of two forest watersheds in the Bavarian Alps. *Forest Ecol. Manage.* 116: 119-128.
- Beasley, R.S. 1979. Intensive site preparation and sediment losses on steep watersheds in the Gulf Coastal Plain. *Soil Sci. Soc. Am. J.* 43: 412-417.
- Beasley, R.S.; Granillo, A.B. 1988. Sediment and water yields from managed forests on flat coastal plain sites. *Water Resour. Bull.* 24: 361-366.
- Beschta, R.L. 1978. Long-term patterns of sediment production following road construction and logging in the Oregon Coast Range. *Water Resour. Res.* 14: 1011-1016.
- Binkley, D.; Brown, T.C. 1993. Forest practices as nonpoint sources of pollution in North America. *Water Resour. Bull.* 29: 729-740.
- Binkley, D.; Burnham, H.; Allen, H.L. 1999. Water quality impacts of forest fertilization. *Forest Ecol. Manage.* 121: 191-213.
- Biswell, H.H.; Schultz, A.M. 1965. Surface runoff and erosion as related to prescribed burning. *J. Forest.* 55: 372-373.
- Boothroyd, I.K.G.; Quinn, J.M.; Langer, E.R.; Costley, K.J.; Steward, G. 2004. Riparian buffers mitigate effects of pine plantation logging on New Zealand streams 1. Riparian vegetation structure, stream geomorphology and periphyton. *Forest Ecol. Manage.* 194: 199-213.
- Bormann, F.H.; Likens, G.E. 1967. Nutrient cycling. *Science* 155: 424-429.
- Bosch, J.M.; Hewlett, J.D. 1982. A review of catchment experiments to determine the effect of vegetation changes on water yield and evapotranspiration. *J. Hydro.* 55: 3-23.
- Bouraouii, F.; Grizzetti, B.; Granlund, K.; Rekolainen, S.; Bidoglio, G. 2004. Impact of climate change on the water cycle and nutrient losses in a Finnish catchment. *Climate Change* 66: 109-126,
- Brooks, K.N.; Ffolliott, P.F.; Gregersen, H.M.; DeBano, L.F. 2003. *Hydrology and the management of watersheds*. Ames, Iowa State Press. 574 p.
- Brown, G.W.; Krygier, J.T. 1970. Effects of clear-cutting on stream temperatures. *Water Resour. Res.* 6: 1133-1139.
- Brown, G.W.; Swank, G.W.; Rothacher, J. 1971. Water temperature in the Steamboat Drainage. USDA Forest Service Res. Pap. PNW-119, Pacific Northwest Forest and Range Experiment Station, Portland, Oregon. 17 p.
- Brown, A.E.; Zhang, L.; McMahon, T.A.; Western, A.W.; Vertessy, R.A. 2005. A review of paired catchment studies for determining changes in water yield resulting from alterations in vegetation. *J. Hydro.* 310: 28-61.
- Campbell, R.E.; Baker, M.B., Jr.; Ffolliott, P.F.; Larson, F.R.; Avery, C.C. 1977. Wildfire effects on a

- ponderosa pine ecosystem: an Arizona case study. USDA Forest Service Res. Pap. RM-191, Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado. 12 p.
- Carignan, R.; D'Arcy, P.; Lamontagne, S. 2000. Comparative impacts of fire and forest harvesting on water quality in Boreal Shield lakes. *Can. J. Fish. Aquat. Sci.* 57: 105-117.
- Corbett, E.S.; Lynch, J.A.; Sopper, W.E. 1975. Forest-management practices as related to nutrient leaching and water quality. Pp. 157-173. In: Conference on Non-point Sources of Water Pollution Proceedings. Virginia Water Resources Research Center; Virginia Polytechnical Institute and State University, Blacksburg, Virginia, May 1-2, 1975.
- Daniel, H.A.; Elwell, H.M.; Cox, M.B. 1943. Investigations in erosion control and the reclamation of eroded land at the Red Plains Conservation Experiment Station, Guthrie, OK, 1930-1940. USDA Soil Conservation Service Tech. Bull. 837 p.
- DeBano, L.F.; Neary D.G.; Ffolliott, P.F. 1998. *Fire Effects on Ecosystems*. John Wiley & Sons, New York, 333 p.
- Diaz-Chavez, R.; Neto, A.E.; Berndes, G.; Neary, D.G. 2011a. Chapter 4: Bioenergy-related water quality issues. Pp. 41-61. In: *The Bioenergy and Water Nexus*. United Nations Environment Program, the Oko-Institut eV. Institute for Applied Ecology, and International Energy Agency Bioenergy Task 43. 173 pp.
- Diaz-Chavez, R.; Berndes, G.; Neary, D.; Nieto, A.; Fall, M. 2011b. Water quality assessment of bioenergy production. *Biofuel. Bioprod. Bior.* 5: 445-463
- Dissmeyer G.E. (ed.) 2000. *Drinking water from forests and grasslands: a synthesis of the scientific literature*. USDA Forest Service, Gen.Tech.Rep. SRS-39, Southern Research Station, Asheville, North Carolina. 246 p.
- Douglass, J.E.; Swank, W.T. 1975. Effects of management practices on water quality and quantity: Coweeta Hydrologic Laboratory, North Carolina. Pp. 1-13. In: USDA Forest Service Gen. Tech. Rep. NE-13, Northeastern Forest Experiment Station, Upper Darby, Pennsylvania.
- Feller, M.C. 1981a. Catchment nutrient budgets and geological weathering in Eucalyptus regnans ecosystems in Victoria. *Aust. J. Ecol.* 6: 65-77.
- Feller, M.C. 1981b. Effects of clearcutting and slash burning on stream temperature in southwestern British Columbia. *Water Resour. Res.* 17: 863-867.
- Feller, M.C.; Kimmons, J.P. 1984. Effects of clearcutting and slash burning on streamwater chemistry and watershed nutrient budgets in southwestern British Columbia. *Water Resour. Res.* 20: 29-40.
- Foran B.; Poldy F. 2002). *Future Dilemmas: Options to 2050 for Australia's Population, Technology, Resources and Environment*. CSIRO Resource Futures Working Paper 02/01. CSIRO Sustainable Ecosystems, Canberra.
- Fox, T.R.; Allen, H.L.; Albaugh, T.J.; Rubilar, R.; Carlson, C.A. 2007. Tree nutrition and forest fertilization of pine plantations in the southern United States. *South. J. Appl. For.* 31: 5-11.
- Fredriksen, R.L.; Moore, D.G.; Norris, L.A. 1975. Impact of timber harvest, fertilization, and herbicide treatment on stream water quality in the Douglas-fir regions. p. 283-313. In: Bernier, B.; Winget, C.H. (eds.) *Forest Soils and Forest Land Management: Proceedings of the Fourth North American Forest Soils Conference*, Laval University, Quebec City, Canada.
- Gleick, P.H. (ed.) 1993. *Water in Crisis: A Guide to the World's Fresh Water Resources*. Oxford University Press, New York. 473 p.
- Glendening, G.E.; Pase, C.P.; Ingebo, P. 1961). Preliminary hydrologic effects of wildfire in chaparral. Pp. 12-15. In: *Proc. 5th Ann. Arizona Watershed Symp.*, September 21, 1961, Phoenix, Arizona, Tucson, University of Arizona.
- Goethe, L.; Soderberg, H.; Sjolander, E. 1993. Effects on water chemistry, benthic invertebrates and brown trout following forest fertilization in central Sweden. *Scand. J. For. Res.* 8: 81-90.
- Gomi T.; Sidle R.C.; Noguchi S.; Negishi J.N.; Abdul Rahim, N.; Sasaki, S. 2006. Sediment and wood accumulations in humid tropical headwater streams: effects of logging and riparian buffers. *Forest Ecol. Manage.* 224:166-175
- Gottfried, G.; Neary, D.; Ffolliott, P.; Koestner, K. 2012. Cascabel prescribed fire long-term watershed study: An opportunity to monitor climate change. *Revisiting Experimental Catchment Studies in Forest Hydrology*, Proceedings of a Workshop held during the XXV IUGG General Assembly in Melbourne, June-July 2011.
- Graynoth, E. 1979. Effects of logging on stream environments and faunas in Nelson. *New Zeal. J. Mar. Fresh.* 13: 79-109.
- Grayson, R.B.; Haydon, S.R.; Jayasuriya, M.D.A.; Finlayson, B.L. 1993. *Water quality in mountain ash*

- forests – separating the impacts of roads from those of logging. *J. Hydrol.* 150: 459-480.
- Grip, H. 1982. Water chemistry and runoff in forest streams at Kloten. UNGI Report No. 58, Uppsala University. Uppsala, Sweden. 144 p.
- Harr, R.D.; Fredriksen, R.L. 1988. Water quality after logging small watersheds within the Bull Run Watershed, Oregon. *J. Am. Water Resour. Assoc.* 24: 1103-1111.
- Harriman, R. 1978. Nutrient leaching from fertilized forest watersheds in Scotland. *J. Appl. Ecol.* 15: 933-942.
- Harris, D.D. 1977. Hydrologic changes after logging in two small Oregon coastal watersheds. U.S. Geological Survey Water Supply Paper 2037. U.S. Geological Survey, Washington, D.C. 31 p.
- Heede, B.H. 1987. Overland flow and sediment delivery five years after timber harvest in a mixed conifer forest. *J. Hydrol.* 91: 627-634.
- Helvey, J.D. 1980. Effects of a north-central Washington wildfire on runoff and sediment production. *Water Resour. Bull.* 16: 627-634.
- Helvey, J.D.; Kochenderfer, J.M.; Edwards, P.J. 1989. Effects of forest fertilization on selected ion concentrations in Central Appalachian streams. Pp. 278-282. In: Proceedings of the 7th Central Hardwood Conference. USDA Forest Service General Technical Report NC-132, North Central Forest and Range Experiment Station, St. Paul, Minnesota.
- Hetherington, E.D. 1976. Dennis Creek: A look at water quality following logging in the Okanagan Basin. Environment Canada, Canadian Forestry Service, p. 1-28.
- Hetherington, E.D. 1985. Streamflow nitrogen loss following fertilization in a southern Vancouver Island watershed. *Can. J. Forest Res.* 15: 34-41.
- Hewlett, J.D.; Doss, R. 1984. Forests, floods, and erosion: A watershed experiment in the Southeastern Piedmont. *Forest Sci.* 30: 424-434.
- Hewlett, J.D.; Fortson, J.C. 1982. Stream temperature under an inadequate buffer strip in the Southeast Piedmont. *Water Resour. Bull.* 18: 983-988.
- Hopmans, P.; Bren, L.J. 2007. Long-term changes in water quality and solute exports in headwater streams of intensely managed radiate pine and natural eucalypt forest catchments in south-eastern Australia. *Forest Ecol. Manage.* 253: 244-261.
- Hornbeck, J.W.; Martin, C.W.; Pierce, R.S.; Bormann, F.H.; Likens, G.E.; Eaton, J.S. 1987. The northern hardwood forest ecosystem: 10 years of recovery from clearcutting. USDA Forest Service Res. Pap. NE-596. Northeastern Forest Experiment Station, Upper Darby, Pennsylvania. 30 p.
- Ice, G. 2004. History of innovative Best Management Practice development and its role in addressing water quality limited waterbodies. *J. Environ. Eng.* 130: 684-689.
- Ice, G.G.; Stednick, J.D. 2004. A Century of Forest and Wildland Watershed Lessons. Society of American Foresters, Bethesda, MD, 292 pp.
- Iwatsubo, G.; Nagayama, Y. 1994. Effects of sewage water spraying on mineral cycling in a forest ecosystem. *Forest Ecol. Manage.* 68: 75-85.
- Krause, H.H. 1982. Nitrate formation and movement before and after clear-cutting of a monitored watershed in central New Brunswick, Canada. *Can. J. Forest Res.* 12: 922-930.
- Lane, P.N.J.; Sheridan, G.J. 2002. Impact of an unsealed forest road stream crossing: water quality and sediment sources. *Hydrol. Proc.* 16: 2559-2612.
- Lane, P.N.J.; Sheridan, G.J.; Noske, P.J. 2006. Changes in sediment loads and discharge from small mountain catchments following wildfire in southeastern Australia. *J. Hydrol.* 331: 495-510.
- Leeks, G.J.L.; Roberts, G. 1987. The effects of forestry on upland streams - with special reference to water quality and sediment transport. Pp. 9-24. In: Good, J.E.G., (ed.) Environmental Aspects of Plantation Forestry in Wales. Grange-over-Sands, NERC/ITE, 64-69. (ITE Symposium, 22).
- Leonard, J.H. 1977. Nitrogen run-off from a radiate pine forest fertilized with urea. *New Zeal. J. For. Sci.* 22: 64-80.
- Levno, A.; Rothacher, J. 1969. Increases in maximum stream temperatures after slash burning in a small experimental watershed. USDA Forest Service Res. Note PNW-110, Pacific Northwest Forest and Range Experiment Station, Portland, Oregon. 7 p.
- Likens, G.E.; Driscoll, C.T.; Buso, D.C. 1996. Long-term effects of acid rain: Response and recovery of a forest ecosystem. *Science* 272: 244-246.
- Likens, G.E.; Bormann, F.H.; Johnson, N.; Fisher, D.; Pierce, R. 1970. Effects of forest cutting and herbicide treatment on nutrient budgets in the Hubbard Brook watershed-ecosystem. *Ecol. Monogr.* 40: 23-47.

- Malmer, A. 1990. Stream suspended sediment load after clear-felling and different forestry treatments in tropical rainforest, Sabah, Malaysia. Pp. 62-71. In: Proc. Fiji Symp. IAHS-AISH Publ. 192. Research Needs and Application to Reduce Erosion and Sedimentation in Tropical Steeplands.
- Meehan, W.R.; Lotspeich, F.B.; Mueller, E.W. 1975. Effects of forest fertilization on two Southeast Alaska streams. *J. Environ. Qual.* 4: 50-55.
- Michael, J.L.; Neary, D.G. 1993. Herbicide dissipation studies in forest ecosystems. *Environ. Tox. Chem.* 12: 405-410.
- Miller, E.L.; Beasley, R.S.; Lawson, E.R. 1988. Forest harvest and site preparation effects on erosion and sedimentation in the Ouachita Mountains. *J. Environ. Qual.* 17: 219-225.
- Moody, J.A.; Martin, D.A. 2001. Initial hydrologic and geomorphic response following a wildfire in the Colorado Front Range. *Earth Surf. Processes* 26: 1049-1070.
- Moore, H.D.; Spittlehouse, D.L.; Story, A. 2005. Riparian microclimate and stream temperature response to forest harvesting: A review. *J. Amer. Water Resour. Assoc.* 41: 813-834.
- Napier, T.L. 2000. Soil and water conservation policies and programs: Successes and failures. CRC Press, Boca Raton, FL, 640 p.
- Neal, C.; Fisher, R.; Smith, C.J.; Hill, S.; Neal, M.; Conway, T.; Ryland, G.P.; Jeffrey, H.A. 1992. The effects of tree harvesting on stream-water quality at an acidic and acid-sensitive spruce forested area: Plynlimon, mid-Wales. *J. Hydrol.* 135: 305-319.
- Neary, D.G. 2000. Changing perceptions of watershed management from a retrospective viewpoint. Pp.167-176. In: Ffolliott, P.F.; Baker, M.B. Jr.; Edminster, C.B.; Dillon, M.C.; Mora, K.L. (tech. cords.) *Land Stewardship in the 21st Century: The Contributions of Watershed Management*. USDA Forest Service Proceedings RMRS-P-13, Rocky Mountain Research Station, Fort Collins, Colorado.
- Neary, D.G. 2002; Chapter 6: Environmental sustainability of forest energy production, 6.3 Hydrologic values. Pp. 36-67. In: Richardson, J.; Smith, T.; Hakkila, P. *Bioenergy from Sustainable Forestry: Guiding Principles and Practices*. Elsevier, Amsterdam. 344 p.
- Neary, D.G. 2011. Impacts of wildfire severity on hydraulic conductivity in forest, woodland, and grassland soils. Pp. 123-142. In: Elango, L. (ed.) *Hydraulic Conductivity - Issues, Determination and Applications*. INTECH, Rijeka, Croatia. ISBN 978-953-307-288-3, 434 p.
- Neary, D.G.; Michael, J.L. 1996. Herbicides—protecting long-term sustainability and water quality in forest ecosystems. *New Zeal. J. For. Sci.* 26: 241-264.
- Neary, D.G.; Bush, P.B.; Douglass, J.E. 1983. Offsite movement of hexazinone in stormflow and baseflow from forest watersheds. *Weed Sci.* 31: 543-551.
- Neary, D.G.; Bush, P.B.; Michael, J.L. 1993. Fate of pesticides in southern forests: a review of a decade of progress. *Environ. Toxicol. Chem.* 12: 411-428.
- Neary, D.G.; Ice, G.G.; Jackson, C.R. 2009. Linkages between forest soils and water quantity and quality. *Forest Ecol. Manag.* 258: 2269-2281.
- Neary, D.G.; Ryan, K.C.; DeBano, L.F. (eds.) 2005. *Fire Effects on Soil and Water*. USDA Forest Service, Gen. Tech. Rep. RMRS-GTR-42, Volume 4, Rocky Mountain Research Station, Fort Collins, Colorado. 250 p.
- Neary, D.G.; Bush, P.B.; Douglass, J.E.; Todd, R.L. 1985. Picloran movement in an Appalachian hardwood forest watershed. *J. Environ. Qual.* 14: 585-592.
- Neary, D.G.; Koestner, K.A.; Youberg, A.; Koestner, P.E. 2012. Post-fire rill and gully formation, Schultz Fire 2010, Arizona, USA. *Geoderma* 191: 97-104.
- Neary, D.G.; Smethurst, P.J.; Baillie, B.; Petrone, K.C. 2011. Water quality and biodiversity effects of harvesting Streamside Management Zone tree plantations. CSIRO Special Report, Canberra, Australia, 101 p.
- Neary, D.G.; Hayes, D.; Rustad, L.; Vose, J.; Gottfried, G.; Sebesteyn, S.; Johnson, S.; Swanson, F.; Adams, M. 2012. U.S. Forest Service Experimental Forests and Ranges Network: A Continental Research Platform for Catchment-Scale Research. Revisiting Experimental Catchment Studies in Forest Hydrology, Proceedings of a Workshop held during the XXV IUGG General Assembly in Melbourne, June-July 2011.
- Norris, L.A. 1970. Degradation of herbicides in the forest floor. pp. 397-412. In: Youngberg, C.T.; Davey, C.B. (eds.), *Tree Growth and Forest Soils*. Oregon State University Press, Corvallis, Oregon.
- Ohruji, K.; Mitchell, M.J. 1998. Stream water chemistry in Japanese forested watersheds and its variability on a small regional scale. *Water Resour. Res.* 34: 1553-

- 1561.
- O'Loughlin, C.L. 1994. The forest and water quality relationship. *New Zeal. For.* 39: 26-30.
- O'Loughlin, C.L.; Pearce, A.J. (1976) Influence of Cenozoic geology on mass movement and sediment yield response to forest removal. *Bull. Int. Assoc. Eng. Geol.* 14: 41-46.
- O'Loughlin, C.L.; Rowe, L.K.; Pearce, A.J. 1980. Sediment yield and water quality responses to clearfelling of evergreen mixed forests in western New Zealand. Pp. 285-292. In: *Proceedings of the Helsinki Symposium on the Influence of Man on the Hydrological Regime with Special Reference to Representative and Experimental Basins, PIAHS-AISH Publication No. 130.*
- Oyarzun, C.; Aracena, C.; Rutherford, P.; Godoy, R.; Deschrijver, A. 2007. Effects of land use conversion from native forests to exotic plantations on nitrogen and phosphorus retention in catchments of southern Chile. *Water Air Soil Pollut.* 179: 341-350.
- Pase, P.C.; Ingebo, P.A. 1965. Burned chaparral to grass: early effects on water and sediment yields from two granitic soil watersheds in Arizona. Pp. 8-11. In: *Proc. 9th Ann. Arizona Watershed Symp., Tempe, AZ.*
- Penman, H.L. 1963. Vegetation and hydrology. Technical Communication No. 53, Commonwealth Bureau of Soils, Commonwealth Agricultural Bureau, Bucks, England, 124 p.
- Perrin, C.J.; Shortreed, K.S.; Stockner, J.G. 1984. An integration of forest and lake fertilization: transport and transformations of fertilizer elements. *Can. J. Fish. Aquatic Sci.* 41: 253-262.
- Petticrew, E.L.; Owens, P.N.; Giles, T.R. 2006. Effects of wildfire on suspended and gravel-stored sediments. *Water Air Soil Poll.* 6: 647-656.
- Pierce, R.S.; Hornbeck, J.W.; Likens, G.E.; Bormann, F.H. 1970. Effect of elimination of vegetation on stream water quality and quantity. *J. Int. Assoc. Hydro. Sci.* 96: 311-328.
- Pigram, J.J. 2006. *Australia's Water Resources: From Use to Management.* CSIRO Publishing, Collingwood, Victoria, Australia, 240 pp.
- Pires, M. 2004. Watershed protection for a world city: The case of New York. *Land Use Policy* 2: 161-175.
- Plamondon, A.P.; Ruiz, R.A.; Morales, C.F.; Gonzalez, M.C. 1991. Influence of protection forest on soil and water conservation (Oxapampa, Peru). *Forest Ecol. Manage.* 38: 277-238.
- Pope, J.B.; Archer, J.C.; Johnson, P.R. 1946. Investigations in erosion control and reclamation of eroded sandy clay lands of Texas, Arkansas, and Louisiana at the Conservation Experiment Station, Tyler, Texas, 1931-1940. USDA Soil Conservation Service Tech. Bull. 916. Washington, DC, 76 p.
- Quinn, J.M.; Stroud, M.J. 2002. Water quality and sediment and nutrient export from New Zealand hill-land catchments of contrasting land use. *New Zeal. J. Mar. Fresh.* 36: 409-430.
- Rango, A. 1995. A look to the future in watershed management. Pp. 15- 22. In: Ward, T.J. (ed.) *Watershed Management: Planning for the 21st Century.* Proceedings of the Symposium, Watershed Management Committee, Water Resources Engineering Division, American Society of Civil Engineers, August 14-16, 1995, San Antonio, Texas. 442 p.
- Reimold, R.J. 1998. *Watershed Management: Practice, Policies, and Coordination.* McGraw-Hill Company, New York, 391 p.
- Riekerk, H.; Swindel, B.F.; Replogle, J.A. 1980. Effect of forestry practices in Florida watersheds. Pp. 706-720. In: *Proceedings of the Symposium on Watershed Management 1980, ASCE, Boise, ID, July 21-23, 1980.*
- Riggan, P.J.; Lockwood, R.N.; Jacks, P.M. 1994. Effects of fire severity on nitrate mobilization in watersheds subject to chronic atmospheric deposition. *Environ. Sci. Tech.* 28: 369-375.
- Ringrose, C.; Meyer, S.; Bren, L.J.; Neilsen, W.A. 2001. hydrology of small catchments in the WARRALTER site: objectives and preliminary analysis. *Tasforests* 13: 31-44.
- Rinne, J.N.; Janisch, J. 1995. Coldwater fish stocking and native fishes in Arizona: Past, present, and future. *Amer. Fish. Soc. Symp.* 15: 397-406.
- Rishel, G.B.; Lynch, J.A.; Corbett, E.S. 1982. Seasonal stream temperature changes following forest harvest. *J. Environ. Qual.* 11: 112-116.
- Rosen, K. 1984. Effect of clear-felling on runoff in two small watersheds in central Sweden. *Forest Ecol. Manage.* 9: 267-281.
- Rosen, K. 1996. Effect of clear-felling on streamwater quality in forest catchments in central Sweden. *Forest Ecol. Manage.* 83: 237-244.
- Rosen, M.R.; Lapham, W.W. 2008. Introduction to the U.S. Geological Survey national water-quality

- assessment (NAWQA) of ground-water quality trends and comparison to other national programs. *J. Environ. Qual.* 37: S-190–S-198.
- Ryan, D.F.; Glasser, S. 2000. Chapter 1: Goals of this report. Pp. 3-6. In: Dissmeyer, G.E. (ed.). *Drinking water from forests and grasslands: a synthesis of the scientific literature*. USDA Forest Service Gen. Tech. Rep. SRS-39. Southern Research Station, Asheville, North Carolina.
- Sahin, V.; Hall, M.J. 1996. The effects of afforestation and deforestation on water yields. *J. Hydro.* 178: 293-309.
- Scott, D.F. 1993. The hydrological effects of fire in South African mountain catchments. *J. Hydrol.* 150: 409-432.
- Scott, D.F.; Lesch, W. 1996. The effects of riparian clearing and clearfelling of an indigenous forest on streamflow, stormflow, and water quality. *S. Afr. For. J.* 175: 1-14.
- Shakesby, R.A.; Doerr, S.H. 2006. Wildfire as a hydrological and geomorphological agent. *Earth-Sci. Rev.* 74: 269-307.
- Sidle, R.C.; Sasaki, S.; Otsuki, M.; Noguchi, S.; Abdul Rahim, N. 2004. Sediment pathways in a tropical forest: effects of logging roads and skid trails. *Hydrol. Process.* 18: 703–720.
- Singh, T.; Kalra, Y.P. 1975. Changes in chemical composition of natural waters resulting from progressive clearcutting of forest catchments in West Central Alberta, Canada. *Int. Assoc. Sci. Hydro. Symp. Proc.*, Tokyo, Japan. December 1970. Publication 117. Pp. 435-449.
- Smith, C.T.; Dyck, W.J.; Beets, P.N.; Hodgkiss, P.D.; Lowe, A.T. 1994. Nutrition and productivity of *Pinus radiata* following harvest disturbance and fertilization of coastal sand dunes. *For. Ecol. Manage.* 66: 5-38.
- Smith, H.G.; Hopmans, P.; Sheridan, G.J.; Lane, P.N.J.; Noske, P.J. 2012. Impacts of wildfire and salvage harvesting on water quality and nutrient exports from *radiata* pine and eucalypt forest catchments in south-eastern Australia. *Forest Ecol. Manage.* 263: 160-169.
- Smith, H.G.; Sheridan, G.J.; Lane, P.N.J.; Nyman, P.; Haydon, S. 2011. Wildfire effects on water quality in forest catchments: A review with implications for water quality. *J. Hydrol.* 396: 170-192.
- Snyder, G.G.; Haupt, H.F.; Belt, G.H. Jr. 1975. Clearcutting and burning slash alter quality of stream water in Northern Idaho. USDA Forest Service Res. Pap. INT-1698, Intermountain Forest and Range Experiment Station, Ogden, Utah. 34 p.
- Stednick, J.D. 1996. Monitoring the effects of timber harvest on annual water yield. *J. Hydro.* 176: 79-95.
- Steen, H.K. 1976. *The U.S. Forest Service, A History*. University of Washington Press, Seattle, 356 p.
- Stottlemyer, R. 1992. Nutrient concentration patterns in streams draining alpine and subalpine catchments, Fraser Experimental Forest, Colorado. *J. Hydro.* 140: 179-208.
- Swank, W. T.; Douglass, J. E. 1975: Nutrient flux in undisturbed and manipulated forest ecosystems in the Southern Appalachian Mountains. *International Association of Scientific Hydrology. Symposium Proceedings, Tokyo, Japan. December 1970. Publication 117. Pp. 445-456.*
- Swank, W.T.; Vose, J.M.; Elliott, K.J. 2001. Long-term hydrologic and water quality responses following commercial clearcutting of mixed hardwoods on a southern Appalachian catchment. *Forest Ecol. Manage.* 143: 163-178.
- Swanson, F.J.; Scatena, F.N.; Dissmeyer, G.E.; Fenn, M.E.; Verry, E.S.; Lynch, J.A. 2000. Chapter 3: Watershed processes –fluxes of water, dissolved constituents, and sediment. Pp. 26-41. In Dissmeyer, G.E. (ed.). *Drinking water from forests and grasslands: a synthesis of the scientific literature*. USDA Forest Service Gen. Tech. Rep. SRS-39. Southern Research Station, Asheville, North Carolina.
- Swanson, R.H.; Golding, D.L.; Rothwell, R.L.; Bernier, P.Y. 1986. Hydrologic effects of clear-cutting at Marmot Creek and Streeter watersheds, Alberta. *Information Report NOR-X-278*, Northern Forestry Centre, Canadian Forestry Service, Edmonton, Alberta, Canada. 27 p.
- Swift, L.W. Jr. 1988. Chapter 23: Forest access roads: Design, maintenance, and soil loss. Pp. 313-324. In: Swank, W.T.; Crossley, D.A. (eds.) *Forest Hydrology and Ecology at Coweeta*. Springer-Verlag, New York, NY. 469 p.
- Swift, L.W. Jr.; Messer, J.B. 1971. Forest cuttings raise temperatures of small streams in the southern Appalachians. *J. Soil Water Conserv.* 26, 111-116.
- Tamm, C.O.; Holmen, H.; Popovic, B.; Wiklander, G. 1974. Leaching of plant nutrients from soils as a consequence of forestry operations. *Ambio.* 3: 211-221.
- Terrene Institute. 1993. *Proceedings of a technical workshop on sediments*. Terrene Institute, Washington, D.C. 143 p.

- Tiedemann, A.; Helvey, J.; Anderson, T.D. 1978. Stream chemistry and watershed nutrient economy following wildfire and fertilization in Eastern Washington. *J. Environ. Qual.* 7: 580-588.
- Trimble, S.W.; Crosson, P. 2000. U.S. soil erosion rates – myth and reality. *Science* 289: 248-250.
- Ursic, S.J. 1970. Hydrologic effects of prescribed burning, and deadening upland hardwoods in northern Mississippi. USDA Forest Service Res. Pap. SO-54. Southern Forest Experiment Station, New Orleans, Louisiana 15 p.
- Vaidya, O.C.; Smith, T.P.; Fernand, H.; McInnis-Leek, N.R. 2008. Forestry Best Management Practices: Evaluation of alternate streamside management zones on stream water quality in Pockwock Lake and Five Mile Lake watersheds in central Nova Scotia, Canada. *Environ. Monit. Assess.* 137: 1-14.
- Van Lear, D.H.; Douglass, J.E.; Fox, S.K.; Augsberger, M.K. 1985. Sediments and nutrient export in runoff from burned and harvested pine watersheds in the South Carolina Piedmont. *J. Environ. Qual.* 14: 169-174.
- Verry, E.S. 1972. Effect of aspen clearcutting on water yield and quality in northern Minnesota. Pp. 276-284. In: Csallany, S.C.; McLaughlin, T.G.; Striffler, W.D. (eds.) *Watersheds in Transition. Proceedings of a Symposium, Fort Collins, CO, June 19-22, 1972, American Water Resources Association and Colorado State University.* 405 p.
- Vertessy, R.A. 1999. The impacts of forestry on streamflows: a review. Pp. 93-109. In: Croke, J.; Lane, P. (eds.) *Forest Management for the Protection of Water Quality and Quantity, Proceedings of the Second Erosion in Forests Meeting, Warburton, 4-6 May 1999, Cooperative Research Centre for Catchment Hydrology, Report 99/6.*
- Vose, J.M.; Sun, G.; Ford, C.R.; Bredemeier, M.; Otsuki, K.; Wei, X.; Zhang, L. 2011. Forest ecohydrological research in the 21st century: what are the critical needs. *Ecohydrology* 4: 146-158.
- Wells, W.G., II. 1981. Some effects of brushfires on erosion processes in coastal southern California. Pp. 305-342. In: Davies, T.R.R. (ed.) *Proceedings, erosion and sediment transport in Pacific rim steeplands; January 25–31, 1981; Christchurch, New Zealand.; Publ. No.132. International Association of Scientific Hydrology, Washington, DC.*
- Westling, O.; Hultberg, H. 1990. Liming and fertilization of acid forest soil: short-term effects on runoff from small catchments. *Water Air Soil Poll.* 54: 391-407.
- World Health Organization 2006. *Guidelines for drinking water quality. Volume:1 Recommendations.* World Health Organization, Geneva, Switzerland, 595 p.
- Wright, H.A.; Churchill, F.M.; Stevens, W.C. 1976. Effect of prescribed burning on sediment, water yield, and water quality from juniper lands in central Texas. *J. Range Manage.* 29: 294-298.