

Correlation of Environmental Factors with Nitrate Concentration in Meadow Plants

F.B. GOMM

Abstract

Earlier studies of meadow forages in eastern Oregon have shown that fertilizer-N increased yields and N concentration. How much of the increased N was $\text{NO}_3\text{-N}$ or how variations in environmental factors affected the accumulation of NO_3 in meadow plants is not known. Plots of native meadow were fertilized on March 31 with ammonium sulfate to provide nitrogen levels of 0, 110, 220, 440 kg/ha. Changes in air temperature; soil temperature at the 0-, 5-, 10-, 20-, and 50-cm depths; luminous flux at canopy level, at 20 cm above ground, and at ground level; solar radiation over vegetation; and soil moisture content were monitored through the growing season. These data were statistically correlated with concentrations of $\text{NO}_3\text{-N}$ extracted from plant tissue. Herbage yields at the end of the growing season increased from 4,275 kg/ha without fertilizer to 9,782 kg/ha with 440 kg N/ha. Total-N concentration was highest (3.33%) on May 5 in herbage fertilized with 440 kg N/ha. As plants matured, concentration of total-N decreased by August 29 to 1.15% in plants receiving the 440 kg/ha rate and to 0.72% in unfertilized plants and was higher in the leaves than in the stems. Conversely, concentration of $\text{NO}_3\text{-N}$ were lowest (10-70 ppm) early in the season. Late in the growing season the $\text{NO}_3\text{-N}$ level was lowest in the leaves and highest (1,060 ppm) in the stems. Correlation analyses indicated that soil-moisture content was the most significant environmental factor correlated with accumulations of $\text{NO}_3\text{-N}$, except for soil fertility as measured by increased fertilization rate. Luminous flux at ground level and air temperature were also contributing factors.

Native plants commonly found on flood meadows of eastern Oregon are tolerant to long periods of flooding. These plants thrive under seasonally moist to saturated soil conditions, yet survive periodic droughts. With nitrogen (N) fertilization, these meadow plants generally have increased yields and increased N content of the herbage (Rumburg

1969 and 1972). In these meadows, the level of herbage-N increased rapidly after fertilizer was applied, then decreased gradually until the N concentration of the fertilized herbage resembled that of nonfertilized plants (Rumburg 1972). How much of the N content in the herbage was NO_3 or how long the NO_3 persisted in the plants as they matured was not studied. In similar fertilizer studies, but with different grass species, Lewis (1960), Slinkard (1964), and Willhite et al. (1955) also observed increased yields and high total-N concentrations.

Although few cattle deaths in Oregon have been attributed to NO_3 poisoning, cattle using high-nitrogen forage in other areas have died. Nitrate poisoning is not limited to cattle grazing plants that have been fertilized with N. It also occurs when cattle graze plants after environmental stress. Primary factors known to affect plant accumulation of NO_3 include nutrient supply, moisture stress, light, soil type, herbicides, and disease (McKee 1962).

The primary objective of this study was to determine the relationship of temperature, light, soil moisture, and fertilizer to NO_3 accumulation in native meadow plants.

Materials and Methods

Plots (3×6 m) arranged in a 4×4 Latin square, were located on a native flood meadow of the Squaw Butte Experiment Station, near Burns, Oregon. The vegetation was a mixture of grasses and sedges, dominantly beardless wildrye (*Elymus triticoides* Buckl.) with slender sedge (*Carex praegracilis* W. Boott) and Nevada bluegrass (*Poa nevadensis* Vasey ex Scribn.). Ammonium sulfate was applied March 31 to provide nitrogen levels of 0, 110, 220, and 440 kg/ha. The soil was flooded with standing water from May 1 until June 26 and then allowed to dry. Rainfall from March 31 to May 1 was less than 0.5 cm, and during the drying period it was less than 1.25 cm.

Ten grab samples of herbage was taken at 3-week intervals from May 5 until August 29 by clipping herbage to ground level. After

Author is range scientist, U.S. Dep. Agr.-SEA-Agr. Res., Crops Research Lab., Utah State University, Logan, 84322. Formerly at Oregon State University, Squaw Butte Agricultural Experiment Station, Burns, Ore.

This study represents the cooperative investigation of the U.S. Dep. Agr.-SEA-Agr. Res. and the Agr. Exp. Sta., Oregon State Univ. Tech. Pap. No. 4915 of the latter. Manuscript received January 8, 1979.

June 2, samples were collected separately from upper leaves, mid-stem, and lower stem of the plant. The final yield was taken on August 29 by harvesting 1-m² sample from each treatment plot. Each herbage sample was dried at 65°C for 48 hours, ground, and analyzed for total-N, NO₃-N, and NH₄-N by Kjeldahl and micro-Kjeldahl methods (Bremner and Keeney 1966, as modified for plants by Oregon State University Soil Testing Laboratory). Phenological development and plant height were recorded periodically.

Soil samples were taken on May 5 and October 6 from the surface 30 cm of soil. The soil was dried, ground, and analyzed for NO₃-N and NH₄-N by the micro-Kjeldahl method for soil (Bremner and Keeney 1966).

The following environmental factors were monitored: air temperatures, daily maxima and minima at the standard 1-m height; luminous flux at canopy level, at 20 cm above ground, and at ground level was recorded at 1-hour intervals on designated days. Luminous flux from blue sky may be converted to photosynthetic irradiance in the 400-700 nm wave band where W/m² klux 4.88 (McCree 1972). Luminous flux was measured with a Kahlsico miniphotometer (35AM200)¹. Soil temperatures were recorded daily at the 10-cm depth using an Instrument Corporation recording thermograph. On selected dates, soil temperature was recorded at 2-hour intervals at 0-, 5-, 10-, 20-, and 50-cm depths on fertilized and unfertilized plots. These temperature measurements were made using thermocouples constructed from Honeywell constantan-copper wire (Model 1W2P6), as indicated on a Honeywell potentiometer (Model 2736). Solar radiation over vegetation was determined with a constant recording pyrheliograph (Belfort Instrument Company, No. 53850). Net radiation was determined with a portable net radiation indicator (C.W. Thornwaite Associated, Model 603). Air temperatures were recorded on a constant recording hygrothermograph (Belfort Instrument Company, No. 5-594). Soil moisture content was evaluated by drying soil samples at 105° C for 48 hours and determining moisture content gravimetrically.

Results and Discussion

Environmental Influences

Soil Moisture

From May 1 to June 26 the soil remained saturated. The flooding depth was never greater than 2.5 cm and was uniform across the study plots. After July 1, the soil gradually dried until the end of the experiment (Table 1).

Soil N

Soil analyses indicated that in early May the residual-N was in the NH₄ form. At that time available-N in the saturated soil was less than 1 ppm NO₃. The NH₄-N concentra-

tion, however, was 13.7 ppm in unfertilized plots and increased in fertilized plots. As the soil dried, the NO₃-N in unfertilized plots increased to 13.9 ppm, and NH₄-N decreased to 2.8 ppm. These changes in N-form agreed with the findings of Van Schreven and Sieben (1972). By October 6, the differences in soil N concentrations were minimal among fertilizer rates.

Radiation.

Solar radiation at the field site range from 1.0 to 1.5 ly/min (Table 1). In May and June 600 to 750 ly/day were received. In July, August, and September solar radiation decreased to 625, 575, and 480 ly/day, respectively. The average net radiation received at the plants from June 2 to August 17 was about 50% of the solar radiation.

Irradiance

Light measurements on June 22 show that about 107,500 lux (525 W/m²) was received at canopy level for about 45 minutes. Within the cover of the unfertilized vegetation, the irradiance level was above 32,000 lux (156 W/m²) for 2 hours, whereas at ground level it was above 10,750 lux (52.5 W/m²) for about 3.3 hours. Within the cover of grass fertilized with 440 kg N/ha, irradiance levels were 8,500-107,750 lux (41.5-52.5 W/m²) for 2.5 hours. At the soil surface the irradiance level was above 1,075 lux (5.2 W/m²) for only 4 hours. For other dates irradiance levels at the soil surface were shown in Figure 1.

The amount of light that reached the lower plant parts was greater earlier than later in the growing season. It was also much greater within unfertilized vegetation than within the fertilized. As the season progressed and the herbage became dense, less light penetrated the canopy (Table 1). By the end of August, the luminous flux level received at the lower plant stem of fertilized vegetation was reduced to 108 lux (0.5 W/m²).

Soil Temperature

Early in the season, soil temperatures were similar under the meadow vegetation of the unfertilized and fertilized plots. By May 31, however, the temperature at the 10-cm soil depth was higher under unfertilized vegetation than it was under fertilized vegetation. From June 14 to September 10, the soil at the 50-cm depth was 1° C warmer under the unfertilized vegetation than it was under the fertilized, whereas the soil surface the temperature was 4 to 5° C warmer under unfertilized vegetation.

From June 22 to August 19, the average temperature of the surface soil in unfertilized plots was 16° C at 0700 hours;

¹Mention of a product does not constitute a guarantee or warranty by the U.S. Dep. Agr.

Table 1. Environmental data from fertilized meadow on various sampling dates.

Date (1972)	Days from May 1 No.	Maximum air temperature ° C	Maximum surface soil temperature ° C	Maximum soil temperature at 10-cm depth ° C	Maximum irradiance at soil surface ¹ lux	Solar radiation ly/min	Soil moisture content %
May 5	5	17	24	15	3762	1.3	54
May 24	24	17	24	15	4300	1.3	55
June 2	33	26	30	20	5375	1.4	54
June 22	53	26	30	18	3225	1.5	59
July 7	68	28	28	18	430	1.5	46
July 24	85	28	31	18	322	1.3	38
August 17	109	20	25	16	161	1.2	28
August 29	121	31	26	18	108	1.0	24

¹Irradiance was measured with a photometer in luminous flux.




Fig. 1. Irradiance at the soil surface under meadow vegetation, unfertilized (a) and fertilized (b). Note scale differences between (a) and (b).

it then rose to a high of 32° C at 1300 hours. Under vegetation fertilized with 440 kg N/ha, the highest temperature was 25° C.

At the 5-cm depth, the temperature was warmed to 25° C by May 31 under unfertilized vegetation. This was 3° C warmer than it was under fertilized vegetation. By August 9, the maximum temperature under unfertilized vegetation was 28° C, and the difference in soil temperature between the two treatments widened to 6° C then narrowed again, until on September 10 the soil from the unfertilized treatment was 1° C warmer than it was under fertilized vegetation.

At the 20-cm depth the soil reached its maximum temperature of 18° C in mid-August at 0900 hours under unfertilized vegetation. The temperature of the soil was 2 to 3° C warmer under the unfertilized treatment than under the fertilized vegetation.

Although soil temperature change lagged in relation to time and depth in sine-wave patterns, the variation in temperature below 20 cm was fairly constant with a daily range of 3 to 5° C.

Phenological Development

Meadow plants began to grow about the first of April, but little difference among fertilizer treatments was observed until May 10. By May 20 plants receiving 220 and 440 kg N/ha were taller than those receiving the 110 kg/ha rate or




Fig. 2. Height and development of beardless wildrye as affected by nitrogen fertilization.

less (Fig. 2). The differences among treatments continued to widen until June 22, when most plants approached their maximum height. Some plants receiving 220 and 440 kg N/ha developed seed heads, but those receiving 0 and 110 kg N/ha developed only vegetative stems. Yields on August 29 increased as rates of fertilizer-N increased (Table 2).

Table 2. Yield and nitrogen recovered from meadow herbage of plants harvested August 29 as affected by fertilizer rate.

Fertilizer rate of N (kg/ha)	Herbage ¹ yield kg/ha	Concentration of N in herbage %	Nitrogen in herbage k/gha	Efficiency of fertilizer %
0	4198	0.80a ²	33a	—
110	6969	0.77a	53.7b	19
220	8046	0.72a	57.9b	11
440	9607	1.15b	110.5c	18

¹LSD_{0.01} = 150 kg/ha.

²Values in columns followed by different letters are significantly different at $P < 0.05$ according to Duncan's multiple range test.

The low irradiance level within the fertilized vegetation late in the season caused the shaded leaves to yellow as in senescence. Salisbury and Ross (1969) suggested that senescence is the result of altered hormone balance that prevents the production of enzymes synthesized when the cells were younger. It is, therefore, reasonable to believe that the irradiance level at the lower parts of these plants may have been too low to continue the life processes of the lower leaves, and yellowing resulted from light deficiency, not maturation.

Chemical Analyses

Ammonium Concentration

The normal $\text{NH}_4\text{-N}$ level in meadow plant tissue, predominantly beardless wildrye, appears to be 50 to 80 ppm (Table 3). Unfertilized plants remained within this concentration range throughout the season. Fertilized plants seemed to contain their highest $\text{NH}_4\text{-N}$ levels early in the season with concentrations decreasing as the season advanced. This consistent decrease was observed particularly in herbage from plants which received fertilizer-N at the 440 and 220 kg N/ha rates.

Table 3. Ammonium and nitrate concentration (ppm) in meadow herbage as affected by fertilizer-N levels and seasonal growth¹.

Sampling date	Concentrations at four rates of N (kg/ha)			
	0	110	220	440
Ammonium-nitrogen				
May 5	50	170	130	170
May 24	80	90	110	160
June 2	60	30	180	180
June 22	50	50	90	130
July 7	70	20	60	100
July 24	60	60	60	80
Aug. 17	60	40	50	80
Aug. 29	70	90	40	70
Nitrate-nitrogen				
May 5	10	40	10	70
May 24	30	20	10	80
June 2	10	10	80	200
June 22	10	10	10	300
July 7	10	10	10	340
July 24	10	10	10	340
Aug. 10	70	10	30	440
Aug. 29	20	70	10	580

¹Beardless wildrye was the predominant species.

Total-N Concentration

The concentration of total-N early in the season was highest in herbage fertilized with 440 kg N/ha (Fig. 3). As plants matured, total-N levels decreased. After June 1, total-N concentrations were similar at rates of N less than 220 kg/ha. Although statistical analyses indicated no differences in total-N from fertilizer treatments on July 24 and August 17, mean concentrations were consistently highest in herbage treatments receiving 440 kg N/ha. From May 24 to July 7, the differences in total-N concentrations among fertilizer treatments were highly significant ($P < 0.01$). This result was primarily due to the high concentration of N in plants fertilized with 440 kg N/ha.




Fig. 3. Concentration of total-N in meadow forage at various dates as affected by fertilizer rate.

Total-N concentrations were highest in leaf tissue in early May with no difference among fertilizer treatments. In early June, the total-N in the stem tissue was relatively high but not as high as it was in the leaves and the concentration was highest with 440 kg N/ha (Fig. 4). As the season advanced, total-N concentrations in both leaves and stems decreased so that near the end of the season and the unfertilized stems contained about 0.6% total-N, and those fertilized with 440 kg N/ha contained 0.9% total-N. At the same times leaves




Fig. 4. Concentration of total-N in tissue of meadow forage as affected by fertilizer rate.

contained 1.2 and 1.5% total-N at the 0- and 440-kg N/ha rates, respectively. Less than 20% of the N applied was used in herbage production (Table 2).

Nitrate N

The normal level of the $\text{NO}_3\text{-N}$ in the meadow herbage, predominantly beardless wildrye, appears to be less than 10 ppm (Table 3). Except for occasional increases, only traces of $\text{NO}_3\text{-N}$ were measured from samples receiving 220 kg N/ha or less. Nevertheless, the 440 kg N/ha rate caused the $\text{NO}_3\text{-N}$ level to increase. During the flooding period, while the soil was saturated, the $\text{NO}_3\text{-N}$ concentration in herbage from the 440 kg N/ha treatment was significantly lower than later when the soil was dry. The highest $\text{NO}_3\text{-N}$ concentration in the herbage occurred during August when the soil was driest (Table 1).

The production and activation of nitrate reductase is a light-stimulated reaction (Hageman and Flesher 1960). Therefore, the NO_3 concentration in the stems and lower, shaded leaves should be higher than that in leaves exposed to full sunlight. Early in June, 80 and 220 ppm $\text{NO}_3\text{-N}$ in the lower stems was detected from the 220 and 440 kg N/ha rates, respectively. During late June under the 440 kg N/ha treatment, the $\text{NO}_3\text{-N}$ level increased to 240 ppm in the upper leaves, 990 ppm in the midstem, and 1,060 ppm in the lower stem. During July and August, the $\text{NO}_3\text{-N}$ level remained higher in the lower stem than in upper leaves, but concentrations were never high enough to be toxic (2,100 ppm $\text{NO}_3\text{-N}$) to livestock.

Simple correlation analyses were used to test the assumption that the $\text{NO}_3\text{-N}$ increase late in the season was related to decreasing soil moisture and increasing shade at ground level—both of which were associated with advance in season. Only data for $\text{NO}_3\text{-N}$ concentration in herbage from plots fertilized with N at 440 kg/ha were used.

Days from May 1 accounted for 94% of the variation (Fig. 5). Decreasing soil moisture content was the most significant factor correlated with $\text{NO}_3\text{-N}$ accumulation (Fig. 6a). Luminous flux at ground level (Fig. 6b) and maximum air temperature (Fig. 6c) were also weakly correlated with the $\text{NO}_3\text{-N}$ accumulation ($P < 0.05$). These three factors, soil moisture, light, and air temperature, are interrelated with stage of growth (days from May 1), seasonal influences, and characteristics of plant growth that tend to be related with time to the NO_3 concentration. Multiple correlation analysis of these factors compared closely with the simple correlation of analysis of days from May 1 (Fig. 5). Multiple correlation accounted for about 97% of the variation. Solar radiation at




Fig. 5. The correlation of time (seasonal development) to the accumulation of NO_3 in meadow forage as expressed by simple and multiple correlation analyses. In the multiple correlation analysis, air temperature (maxima at 1300 hr), light intensity (maxima luminous flux), and soil moisture content on sampling dates were used to derive computed multiple correlation values.

canopy level (Fig. 6e) and the soil temperature at the soil surface (Fig. 6f) and at the 10-cm depth (Fig. 6d) were nonsignificant factors in simple correlation analyses.

Considering the simple correlation coefficients of NO_3 -N concentration with luminous flux ($r_{1.2} = -0.818$) and with soil moisture stress ($r_{2.1} = 0.885$), both variables seemed to be strongly related to NO_3 -N accumulation in the plants, and the corresponding coefficient of multiple correlation ($R = 0.89$) suggests that 80% of the deviation from the mean was explained by the influence of these two variables. A comparison of the standard partial regression coefficients for NO_3 -N concentration by luminous flux ($b_{1.2} = -0.010$) and by soil-moisture stress ($b_{2.1} = 0.793$) showed that the YX_2 relationship was strong whereas the YX_1 relationship was less strong. The F test ($P = 0.05$) indicated with good confidence that most of the deviation was due to the influence of soil-moisture stress and luminous flux of the light source.

The overriding effect of soil moisture on NO_3 concentration is well demonstrated. The normal response is for the NO_3 level to decrease as plants mature. However, plants common to the meadows of eastern Oregon contained relatively low NO_3 levels in the early summer, when the soil was saturated, as compared with the increased NO_3 levels when the soil dried. To high correlation of NO_3 level with soil moisture content substantiated this observation (Fig. 6a). Several authors have reported studies which help explain




Fig. 6. Nitrate concentration in meadow forage as correlated with environmental variables determined at 1300 hr on date of sampling.

these results. Spratt and Gasser (1970), Plaut (1973), Terman et al. (1969), and Tisdale and Nelson (1966) have shown the importance of adequate soil moisture for efficient use of N by plants. Soil moisture influences the N available to plants as it affects the behavior of soil microorganisms. The activity of nitrifying organisms is inhibited at extremely low or extremely high moisture levels, which may reduce the supply of available N. When soil becomes waterlogged, as it was during the flooding period of this experiment, oxygen is excluded and anaerobic decomposition occurs. The exclusion of oxygen and the activity of the anaerobic bacteria induced denitrification. Under the waterlogging conditions, NO_3 could have been reduced and the N lost to the atmosphere. Also, because NO_3 is extremely soluble in water, much of it could have been lost through leaching processes. As the soil moisture level decreased and oxygen became more available, ammonification of organic N could have taken place and the nitrifying processes increased the supply of NO_3 in the soil, thus increasing the NO_3 concentration in the plant material. Soil moisture not only controls the behavior of the N in the soil, but also affects N uptake and utilization by the plant. D'Aoust and Tayler (1968) and Holliday (1970) showed that the N uptake is inefficient when the root zone is depleted of water. Also, Garcia and Gomez (1972) observed that the variation in NO_3 concentration depends on the water content of the herbage. They reported that the NO_3 level in alfalfa plants increased as the water content decreased, and Gomm (1979) observed that meadow plants, grown under controlled environmental conditions, could accumulate potentially toxic levels of NO_3 when

soil moisture conditions were less than saturated. Plants grown in saturated soil contained levels of NO₃ considered to be safe for cattle, even when plants were fertilized with 440 kg N/ha.

Conclusion

The rate of applied fertilizer-N was the most important of the factors considered that determined the level of NO₃ accumulation in native meadow plants. At normal application rates (90-130 kg N/ha), however, it is unlikely that NO₃ concentrations would accumulate to toxic levels. The primary reason that nitrate poisoning has not been associated with forage from fertilized meadows of eastern Oregon seems to be because of the saturated soil conditions during plant growth. Other environmental factors of secondary importance responsible for maintaining low levels of NO₃ in the forage seems to be the high level of irradiance and the relatively low air temperatures characteristic of eastern Oregon.

Literature Cited

- Bremner, J.M., and D.R. Keeney. 1966.** Determination of isotope-ratio analysis of different forms of nitrogen in soils:3. Exchangeable ammonium nitrate and nitrite by extraction-distillation methods. *Soil Sci. Soc. Amer. Proc.* 30: 577-582.
- D'Aoust, M.J., and R.S. Tayler. 1968.** The interaction between nitrogen and water in the growth of grass swards. 1. Methods and dry matter results. *J. Agr. Sci.* 70: 11-17.
- Garcia, B., and J.M. Gomez. 1972.** The effect of various factors on the NO₃ content of lucerne (*Medicago sativa* L.). *Agrochimica* 16: 409-417. (Herb Abstr. 43: 1187).
- Gomm, F.B. 1979.** Herbage yield and nitrate concentration in meadow plants as affected by environmental variables. *J. Range Manage.* 32: 359-364.

- Hageman, R.H., and D. Fleisher. 1960.** Nitrate reductase activity in corn seedlings as affected by light and nitrate content of nutrient media. *Plant Physiol.* 35: 700-708.
- Holliday, R. 1970.** Soil profile moisture and nitrogen availability. *In:* Kirkby, E.A. (ed.) *Nitrogen Nutrition of the Plant.* A Wilely and Sons. Leeds.
- Lewis, R.D. 1960.** Mountain meadow problems—with suggestions for improvement. Eleventh Fertilizer Conf. of the Pacific Northwest Proc. 59-64. Mimeo.
- McCree, K.J. 1972.** Test of current definitions of photosynthetically active radiation against leaf photosynthesis data. *Agr. Meteorol.* 10: 443-453.
- McKee, H.S. 1962.** *Nitrogen Metabolism in Plants.* Oxford Univ. Press. London and New York. 728 p.
- Plaut, Z. 1973.** Effect of soil moisture tension and nitrogen supply on nitrate reduction and accumulation in wheat seedlings. *Plant Soil* 38: 81-94.
- Rumburg, C.B. 1969.** Yield and N concentration of meadow hay fertilized with three N sources. *Agron. J.* 61: 824-825.
- Rumburg, C.B. 1972.** Yield and N accumulation of meadow forage fertilized at advancing maturity with N. *Agron. J.* 64: 187-189.
- Salisbury, F.B., and C. Ross. 1969.** *Plant Physiology.* Wadsworth Publishing Co., Inc. 761 p.
- Slinkard, A.E. 1964.** Mountain meadow fertilization. *Idaho Agr. Exp. Sta. Prog. Rep.* 90. 7 p. Mimeo.
- Spratt, E.D., and J.K.R. Gasser. 1970.** Effect of fertilizer-nitrogen and water supply on distribution of dry matter and nitrogen between the different parts of wheat. *Can. J. Plant Sci.* 50: 613-625.
- Terman, G.L., R.E. Ramig, A.F. Dreier, and R.A. Olson. 1969.** Yield-protein relationships in wheat grain as affected by nitrogen and water. *Agron. J.* 61: 755-759.
- Tisdale, S.L., and W.L. Nelson. 1966.** *Soil Fertility and Fertilizers.* 2nd ed. The MacMillan Co., New York. 694 p.
- Van Schreven, D.A., and W.H. Sieben. 1972.** The effect of storage on soils under waterlogged conditions upon subsequent mineralization of nitrogen, nitrification, and fixation of ammonia. *Plant Soil* 37: 245-253.
- Willhite, F.M., H.K. Rouse, and D.E. Miller. 1955.** High altitude meadows in Colorado. 3. The effect of nitrogen fertilization on crude protein. *Agron. J.* 47: 117-121.

THINK BAUSCH & LOMB... FOR GRAPHICAL DATA TRANSFER WORK


With a Bausch & Lomb ZOOM TRANSFER SCOPE* instrument, you can quickly update a data base or prepare a special purpose thematic map without expensive data transfer, projection or darkroom equipment. These instruments work on the camera lucida principle that optically mix aerial photographs and data base without projection. Compare these performance features with what you are now using.

- Built-in scale and stretch correction systems compensate for photographic distortions.
- Compact, lightweight design offers total portability. Can be used in a lighted office on any flat surface.
- Zoom control instantly matches photograph to data base scale.
- Accommodates all photographic formats including transparencies and prints (black and white or color). Stereo model (illustrated) helps you examine topographical and structural features.
- Two models feature a photographic mode for permanent documentation.

CALL US AT 716 338-6000

And let a Bausch & Lomb representative recommend the instrument that best meets your needs. Or, send coupon for illustrated literature.

*ZOOM TRANSFER SCOPE and Z.T.S. are trademarks of Bausch & Lomb Incorporated for graphical data transfer instruments.



BAUSCH & LOMB

Scientific Optical Products Division

ROCHESTER, NEW YORK 14602 USA
716-338-6000, TWX 510-253-6189
TELEX 97-8231. CABLE: BAUSCH & LOMB

In Canada: Bausch & Lomb Canada, Ltd. Scientific Optical Products Division, 2001 Leslie Street, Don Mills, Ontario, Canada M3B 2M5

Name	Organization	Address	City	State/Province	Zip
<input type="checkbox"/> Please send brochure describing the ZOOM TRANSFER SCOPE instruments (33-5016). <input type="checkbox"/> Please send literature on other B&L photointerpretation instruments (53-2290) (53-2288). <input type="checkbox"/> Please arrange a demonstration.					

Bausch & Lomb Incorporated
Scientific Optical Products
Division
Department 6606
1400 North Goodman Street
Rochester, New York 14602