

Near Infrared Spectroscopy and Fecal Chemistry as Predictors of the Diet Composition of White-Tailed Deer

Pierre-Olivier Jean,¹ Robert L. Bradley,² Marie-Andrée Giroux,³ Jean-Pierre Tremblay,⁴
and Steeve D. Côté⁵

Authors are ¹PhD student and ²Professor, Dép. de biologie, Université de Sherbrooke, QC, J1K2R1, Canada; ³PhD student and ^{4,5}Professors, Dép. de biologie, Université Laval, Québec, QC, G1V0A6, Canada.

Abstract

Overbrowsing by white-tailed deer (*Odocoileus virginianus* Zimmermann) on Anticosti Island (Canada) created a need to develop efficient methods for estimating their foraging patterns. We tested the ability of near infrared (NIR) spectra of feces and of fecal chemical properties to predict diet composition of different individuals. We first used a principal component-based discriminant analysis to sort the NIR spectra of fecal samples ($n=102$) obtained from two groups of captive deer that had been fed two different diets. The diets differed only in their relative abundance of balsam fir (*Abies balsamea* [L.] P.Mill.) and white spruce (*Picea glauca* [Moench] Voss.) foliage. The calibrated model allowed us to assign 28 of 30 validation fecal samples (93.3%) to the correct diet. In a second study, we attempted to estimate the proportion of coniferous, deciduous, herbaceous, and lichenous forages in diets of free-ranging white-tailed deer, as determined by fecal microhistology. Both NIR spectra and chemical properties of feces were used as predictors of diet composition. NIR spectra were analyzed using partial least-squares regression (PLSR), whereas fecal chemical properties were analyzed using mixed-linear regressions (MLRs). The PLSR models were robust ($R^2=0.89$; ratio of prediction to deviation=3.2) for predicting the amount of coniferous fragments, but not for predicting the relative amounts of balsam fir, white spruce, and deciduous and lichenous fragments within feces. MLR models revealed a positive relationship (47% variance explained) between acid detergent lignin (ADL) and coniferous fragments within feces. ADL and cellulose explained 24% of variance in deciduous fecal fragments, whereas ADL alone explained 22% of variance in balsam fir fecal fragments. These results suggest that NIR spectroscopy and fecal chemical properties have several applications on Anticosti Island, such as measuring the degree of variation in diets within a given home range or determining dietary conifer intake during winter.

Key Words: diet quality, fecal analysis, fecal microhistology, near infrared spectroscopy, overbrowsing, white-tailed deer

INTRODUCTION

Studying the foraging behavior of large wild herbivores in heterogeneous landscapes informs us on habitat selection, seasonal movements, or the impacts of herbivory on plant communities. To this end, a variety of methods have been developed over past decades to describe the diet of wild herbivores. These include fecal analyses (Stewart 1967), the probing of stomach contents (McInnis et al. 1983), browse surveys (Shiple et al. 1998), behavioral observations in the field (Parker et al. 1999), as well as isotope fractionation of animal tissues (Dalerum and Angerbjörn 2005). Among these techniques, fecal analyses are an interesting option because fecal samples are easy to obtain in high-density populations and do not require the handling or killing of animals. The main disadvantage of this approach is that highly digestible plant species could be underrepresented in the estimated diets, whereas less digestible plants could be overrepresented.

Nevertheless, Hanley and McKendrick (1985) showed 65% to 77% similarity between the estimated plant species composition of rumen contents and fecal samples of Sitka black-tailed deer (*Odocoileus hemionus sitkensis* Rafinesque).

Estimating the diets of large herbivores through fecal analyses usually entails the microhistological analysis of plant fragments (Dearden et al. 1975; Lefort 2002). This method is, however, expensive and time consuming, and it requires a good knowledge of the micromorphology of different plant tissues. More rapid proxy methods have been developed, such as determining fecal n-alkanes (Bugalho et al. 2004; Dove and Mayes 2005) or DNA barcoding of plant fragments within feces (Valentini et al. 2009a), but these remain relatively expensive, and their ability to accurately quantify the proportion of different plant species in complex diets is uncertain (Bugalho et al. 2002; Valentini et al. 2009b). There is, therefore, a need to develop rapid and cost-efficient ways for estimating the diets of large herbivores through fecal analyses.

Recent research has shown that near infrared spectroscopy (NIRS) of fecal samples could be used to assess the dietary composition of free-ranging domestic livestock (Landau et al. 2006; Dixon and Coates 2009; Walker et al. 2010). NIRS is a nondestructive procedure requiring minimal sample preparation, and it produces a spectrum expressing the entire chemical makeup of the sample. NIRS may thus be used to predict both the chemical (Malley et al. 2002; Stuth et al. 2003) and

The project was funded by an NSERC-RDC team grant (S.D.C.) and an NSERC-Discovery Grant (R.L.B.).

Correspondence: Pierre-Olivier Jean, Dép. de biologie, Université de Sherbrooke, 2500 boul. de l'Université Sherbrooke, QC, J1K 2R1, Canada. Email: p-o.jean@usherbrooke.ca

Manuscript received 25 July 2013; manuscript accepted 6 January 2014.

© 2014 The Society for Range Management

functional (McIlwee et al. 2001; Stolter et al. 2006) properties of materials. There is an opportunity, therefore, to explore the potential and limitations of NIRS to discriminate different diets of wild herbivores through the spectral analysis of their feces.

A second potentially rapid and cost-effective approach to assess the diets of wild herbivores would be to use fecal chemical properties such as fecal nitrogen (N) or proximate carbon (C) fractions. These have traditionally been used to assess the nutritional and energetic quality of different diets or habitats, but not to estimate the actual botanical composition of the diet. For example, fecal N may be a good index of seasonal variations in diet quality (Verheyden et al. 2011), provided that the amount of tannins in the diet is low (Leslie et al. 2008). Likewise, Hodgman and Davitt (1996) proposed that neutral detergent fiber (NDF) content of feces be used as a predictor for digestible energy of the diet. Given that different forage types vary in their nutritional and energetic qualities, there is an opportunity to explore the ability of fecal chemical properties to estimate the actual botanical composition of the diets of wild herbivores. For example, fecal N might be used to detect the intake of nitrophilous plant species, whereas NDF might be used to identify the most digestible plant species. We posit that this potential to detect diet composition via fecal chemical properties increases as the number of available forage types decreases.

Using NIRS or fecal chemical properties to estimate the diets of wild herbivores first requires a calibration data set of fecal samples from animals with known diets. Knowledge of an animal's recent food intake may be obtained directly through feeding trials on captive individuals, or by reliable microhistological data. Feeding trials may yield more accurate data than microhistological analyses, but are relatively expensive and logistically complicated to perform on wild animals in remote regions. Furthermore, feeding trials usually comprise a limited number of distinct diets, which underrepresents the wide range of diets that are possible in the wild. On the other hand, microhistological data have been criticized as being imprecise (e.g., Wam and Hjeljord 2010), leading Walker et al. (2010) to speculate that these cannot be used as calibration data sets for NIRS. This remains, however, a contentious matter given that Taillon et al. (2006) found an almost perfect concordance between the amounts of fir and spruce needles fed to captive white-tailed deer (*Odocoileus virginianus* Zimmermann) fawns and the percentages of these two forage types within feces. There is, therefore, a need to test the resolution at which NIRS or fecal chemical properties can predict simple diets of wild herbivore based on both types of calibration data sets (i.e., feeding trials and microhistological data).

We report on a study where we evaluated both the potential and limitations of NIRS and fecal chemical properties to predict the proportion of different plant groups composing the winter diets of white-tailed deer on Anticosti Island, Canada. The island boasts a high population of this herbivore, which makes it a prized area for game hunting. There are concerns, however, that overbrowsing of certain winter forage types could result in an important crash of the population before 2100 (Potvin et al. 2004). For this reason, it is important to propose rapid and cost-effective tools that could help rangeland managers on Anticosti Island to monitor the depletion of resources during winter. Fecal samples were obtained, there-

fore, from two prior studies; one in which distinct diets had been administered to captive deer, and the other in which diets of free-ranging deer had been estimated from microhistological analyses.

METHODS

Study Area

Anticosti Island (lat 49°28'N, long 63°00'W) is located in the Gulf of St. Lawrence, Canada. The entire island (7943 km²) is located in the Eastern balsam fir-White birch bioclimatic zone (Grondin et al. 1996). The subboreal maritime climate provides cool summers (mean of 16°C in July) and cold winters (mean of -11°C in January), with an average annual precipitation of 630 mm of rain and 406 cm of snow (Environment Canada 2006).

About 200 white-tailed deer were introduced onto the island in 1896, and the population increased rapidly in the absence of natural predators. Accordingly, the average white-tailed deer density over different sectors of the island has been 20 to 60 individuals · km⁻² for the past 80 yr (Potvin et al. 2004). Heavy browsing has resulted in a loss of palatable species in the shrub layer such as Mountain maple (*Acer spicatum* Lam.), Beaked hazel (*Corylus cornuta* Marsh.), and Canadian yew (*Taxus canadensis* Marsh; Potvin et al. 2003), and in the spread of browse tolerant graminoids such as Canada bluejoint (*Calamagrostis canadensis* [Michx.] Dufresne et al. 2009). Balsam fir (*Abies balsamea* [L.] P.Mill.) is the preferred winter forage (Lefort et al. 2007), such that the recruitment of balsam fir seedlings has substantially diminished in the understory giving way to a higher recruitment of less palatable white spruce seedlings (*Picea glauca* [Moench] Voss.; Hidding et al. 2013). In accordance with this shift in the understory vegetation, it is expected that white-tailed deer will increase their winter consumption of white spruce seedlings in coming years (Sauvé 2005; Taillon et al. 2006).

Fecal Samples

In study 1, Taillon et al. (2006) tested the effects of two winter diets on the behavior and body condition of captive white-tailed deer fawns. In 2004, seven fawns were fed a control winter diet, approximating the proportional intake of balsam fir (70%), white spruce (20%), and arboreal lichens (10%) for white-tailed deer on Anticosti Island (Lefort et al. 2007). Six other fawns were fed a "poor quality" diet consisting of 50% balsam fir, 40% white spruce, and 10% arboreal lichens. Feces were collected on a daily basis (control diet: $n=69$, poor diet: $n=63$) and kept frozen at -20°C until further analyses.

In study 2, performed in 2009–2010, Giroux et al. (2012) conducted a field survey on Anticosti Island to determine the influence of winter diet on body condition of free-ranging white-tailed deer. Using a combination of telemetry and the bio-marking of feces with food dyes, they followed 27 individuals and recovered fresh feces ($N=90$). Within 6 h after collection, fecal samples were dried at 20°C and subsequently stored at -4°C. A subsample of each fecal sample was kept for NIRS scanning and chemical analyses (explained below), whereas the remaining fecal material was analyzed for its plant content by microhistological analysis (Lefort et al. 2007).




Figure 1. Results of principal component-based discriminant analysis showing the Mahalanobis distances between the near infrared spectra of fecal samples collected from captive deer that had been fed two different diets (Control diet=70% balsam fir+20% white spruce+10% lichens; Poor diet=50% balsam fir+40% white spruce+10% lichens). A total of 102 samples were used for model calibration and 30 samples for validation.

Coniferous, deciduous, as well as herbaceous and lichenous fragments were identified by comparing with reference samples under a microscope. Among the coniferous fragments, balsam fir and white spruce needles were distinguished based on their micromorphology. The proportion of lichens was estimated at 16 \times magnification, whereas the proportions of the other forage types were estimated at 100 \times magnification, according to the protocol developed by Lefort (2002).

Fecal Chemical Properties

The stored fecal subsamples from study 2 (Giroux et al. 2012) were thawed, dried again in an air-draft oven (50 $^{\circ}$ C), and ground in a mortar to pass a 2-mm sieve. Fecal N was measured by high temperature combustion followed by gas analysis, using a Vario Macro C&N Analyzer (Elementar Analysensysteme Corp, Hanau, Germany). NDFs, acid detergent fibers (ADFs), and acid detergent lignin (ADL) were measured following the protocol of Goering and Van Soest (1970), using an Ankom Fiber Analyzer 200 (Ankom Technology, Macedon, NY). Cellulose was estimated as the difference between ADL and ADF (Van Soest 1994).

Near Infrared Spectroscopy

Air-dried and sieved fecal subsamples of both study 1 and study 2 were scanned for the reflectance of wavelengths ranging from 1 000 to 2 500 nm, using a Nicolet Antaris FT-NIR spectroscope supported with Omnic software (ThermoFisher Scientific, Waltham, MA). In study 1, a principal component (PC) based discriminant analysis using Mahalanobis distances was used to discriminate the NIR spectra of feces from white-tailed deer fed with control versus poor quality winter diets. This predictive model was calibrated using 102 fecal subsamples, and the remaining 30 subsamples were used for its validation. The

nonrandomness in the classification of fecal samples among the two diets was evaluated using a χ^2 test. For study 2, we used partial least-squares regressions (PLSR; Wold 1966) to build predictive models relating microhistological data to the NIR spectra of 75 fecal samples. First derivatives and mathematical smoothing features from the TQ Analyst software (ThermoFisher Scientific) were used to improve the predictive power of the calibrated PLSR models (Savitzky and Golay 1964; Williams and Norris 1987). The remaining spectra from 15 randomly selected fecal samples were used to validate each PLSR model and evaluate its robustness based on the ratio of prediction to deviation (RPD=ratio of the standard deviation of predictions to the observed standard deviation). According to Williams (2001), model calibrations are deemed reliable only if $R^2 > 0.49$ and $RPD > 2.3$.

Fecal Chemistry Models

Linear mixed-effects models (*nlme* R package; Pinheiro and Bates 2000) were used to relate fecal chemistry to microhistological data stemming from study 2 (Giroux et al. 2012). We considered the identity of the deer as a random effect variable, which required us to drop one fecal sample, the identity of which was unknown ($N=89$). We limited the number of explanatory variables to ADL, cellulose, and fecal N in order to avoid multicollinearity among fecal chemical properties and to maintain variance inflation factors below 2.5 (Graham 2003). ADL and cellulose were expressed on an NDF basis in order to reduce the bias induced by variations in labile substrates in the feces (e.g., microbial biomass, endodermal losses, etc.; Van Soest 1994). Microhistological data were either arcsine or arcsine square root-transformed, in order to meet the assumption of homoscedasticity (Gilbert 1989). We used the Akaike information criterion (Akaike 1974; Hurvich and Tsai 1989) corrected for small sample size (AICc) to select the best fitting model for each forage type.

RESULTS

PC-based discriminant analysis is a modeling method that classifies each fecal sample into one of two predefined classes (i.e., diets), based on a set of PCs that were derived from NIR absorbance values. The classification of a given fecal sample is based on the shortest Mahalanobis distance (i.e., H value) to the mean of each class. Thus, PC-based discriminant analysis of NIR spectra of feces from two different diets used in study 1 correctly classified 28 of 30 validation samples (Fig. 1). Altogether, 115 of 132 fecal samples were correctly assigned to the correct diet ($\chi^2=72.8$, $P < 0.001$).

Microhistological analyses from study 2 reported fecal samples with 47–100% conifer content ($\bar{X}=81\%$), 0–44% deciduous content ($\bar{X}=7\%$), and 0–17% lichen content ($\bar{X}=5\%$). Among the conifer fragments, balsam fir and white spruce respectively represented 0–93% ($\bar{X}=40\%$) and 0–90% ($\bar{X}=41\%$) of the total forage fragments in the feces. PLSR regression models using NIR spectral information successfully predicted ($R^2=0.89$, $RPD=3.2$) the proportion of conifer fragments as determined by microhistological analyses (Fig. 2). However, PLSR models were not robust for predicting the proportions of deciduous ($R^2=0.64$, $RPD=1.1$), balsam fir




Figure 2. Correlation between coniferous content of fecal samples as determined by microhistology and that predicted by a partial least-squares regression model of fecal near infrared spectra. A total of 75 samples were used for model calibration and 15 samples for validation. RMSEC indicates root mean squared error of calibration; RMSEP, root mean squared error of prediction.

($R^2=0.68$, $RPD=1.4$), white spruce ($R^2=0.55$, $RPD=1.1$) or lichen ($R^2=0.36$, $RPD=1.0$) content of fecal samples.

Linear mixed-models applied to fecal chemistry data from study 2 best predicted the conifer content of fecal samples, with 47% of the variance explained when using ADL as the only explanatory variable (Table 1). For predicting the deciduous content of feces, the best fitting model explained 24% of the variance, and included ADL and cellulose as explanatory variables. Fecal ADL explained 22% of the variance in balsam fir content, whereas ADL and cellulose could only explain 8% of the variance for white spruce. No model could explain the variance in lichen content by fecal chemistry. Fecal N was not a predictor of microhistological data in any of the models.

DISCUSSION

Our study showed that NIR spectra of feces from study 1 could discriminate between animals consuming two diets differing only in the relative proportion of balsam fir and white spruce. In the context of Anticosti Island, this may have practical implications, given that the relative abundance of white spruce to balsam fir should continue to increase over coming years. NIRS could, therefore, be used to track a possible concomitant shift in white-tailed deer winter diets. For example, it would be relatively inexpensive and easy to collect a large number of winter fecal samples over consecutive years, and to scan these by NIRS. Using our PC-based discriminant model, the average statistical distance of fecal samples to each calibration diet type could then be computed in order to detect an increasing use of white spruce in the winter diets of white-tailed deer.

The potential for NIRS to discriminate between two related simple diets could also be applied to answer more fundamental questions on the ecology of white-tailed deer on Anticosti Island. For example, one of the outcomes of the “Optimal

Foraging Theory” (MacArthur and Pianka 1966) is that individuals sharing a common home range should converge towards the same diet. Empirical evidence has shown, however, that nonrandom variations may occur in the feeding behavior of neighboring individuals of the same age group (Searle et al. 2010). Hanley (1997) discussed the possible role of learned behavior and other social factors that may create this variation in diet choices among individuals. Thus, analyzing the multivariate dispersion among fecal NIR spectra (e.g., Anderson 2004) could be a novel and cost-efficient way of measuring the degree of variation in diets within a given home range.

Although PC-based discriminant analysis successfully sorted fecal spectral data according to two different diets used in study 1, a regression approach to analyzing both NIR spectra and fecal chemical properties in study 2 had limited success in estimating the relative abundance of different forage types. This is consistent with past studies that had limited success in developing robust predictive models for estimating the specific forage composition of diets based on NIR spectra (e.g., Walker et al. 1998; Landau et al. 2004). It may be argued that our limited success was due to the unreliability of microhistological data as a calibration tool for NIRS, as suggested by Walker et al. (2010). But, as previously mentioned, Taillon et al. (2006) reported a high correlation between the spruce and fir contents of fecal samples as determined by microhistology, and the actual proportion of these two forage types in the diets of captive white-tailed deer on Anticosti Island. These forage types are especially relevant to rangeland management on Anticosti Island, as they are among the main winter staples for white-tailed deer and because they differ in nutritional quality (Sauvé and Côté 2007). It is, therefore, unfortunate that NIRS and fecal chemical properties could not accurately predict the proportion of these two forage fragments in fecal samples from study 2.

The salient finding of our study is that both NIRS and fecal chemical properties were relatively good at predicting the coniferous content of feces from free ranging white-tailed deer (i.e., study 2). We speculate that coniferous forage is a lot less digestible than the other forage types found on Anticosti Island,

Table 1. Best-fitting mixed-linear regression models predicting the fecal content of various forages based on fecal chemical properties ($N=89$).

Response variable	Explanatory variable	Coefficient	Standard error	<i>P</i>	Explained variance (%)
Conifers	ADL ¹	5.7	0.9	< 0.001	47
	Intercept	-1.0	0.3	0.003	
Deciduous	ADL	-4.2	1.3	0.001	24
	Cellulose	3.2	1.6	0.049	
Balsam fir	Intercept	0.6	0.8	0.468	22
	ADL	7.3	2.1	< 0.001	
White spruce	Intercept	-1.4	0.7	< 0.001	8
	ADL	-2.6	1.6	0.111	
Lichens	Cellulose	-4.9	2.2	0.028	
	Intercept	4.0	1.0	< 0.001	
Lichens	Intercept (null model)	0.38	0.02	< 0.001	

¹ADL indicates acid detergent lignin.

which facilitated its detection by NIRS. This is a relatively new application for fecal NIR spectra, as previous research on wild ruminants has so far only succeeded in using NIRS to estimate diet nutritional quality rather than the botanical composition of different diets (Kamler et al. 2004; Showers et al. 2006). The fact that we succeeded is probably due to the fact that ecosystems on Anticosti Island are relatively simple, with only a few winter forage types available to browsers. Although we were hoping to predict specific forage types, being able to predict the percent conifer content is nevertheless useful, as it is a good indicator of quality food shortages during the winter (Häsler and Senn 2012).

The fact that fecal ADL was positively correlated to conifer fragments (i.e., needles and twigs), and negatively correlated to deciduous fragments (i.e., twigs and young bark), suggests that the former may have been richer in lignin than the latter. We have difficulty explaining, however, why fecal ADL correlated to balsam fir fragments but not to white spruce fragments, as both forage types contain approximately the same amounts of lignin (Sauvé and Côté 2007). Dumont et al. (2005) demonstrated a positive relationship between different assumed bite sizes of balsam fir forage by white-tailed deer and the ADL concentration of each morsel. Thus, the relationship between fecal ADL and balsam fir fragments may reflect larger bites by white-tailed deer when browsing balsam fir, their preferred winter forage type.

IMPLICATIONS

Our study has shown the potential and limitations of fecal NIR spectra, and of fecal chemical properties, as rapid and cost-efficient indices of the winter diet composition of white-tailed deer on Anticosti Island. Both of these tools could be used, therefore, to launch a low cost monitoring program of winter resources for the white-tailed deer in Anticosti. Only a few fecal samples per relevant home range would be required each year to detect changes in winter-feeding behaviors. The low costs associated to both analytical methods allow rangeland managers to substantially increase their sampling efforts.

ACKNOWLEDGMENTS

We thank J. Taillon, W. F. J. Parsons, S. Rioux-Paquette, S. Rivard, B. Baillargeon, S. De Bellefeuille, C. Hins, D. Duteau, J.-F. Adam, B. Chénard, J.-P. Harnois, A. Hemery, J. Hénault-Richard, K. Ladhiri, G. Laprise, L. L'Italien, J. Malouin, D. Morin, G. Reuchet, L. Richard, M. O'Connor, M. Sicilia, R. Théoret-Gosselin, and A. Veillette for technical and logistical support.

LITERATURE CITED

AKAIKE, H. 1974. A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 19:716–723.

ANDERSON, M. J. 2004. PERMDISP: a FORTRAN computer program for permutational analysis of multivariate dispersions (for any two-factor ANOVA design) using permutation tests. Auckland, New Zealand: Department of Statistics, University of Auckland.

BUGALHO, M. N., H. DOVE, AND W. KELMAN. 2004. Plant wax alkanes and alcohols as herbivore diet composition markers. *Rangeland Ecology & Management* 57:259–268.

BUGALHO, M. N., R. W. MAYES, AND J. A. MILNE. 2002. The effects of feeding selectivity on the estimation of diet composition using the n-alkane technique. *Grass and Forage Science* 57:224–231.

DALERUM, F., AND A. ANGERBJÖRN. 2005. Resolving temporal variation in vertebrate diets using naturally occurring stable isotopes. *Oecologia* 144:647–658.

DEARDEN, B. L., R. E. PEGAU, AND R. M. HANSEN. 1975. Precision of microhistological estimates of ruminant food habits. *The Journal of Wildlife Management* 39:402–407.

DIXON, R., AND D. COATES. 2009. Review: near infrared spectroscopy of faeces to evaluate the nutrition and physiology of herbivores. *Journal of Near Infrared Spectroscopy* 17:1–31.

DOVE, H., AND R. W. MAYES. 2005. Using n-alkanes and other plant wax components to estimate intake, digestibility and diet composition of grazing/browsing sheep and goats. *Small Ruminant Research* 59:123–139.

DUFRESNE, M., R. L. BRADLEY, J.-P. TREMBLAY, M. POULIN, AND S. PELLERIN. 2009. Clearcutting and deer browsing intensity interact in controlling nitrification rates in forest floor. *Ecoscience* 16:361–368.

DUMONT, A., J.-P. OUELLET, M. CRÉTE, AND J. HUOT. 2005. Winter foraging strategy of white-tailed deer at the northern limit of its range. *Ecoscience* 12:476–484.

ENVIRONMENT CANADA. 2006. Climate normals and averages daily data reports of Port-Menier's station from 1995 to 2005. Available at: <http://www.climate.weatheroffice.ec.gc.ca>. Accessed 8 June 2006.

GILBERT, N. 1989. Biometrical interpretation. Making sense of statistics in biology. New York, NY, USA: Oxford University Press. 146 p.

GIROUX, M.-A., C. DUSSAULT, N. LECOMTE, J.-P. TREMBLAY, AND S. D. CÔTÉ. 2012. A new way of assessing foraging behaviour at the individual level using faeces marking and satellite telemetry. *PloS one* 7:e49719.

GOERING, H. K., AND P. J. VAN SOEST. 1970. Forage fiber analysis apparatus (reagents, procedures, and some applications). Agriculture handbook No.379. Washington, DC, USA: US Department of Agriculture, Agricultural Research Service. 20 p.

GRAHAM, M. H. 2003. Confronting multicollinearity in ecological multiple regression. *Ecology* 84:2809–2815.

GRONDIN, P., C. ANSSEAU, L. BÉLANGER, J.-F. BERGERON, Y. BERGERON, A. BOUCHARD, J. BRISSON, L. DE GRANDPRÉ, G. GAGNON, C. LAVOIE, G. LESSARD, S. PAYETTE, P. J. H. RICHARD, J. P. SAUCIER, L. SIROIS, AND L. VASSEUR. 1996. Cadre bioclimatique de référence des régions écologiques du Québec. In: *Manuel de Foresterie*. Québec, Canada: Les Presses de l'Université Laval. p. 134–279. (in French)

HANLEY, T. A. 1997. A nutritional view of understanding and complexity in the problem of diet selection by deer (Cervidae). *Oikos* 79:209–218.

HANLEY, T. A., AND J. D. MCKENDRICK. 1985. Potential nutritional limitations for black-tailed deer in a spruce-hemlock forest, southeastern Alaska. *The Journal of Wildlife Management* 49:103–114.

HÄSLER, H., AND J. SENN. 2012. Ungulate browsing on European silver fir *Abies alba*: the role of occasions, food shortage and diet preferences. *Wildlife Biology* 18:67–74.

HIDDING, B., J.-P. TREMBLAY, AND S. D. CÔTÉ. 2013. A large herbivore triggers alternative successional trajectories in the boreal forest. *Ecology* 94:2852–2860.

HODGMAN, T. P., AND B. B. DAVITT. 1996. Monitoring mule deer diet quality and intake with fecal indices. *Journal of Range Management* 49:215–222.

HURVICH, C. M., AND C. L. TSAI. 1989. Regression and time series model selection in small samples. *Biometrika* 76:297–307.

KAMLER, J., M. HOMOLKA, AND D. CIZMÁR. 2004. Suitability of NIRS analysis for estimating diet quality of free-living red deer *Cervus elaphus* and roe deer *Capreolus capreolus*. *Wildlife Biology* 3:235–240.

LANDAU, S., T. GLASSER, AND L. DVASH. 2006. Monitoring nutrition in small ruminants with the aid of near infrared reflectance spectroscopy (NIRS) technology: A review. *Small Ruminant Research* 61:1–11.

LANDAU, S., T. GLASSER, L. DVASH, AND A. PEREVOLTSKY. 2004. Fecal NIRS to monitor the diet of Mediterranean goats. *South African Journal of Animal Science* 34:76–80.

LEFORT, S. 2002. Habitat hivernal du cerf de Virginie (*Odocoileus virginianus*) à l'île d'Anticosti [thesis]. Québec, Canada: Université Laval. 87 p. (in French)

LEFORT, S., J.-P. TREMBLAY, F. FOURNIER, F. POTVIN, AND J. HOUT. 2007. Importance of balsam fir as winter forage for white-tailed deer at the northeastern limit of their distribution range. *Ecoscience* 14:109–116.

- LESLIE, D. M., JR., R. T. BOWYER, AND J. A. JENKS. 2008. Facts from feces: nitrogen still measures up as a nutritional index for mammalian herbivores. *Journal of Wildlife Management* 72:1420–1433.
- MACARTHUR, R. H., AND E. R. PIANKA. 1966. On optimal use of a patchy environment. *American Naturalist* 100:603–609.
- MALLEY, D. F., L. YESMIN, AND R. G. EILERS. 2002. Rapid analysis of hog manure and manure-amended soils using near-infrared spectroscopy. *Soil Science Society of America Journal* 66:1677.
- MCLLWEE, A. M., I. R. LAWLER, S. J. CORK, AND W. J. FOLEY. 2001. Coping with chemical complexity in mammal-plant interactions: near-infrared spectroscopy as a predictor of Eucalyptus foliar nutrients and of the feeding rates of folivorous marsupials. *Oecologia* 128:539–548.
- MCLNNIS, M. L., M. VAVRA, AND W. C. KRUEGER. 1983. A comparison of four methods used to determine the diets of large herbivores. *Journal of Range Management* 36:302–306.
- PARKER, K. L., M. P. GILLINGHAM, T. A. HANLEY, AND C. T. ROBBINS. 1999. Energy and protein balance of free-ranging black-tailed deer in a natural forest environment. *Wildlife Monographs* 143:3–48.
- PINHEIRO, J. C., AND D. M. BATES. 2000. Mixed-effects models in S and S-PLUS. New York, NY, USA: Springer. 528 p.
- POTVIN, F., P. BEAUPRÉ, AND C. DUSSAULT. 2004. La population de cerfs de l'île d'Anticosti: élaboration de modèles densité-habitat et prévisions pour l'horizon 2000–2100, Report; November 2004; Québec, Canada: NSERC-Produits forestiers Anticosti Industrial Research Chair. 60 p. (in French)
- POTVIN, F., P. BEAUPRÉ, AND G. LAPRISE. 2003. The eradication of balsam fir stands by white-tailed deer on Anticosti Island, Québec: a 150-year process. *Ecoscience* 10:487–495.
- SAUVÉ, D. 2005. Les effets de la détérioration de la qualité de l'alimentation d'hiver sur la condition corporelle du cerf de Virginie de l'île d'Anticosti [thesis]. Québec, Canada: Université Laval. 116 p. (in French)
- SAUVÉ, D. G., AND S. D. COTÉ. 2007. Winter forage selection in white-tailed deer at high density: Balsam fir is the best of a bad choice. *Journal of Wildlife Management* 71:911–914.
- SAVITZKY, A., AND M. J. E. GOLAY. 1964. Smoothing and differentiation of data by simplified least squares procedures. *Analytical Chemistry* 36:1627–1639.
- SEARLE, K. R., L. P. HUNT, AND I. J. GORDON. 2010. Individualistic herds: individual variation in herbivore foraging behavior and application to rangeland management. *Applied Animal Behaviour Science* 122:1–12.
- SHIPLEY, L. A., S. BLOMQUIST, AND K. DANELL. 1998. Diet choices made by free-ranging moose in northern Sweden in relation to plant distribution, chemistry, and morphology. *Canadian Journal of Zoology* 76:1722–1733.
- SHOWERS, S. E., D. R. TOLLESON, J. W. STUTH, J. C. KROLL, AND B. H. KOERTH. 2006. Predicting diet quality of white-tailed deer via NIRS fecal profiling. *Rangeland Ecology & Management* 59:300–307.
- STEWART, D. R. M. 1967. Analysis of plant epidermis in faeces: a technique for studying the food preferences of grazing herbivores. *Journal of Applied Ecology* 4:83–111.
- STOLTER, C., R. JULKUNEN-TIITTO, AND J. U. GANZHORN. 2006. Application of near infrared reflectance spectroscopy (NIRS) to assess some properties of a sub-arctic ecosystem. *Basic and Applied Ecology* 7:167–187.
- STUTH, J. W., A. JAMA, AND D. R. TOLLESON. 2003. Direct and indirect means of predicting forage quality through near infrared reflectance spectroscopy. *Field Crops Research* 84:45–56.
- TAILLON, J., D. G. SAUVÉ, AND S. D. COTÉ. 2006. The effects of decreasing winter diet quality on foraging behavior and life-history traits of white-tailed deer fawns. *Journal of Wildlife Management* 70:1445–1454.
- VALENTINI, A., C. MIQUEL, M. A. NAWAZ, E. BELLEMANN, E. COISSAC, F. POMPANON, L. GIJELY, C. CRUAUD, G. NASCETTI, P. WINCKER, J. E. SWENSON, AND P. TABERLET. 2009a. New perspectives in diet analysis based on DNA barcoding and parallel pyrosequencing: the trnL approach. *Molecular Ecology Resources* 9:51–60.
- VALENTINI, A., F. POMPANON, AND P. TABERLET. 2009b. DNA barcoding for ecologists. *Trends in Ecology and Evolution* 24:110–117.
- VAN SOEST, P. J. 1994. Nutritional ecology of the ruminant. Ithaca, NY, USA: Cornell University Press. 476 p.
- VERHEYDEN, H., L. AUBRY, J. MERLET, P. PETIBON, B. CHAUVEAU-DURIOT, N. GUILLON, AND P. DUNCAN. 2011. Faecal nitrogen, an index of diet quality in roe deer *Capreolus capreolus*? *Wildlife Biology* 17:166–175.
- WALKER, J. W., E. J. CAMPBELL, R. W. KOTT, S. LANDAU, C. J. LUPTON, C. B. SCOTT, L. SURBER, C. A. TAYLOR, JR., AND W. R. WHITWORTH. 2010. Chapter 5. Fecal NIRS for predicting botanical composition of herbivore diets. In: J. W. Walker, D. R. Tolleason, and S. Byrns [eds.]. Shining light on manure improves livestock and land management. Jefferson, MO, USA: Texas AgriLife Research and Society for Range Management. p. 53–65.
- WALKER, J. W., D. H. CLARK, AND S. D. MCCOY. 1998. Fecal NIRS for predicting percent leafy spurge in diets. *Journal of Range Management* 51:450–455.
- WAM, H. K., AND O. HJELJORD. 2010. Moose summer diet from feces and field surveys: a comparative study. *Rangeland Ecology & Management* 63:387–395.
- WILLIAMS, P., AND K. H. NORRIS. 1987. Near-infrared technology in the agricultural and food industries. St. Paul, MN, USA: American Association of Cereal Chemists. 330 p.
- WILLIAMS, P. C. 2001. Implementation of near-infrared technology. In: P. C. Williams and K. H. Norris [eds.]. Near-infrared technology in the agricultural and food industries. St. Paul, MN, USA: American Association of Cereal Chemists. p. 145–170.
- WOLD, H. 1966. Estimation of principal components and related models by iterative least squares. In: P. R. Krishnaiah [ed.]. Multivariate analysis. New York, NY, USA: Academic press. p. 391–420.