

# Damage from intraspecific combat is costly

Zachary Emberts<sup>1</sup>, Ummat Somjee<sup>2</sup>, John J. Wiens<sup>1</sup>

<sup>1</sup> Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, 85721,  
USA

<sup>2</sup> Smithsonian Tropical Research Institute, Balboa, Ancón, 211-8000, Panamá

*Corresponding Author*

Zachary Emberts

<sup>1</sup> Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, 85721,  
USA

E-mail: [emberts@email.arizona.edu](mailto:emberts@email.arizona.edu)

## **Abstract**

When individuals engage in fights with conspecifics over access to resources, injuries can occur. Most theory suggests that the costs associated with these injuries should influence an individual's decision to retreat from a fight. Thus, damage from intraspecific combat is frequently noted and quantified. However, the fitness-related costs associated with this damage are not. Quantifying the cost of fighting-related damage is important because most theory assumes that it is the cost associated with the damage (not the damage itself) that should influence an individual's decision to retreat. Here, we quantified the cost of fighting-related injuries in the giant mesquite bug, *Thasus neocalifornicus*. We demonstrate that experimentally simulated fighting injuries result in metabolic costs and costs to flight performance. We also show that flight costs are more severe when the injuries are larger. Overall, our results provide empirical support for the fundamental assumption that damage acquired during intraspecific combat is costly.

*Keywords:* damage, fight, injury, intraspecific competition, intrasexual competition, male-male combat

## INTRODUCTION

Many animals fight with conspecifics over access to limited resources (e.g. food, shelter, and mates; Smith and Price 1973; West- Eberhard 1983; Rico-Guevara and Hurme 2019). During these fights, injuries can occur. Examples include bite wounds in lizards (Lailvaux et al. 2004), puncture wounds in shrimp (Rojas et al. 2012), and scratch wounds in lemurs (Kappeler 1997). Most theoretical models assume that such injuries influence an individual's decision to retreat from a fight, which ultimately influences who wins (Parker and Rubenstein 1981; Hammerstein and Parker 1982; Enquist and Leimar 1983; Payne 1998; Lane and Briffa 2017; Palaoro and Briffa 2017). Experimental studies have recently found support for this assumption (Emberts and Wiens 2021). Damage (that is accrued during combat) is thought to influence an individual's willingness to persist in a fight because damage is assumed to have a fitness-related cost (Parker and Rubenstein 1981; Hammerstein and Parker 1982; Enquist and Leimar 1983; Payne 1998). Thus, understanding the costs associated with fighting-related damage is crucial to understanding intraspecific contests. Despite their importance, much remains unknown about the costs associated with fighting-related injuries.

The assumption that damage from intraspecific combat is costly is a reasonable one. Acute injuries can result in reduced function of damaged body parts, blood loss, and energetically expensive immune responses (Lochmiller and Deerenberg 2000; Ardia et al. 2012). These injuries could reduce an individual's ability to escape predators (e.g. Krause et al. 2017) and secure mates (e.g. Salvador et al. 1995; Emberts et al. 2018). Thus, they may have serious fitness consequences. Nevertheless, the severity of the costs associated specifically with fighting-related injuries are largely unstudied. This is an important omission because it is these costs (not the injuries per se) that are theorized to influence an individual's fighting behavior (Parker and

Rubenstein 1981; Hammerstein and Parker 1982; Enquist and Leimar 1983; Payne 1998; Lane and Briffa 2017; Palaoro and Briffa 2017). Thus, it is crucial to test if combat-related damage is costly and determine whether some injuries are more costly than others.

To investigate the costs of damage from intraspecific fighting, we used the giant mesquite bug, *Thasus neocalifornicus* (Insecta: Hemiptera: Coreidae, Figure 1). Male *T. neocalifornicus* use their enlarged and spined hind legs in fights with conspecific males over access to territories (Embets and Wiens 2021). During these fights, males can puncture their rival's forewings (Figure 1; see also Embets and Wiens 2021). Similar forewing punctures are not found in coreids that lack sexually selected weapons (Embets, Hwang, et al. 2021), suggesting that this type of damage (in males) is unique to intrasexual combat.

There is some *a priori* evidence that forewing punctures are costly. First, melanization often occurs around the wounds (Embets and Wiens 2021), indicating a metabolically expensive immune response (Cerenius and Soderhall 2004; Ardia et al. 2012). Such an immune response may temporarily increase an individual's resting metabolic rate (RMR; Ardia et al. 2012). Second, damage occurs on their wings (Figure 1), which may reduce flying ability (Combes et al. 2010; Mountcastle et al. 2016). Thus, we predicted that damage associated with male-male competition should have both metabolic and locomotory costs. Moreover, the size of the punctures vary considerably (Embets, Hwang, et al. 2021; Figure 1). Thus, we also predicted that larger punctures should be more costly than smaller ones. To investigate these predictions, we first tested whether damage influences an individual's metabolic rate during a resting phase (i.e. standard or resting metabolic rate; Mathot and Dingemanse 2015). We then tested whether forewing damage influences how far an individual can fly, and if larger punctures had greater impact on flying performance.

## **MATERIALS AND METHODS**

### **Experiment 1: metabolic costs**

#### ***Animal collection and husbandry***

Adult male *Thasus neocalifornicus* used for this first experiment were collected in Santa Cruz County, Arizona (south of Tubac; 31.5294, -111.0218). Individuals were collected by hand September 8–16, 2020, given the availability of a FoxBox Respirometry System (Sable Systems Hendersons, NV) at that time. Some individuals had minor wing damage before the experimental treatment. This should not be problematic because metabolic costs of injury were assessed using a before-and-after experimental design (see below). All individuals had their pronotums uniquely marked with paint pens and permanent markers for identification (regardless of treatment, see below). Collected individuals were housed together in large, mesh insect-rearing containers (305x305x607 mm; LxWxH) for up to 96 h (up to 20 individuals per container) at room temperature (~26 °C), and provided with several fresh cuttings of velvet mesquite (*Prosopis velutina*) for food. Interactions within the rearing container are limited under these conditions (e.g. males do not appear to fight, see Emberts and Wiens 2021). Thirty minutes before measuring CO<sub>2</sub> production, individuals were moved into separate containers without food or water.

#### ***Measuring resting metabolic rate***

Flow-through respirometry was used to measure the rate of CO<sub>2</sub> production as a proxy for resting metabolic rate (RMR) for each individual ( $n=27$ ). The CO<sub>2</sub> analyzer was calibrated using CO<sub>2</sub> free dry air and 2000 ppm CO<sub>2</sub> in N<sub>2</sub>. Incurrent air was pulled through the apparatus and first

scrubbed of moisture using a silica bead column, then scrubbed of CO<sub>2</sub> and further moisture using an ascarite and drierite column (WA Hammond Drierite Company, Xenia, OH, USA). This dry CO<sub>2</sub>-free air then passed through an inbuilt pump and flow control system (FoxBox, Sable Systems Hendersons, NV) before being passed through a custom glass cylindrical insect chamber (50ml), and finally through the CO<sub>2</sub> analyzer with a sampling frequency of 1Hz. Gas circulation between instruments occurred through 3mm diameter plastic tubing and connecting plastic valves (Bevaline-IV, Cole Parmer, Vernon Hills, IL, USA).

An individual was placed inside the insect chamber and a measurement was taken once it stopped moving and readings stabilized, a process that took approximately 8 minutes (Lighton 2008). The chamber was placed in a dark foam box with a viewing window to minimize disturbance and allow individuals to reach a phase where movement ceased. Flow rates were set to maximize the temporal resolution of CO<sub>2</sub> production and detectability at a standard temperature and pressure (200ml/min). Temperature was checked throughout the trial with readings from a separate thermometer (Oregon Scientific thermometer) placed beside the measurement chamber, but note that we do not have complete temperature data for two individuals (Supplementary Dataset S1).

Respirometry data were extracted from Expedata software (v1.9.27; Sable Systems), which received signals directly from a built-in analog-to-digital converter in the FoxBox. Molar rates of CO<sub>2</sub> production were then calculated from raw measures of CO<sub>2</sub> (ppm), using the known flow rate and variables in the Ideal Gas Law equation:

$$\text{CO}_2 = \frac{[P \times \text{FR} \times (\text{Fi} - \text{Fe})]}{R \times T}$$

Where  $\dot{V}_{CO_2}$  is the rate of  $CO_2$  production (mols/min), FR is the flow rate (200ml/min),  $F_i$  and  $F_e$  are the fractional concentrations of incurrent and excurrent  $CO_2$ , respectively, P is the pressure (1 standard atmosphere), R is the gas constant at standard temperature and pressure (0.08206 L atm  $K^{-1}mol^{-1}$ ), and T is the temperature (Kelvin). We multiplied moles of  $CO_2$  by the molar volume of an ideal gas (22400 ml/mol at STP) to get volume of  $CO_2$  (ml/sec), and multiplied this by 21.1 J/ml and divided by a respiratory quotient (RQ) of 1 to get our respiration rate in J/s or Watts (Lighton 2008). A RQ of 1 was used for this conversion because *Thasus neocalifornicus*, like many hemipterans, is a sap feeder (Olivier-Espejel et al. 2011), indicating a high carbohydrate diet (sensu Somjee et al. 2018). Baseline measurements with empty chambers were taken at the beginning and end of each run and subtracted from periods of stable minimal readings when individuals were not visibly moving to maximize accuracy and account for potential sensor drift (Lighton 2008).

After measuring RMR, individuals were randomly assigned to one of two treatments using a random number generator: a comparative baseline treatment ( $n=13$ ) or an experimental manipulation ( $n=14$ ). Each individual in the experimental manipulation treatment received two puncture wounds, each 3 mm in diameter. One puncture was placed in the corium of each forewing (one per side), where damage from male-male combat normally occurs (Figure 1). Moreover, two puncture wounds were used as opposed to one because wild-caught *T. neocalifornicus* males that are damaged rarely have a single puncture wound and most have 7 or more (Embets, Hwang, et al. 2021). Finally, the puncture wounds were 3 mm in diameter because previous studies found that wild-caught *T. neocalifornicus* males can have puncture wounds  $>3$  mm in diameter (maximum=3.150 mm; Embets, Hwang, et al. 2021). The average diameter of puncture wounds in wild-caught male *Thasus neocalifornicus* is 0.440 mm (note:

measured individuals in that study were from the same site or nearby those used here). Therefore, 3 mm puncture wounds are relatively large but still biologically relevant. Wounds were made using a XOOOL leather hole puncher (i.e. experimentally induced) to mimic those received during a fight (Supplementary Figure S1). Wounds were made experimentally so that we could isolate the costs associated with the damage itself. Resting metabolic rate was measured (for a second time) approximately 30 minutes after injury because our previous observations suggested that if there was an energetic cost (associated with the injury) we would most likely see it at this time (Somjee et al. 2018). Individuals in the comparative baseline treatment were not punctured, but were handled similarly to those that were. Specifically, these control individuals had their forewings spread and placed on the hole puncher for 3 seconds (per forewing). However, these individual's did not actually have their wings punctured. The same methods described above were used to measure RMR for the second time, and they were taken within 24 hours of the first measurement. The first measurement was then subtracted from the second to calculate each individual's change in RMR. Individuals were then massed to the nearest hundredth of a gram using a TPO-100 digital pocket scale. However, three individuals were not massed because the scale was unavailable. Pronotal width was also measured to the nearest hundredth of a millimeter using a Mitutoyo digital caliper. Pronotal width is a widely used proxy for body size in this clade (Proctor et al. 2012; Emberts et al. 2020).

## **Experiment 2: locomotory costs**

### ***Animal collection and husbandry***

Adult male *Thasus neocalifornicus* used for the second experiment were collected near those for the first, in Pima County, Arizona (SE of Green Valley: 31.7921, -110.8813; S of Vail: 31.9640,

-110.6729). All individuals were collected by hand between June 23<sup>rd</sup> and July 8<sup>th</sup>, 2020. This is when adult *Thasus neocalifornicus* begin to emerge (Emberts and Wiens 2021), making it easier to find fully sclerotized males without wing damage. Individuals were marked, measured (pronotal width), housed, and fed as described above.

### ***Flight trials***

Three hours before experimentation, each individual was randomly assigned (with a random number generator) to one of five treatments ( $n=20$  per treatment). In these treatment they received either: (i) no wing punctures (i.e. comparative baseline), or two wing punctures (one per wing) that were either: (ii) 1 mm (diameter), (iii) 2 mm, (iv) 3 mm, or (v) 4 mm. The size, number, and location of punctures (on the corium) followed the reasoning explained above. The 1 mm punctures were made using a 20 gauge sewing needle, whereas 2, 3, and 4 mm punctures were made using a XOOD leather hole puncher. Individuals in the comparative baseline treatment were not punctured, but were handled similarly to those that were (see above).

Two hours after their treatment, individuals participated in a flying trial. These trials were conducted outside on a manicured grass field (Video S1) in a randomized order. To initiate flight, individuals were held in an observer's hand 1.5 m above the ground. This was a height at which individuals were found naturally (although they can also be found much higher). Individuals were then lightly tapped by the observer to initiate a flight response. Thus, these trials may be considered an anti-predatory response to a human predator. After individuals took off, the horizontal distance from the observer's hand to their landing location was recorded in meters with a Champion Sports open reel measuring tape. Most individuals landed on the ground. However, for individuals that flew over 23 m, it was possible for them to land on an

object (e.g. a tree, a fence). When this happened, the horizontal distance from the starting location to the landing site was recorded. Flying trials were conducted between 10 am and 7 pm at air temperatures from 31–38 °C. Time and temperature were recorded for use as factors in the statistical analyses.

## **Statistical analyses**

### ***Experiment 1: metabolic costs***

First, we tested whether the difference between an individual's first and second RMR measurements (i.e. the dependent variable) could be explained by treatment (i.e. whether they received experimental wing punctures, independent variable) using a generalized linear model (GLM). This model also included pronotal width (body-size proxy) and air temperature (when measuring resting metabolic rates) as covariates. We confirmed that including or excluding these covariates produced qualitatively similar results (Supplementary Tables S1 and S2), as did using an alternative body-size proxy (i.e. mass; Supplementary Tables S1 and S3). Given that individuals in our comparative baseline treatment appeared to have slightly lower resting metabolic rates after their sham treatment (see Results), we also conducted some follow-up analyses. Specifically, we conducted two paired t-tests (since each individual was measured twice) to determine whether (1) the decrease observed in our comparative baseline treatment was statistically significant and (2) whether wing damage itself significantly increased an individual's resting metabolic rate.

### ***Experiment 2: locomotory costs***

We then examined whether wing damage (i.e. treatment, independent variable) influenced how far individuals flew (dependent variable) using another GLM. For this analysis, we included pronotal width (9.46–14.13 mm), air temperature (31–38 °C), and time of day at the start of the trials (10–18.5 h) as covariates. Given a significant result, we then performed a Tukey pairwise comparison to determine which treatments were responsible for driving the observed effect. We confirmed that including or excluding our covariates produced qualitatively similar results (see Results and Supplementary Tables S4–S7), as did including or excluding the 4 mm treatment (i.e. with punctures larger than in wild populations; see Results and Supplementary Tables S8 and S9). All statistical analyses were conducted in R v 4.0.3 (R Development Core Team 2020).

## RESULTS

We found that damage accrued during intrasexual combat can be costly. First, individuals that experimentally received wing punctures appeared to increase their RMR ( $t=2.701$ ,  $df=24$ ,  $P=0.014$ ; Supplementary Table S1; Figure 2). However, visualization of the data (Figure 2) suggested that this significant result may have been driven by individuals in the comparative baseline treatment (which appeared to have lowered their RMR). We confirmed that this was not the case. A paired t-test analysis revealed that individuals in the comparative baseline treatment had similar resting metabolic rates before and after receiving the sham treatment ( $t=-1.755$ ,  $df=12$ ,  $P=0.105$ ). Moreover, individuals in our experimental manipulation treatment had significantly higher RMR after receiving their wing punctures (paired t-test:  $t=2.227$ ,  $df=13$ ,  $P=0.044$ ). These analyses collectively show that receiving forewing damage increased an individual's resting metabolic rate.

The presence and severity of wing punctures also influenced how far individuals could fly (Figure 3). Non-damaged individuals (i.e. comparative baseline treatment) flew 60.4 m on average, whereas individuals with 4 mm puncture wounds flew 0.61 m on average (i.e. a 99% reduction;  $z=-5.904$ ,  $P<0.001$ ; Supplementary Table S6). Individuals with 2 and 3 mm puncture wounds both flew ~24 m on average (i.e. a 60% reduction from non-damaged individuals; 2 mm:  $z=-3.501$ ,  $P=0.004$ ; 3 mm:  $z=-3.766$ ,  $P=0.002$ ; Supplementary Table S6). Finally, individuals with 1 mm puncture wounds flew 40.1 m. This distance was not statistically different from our comparative baseline (40.1 m compared to 60.4;  $z=-1.761$ ,  $P=0.397$ ), but was statistically different from individuals with 4 mm puncture wounds (40.1 m compared to 0.6 m;  $z=-4.184$ ,  $P<0.001$ ; Supplementary Table S6). This latter result showed that the severity of damage influenced flight performance costs.

## DISCUSSION

We investigated whether damage accrued during intrasexual combat is costly using the giant mesquite bug, *Thasus neocalifornicus*. In this species, intrasexual combat frequently results in wing damage (Figure 1; Emberts and Wiens 2021). Thus, we experimentally induced injuries (which mimicked those received during a fight; Supplementary Figure S1) and investigated whether this damage had a metabolic and/or locomotory cost (i.e. two potential indirect fitness costs). First, we found that there was an energetic cost associated with fighting-related injuries. Second, we found that wing damage decreased flying ability, and that this performance cost was more severe when puncture wounds were larger. Overall, these results provide empirical support for the fundamental assumption that damage acquired during intraspecific combat is costly, and illustrate that some fighting-related injuries can be far more costly than others.

Understanding the costs associated with intraspecific fighting is crucial, because theory suggests that these costs influence contest outcomes and behavior (Parker and Rubenstein 1981; Hammerstein and Parker 1982; Enquist and Leimar 1983; Payne 1998). Costs typically associated with intraspecific combat include energetic consumption (Smith and Taylor 1993; Hack 1997; Rovero et al. 2000), reduced function of damaged body parts (Berzin and Caldwell 1983; but see Johnson et al. 2005; Jennings et al. 2017), and predation risk (Hammerstein and Riechert 1988; Jakobsson et al. 1995; Brick 1998). The results of this study further contribute to our understanding of these costs. For example, studies that have previously quantified the energetic costs of combat frequently quantified the costs associated with physical activity (e.g. the amount of energy it takes a cricket to charge at their opponent; Hack 1997). However, our study reveals that physical activity is not the only factor that can contribute to the energetic costs of combat.

Our finding that an acute injury has an energetic cost is not surprising (Lochmiller and Deerenberg 2000; Ardia et al. 2012). However, the observed effect size was limited (an approximately 3% increase in an individual's RMR), especially given the large size of the puncture wounds. In other insects, for example, wounding can increase resting metabolic rates by 28% (Ardia et al. 2012). Moreover, when birds are presented with an immune challenge their resting metabolic rate increases 5 to 15% (Hasselquist and Nilsson 2012). The metabolic costs measured here may be low because fighting-related damage in coreids mainly occurs more-or-less peripherally, on the wings (specifically the corium; Emberts and Wiens 2021), as opposed to the main body. Moreover, given that approximately half of the males sampled from a population have wing punctures (i.e. they occur frequently; Emberts, Hwang, et al. 2021), selection may have promoted physiological mechanisms to minimize their metabolic cost. Another possibility

is that we underestimated how much wing punctures increase resting metabolic rates. Our control group had slightly decreased metabolic rates between the first and second measurements. This pattern suggests that individuals in our experimental manipulation treatment may have also decreased their resting metabolic rates if they had not been punctured. One reason resting metabolic rates might have decreased between the first and second measurements is because the insects had spent more time in the lab (where individuals fed on cuttings as opposed to live plants). Thus, the fact that we still saw an increase in resting metabolic rate for individuals in the experimental treatment suggests that the metabolic cost associated with wing punctures may be even larger than we quantified here. Nevertheless, even small increases in energetic costs can significantly contribute to daily energy expenditure over longer periods of time (Somjee et al. 2018; Somjee 2021). Thus, future studies should investigate how long this increase in RMR is maintained (now that a metabolic cost has been identified). Moreover, since damage from male-male combat can occur in other locations besides the wings (e.g. on the weapons), future studies should compare how the location of an injury influences its energetic costs.

We also found that injuries can reduce the function of the damaged body part. Here, we found that wing damage decreased flying ability. Whether these injuries would have actually influenced flying ability was unclear, because previous studies investigating the functional costs of having a damaged body part have had inconsistent results. For example, damage to weapons (which may have occurred during previous fights) was found to decrease an individual's future fighting ability in some cases (e.g. Espmark Y. 1964; Lincoln G. 1972; Berzin and Caldwell 1983), but not in others (e.g. Johnson et al. 2005; Jennings et al. 2017). These inconsistencies highlight that injuries should not always be assumed to reduce the function of the damaged body.

Another major finding of this study is that the degree of injury directly influenced the costs (Figure 3). Specifically, individuals with 4 mm puncture wounds did not fly as far as individuals with 1 mm puncture wounds (Figure 3). The differing costs associated with different puncture sizes has important implications because theory suggest that it is the cost associated with the damage that should influence contest outcomes and behavior (Parker and Rubenstein 1981; Hammerstein and Parker 1982; Enquist and Leimar 1983; Payne 1998; Lane and Briffa 2017; Palaoro and Briffa 2017). Thus, all else being equal, our study suggests that individual's that inflict larger puncture wounds during a fight should be better at making their rivals flee. Therefore, future work should make sure to quantify the frequency and severity of damage that occurs during a fight, as opposed to simply noting that damage occurred.

Our results also show that damage from some wounds appears to be catastrophic. Specifically, we found that the furthest that individuals flew with 4 mm puncture wounds was 3.2 m, and most of these individuals flew  $<0.5$  m. On the other hand, an individual with 3 mm puncture wounds was observed flying  $>90$  m, and most individuals with 3 mm puncture wounds flew  $>12$  m. Thus, 4 mm puncture wounds appear to prevent individuals from effectively flying. This cost is likely permanent because adult *Thasus neocalifornicus* are in their terminal molts, and hemimetabolous insects cannot regenerate without molting (Gross 1969; Maginnis 2006). The fact that 4 mm puncture wounds appear to be catastrophic is particularly interesting because the largest puncture wound previously reported in wild caught *Thasus neocalifornicus* is 3.150 mm in diameter (Embets, Hwang, et al. 2021). The data from wild-caught populations coupled with the result from our flying trial suggest one of two possibilities. First, that individuals with punctures that are  $>3.150$  mm in diameter do not survive long, making it unlikely for them to be captured and quantified when sampling populations. Alternatively, fighting might never result in

punctures  $>3.150$  mm. To distinguish between these two alternatives, future studies should quantify the size of punctures wounds that occur during controlled fighting trials.

In conclusion, understanding the costs associated with intraspecific fighting is crucial, because most theory suggests that the cost of engaging in combat influences contest outcomes and behavior (Parker and Rubenstein 1981; Hammerstein and Parker 1982; Enquist and Leimar 1983; Payne 1998; Lane and Briffa 2017; Palaoro and Briffa 2017). One hypothesized cost of engaging in a fight is being injured. However, this assumption has largely gone untested. In this study, we provide empirical evidence suggesting that fighting-related injuries are costly, and that some injuries can be more costly than others.

## **FUNDING**

This work was supported by the National Science Foundation (grant number DBI-1907051) awarded to ZE. We have no conflicts of interest to declare.

## **ACKNOWLEDGEMENTS**

We thank Cody Howard for his help with data collection. We also thank the anonymous reviewers for providing helpful feedback on an earlier version of the manuscript.

## **DATA AVAILABILITY**

Analyses reported in this article can be reproduced using the data provided by Emberts, Somjee, et al. (2021).

## **REFERENCES**

- Ardia DR, Gantz JE, Strebel S. 2012. Costs of immunity in insects: an induced immune response increases metabolic rate and decreases antimicrobial activity. *Funct Ecol.* 26:732–739.
- Berzins IK, Caldwell RL 1983. The effect of injury on the agonistic behavior of the stomatopod, *Gonodactylus bredini* (Manning). *Mar. Freshwater Behav Physiol.* 10:83–96.
- Brick O. 1998. Fighting behaviour, vigilance and predation risk in the cichlid fish *Nannacara anomala*. *Anim Behav.* 56:309–317.
- Cerenius L, Söderhäll K. 2004. The prophenoloxidase-activating system in invertebrates. *Immunol Rev.* 198:116–126.
- Combes SA, Crall JD, Mukherjee S. 2010. Dynamics of animal movement in an ecological context: Dragonfly wing damage reduces flight performance and predation success. *Biol Lett.* 6:426–429.
- Cooper WE. 1999. Tradeoffs between courtship, fighting, and antipredatory behavior by a lizard, *Eumeces laticeps*. *Behav Ecol Sociobiol.* 47:54–59.
- Dunn M, Copelston M, Workman L. 2003. Trade-offs and seasonal variation in territorial defence and predator evasion in the European Robin *Erithacus rubecula*. *Ibis.* 146:77–84.
- Embets Z, St Mary CM, Herrington TJ, Miller CW. 2018. Males missing their sexually selected weapon have decreased fighting ability and mating success in a competitive environment. *Behav Ecol Sociobiol.* 72:1–9.
- Embets Z, St. Mary CM, Howard CC, Forthman M, Bateman PW, Somjee U, Hwang WS, Li D, Kimball RT, Miller CW. 2020. The evolution of autotomy in leaf-footed bugs. *Evolution.* 74:897–910.

- Emberts Z, Wiens JJ. 2021. Defensive structures influence fighting outcomes. *Funct Ecol.* 35:696–704.
- Emberts Z, Hwang WS, Wiens JJ. 2021. Weapon performance drives weapon evolution. *Proc R Soc B* 288:20202898.
- [dataset]\* Emberts Z, Somjee U, Wiens JJ. 2021. Damage from intraspecific combat is costly. Dryad, Dataset. <https://doi.org/10.5061/dryad.0rxwdb0z>
- Enquist M, Leimar O. 1983. Evolution of fighting behaviour: Decision rules and assessment of relative strength. *J Theor Biol.* 102:387–410.
- Espmark Y. 1964. Studies in dominance-subordination relationship in a group of semi-domestic reindeer (*Rangifer tarandus L.*). *Anim Behav.* 12:420–426
- Goss RJ. 1969. The principles of regeneration. New York, NY: Academic Press.
- Hack MA. 1997. The energetic costs of fighting in the house cricket, *Acheta domesticus* L. *Behav Ecol.* 8:28–36.
- Hammerstein P, Riechert SE. 1988. Payoffs and strategies in territorial contests: ESS analyses of two ecotypes of the spider *Agelenopsis aperta*. *Evol Ecol.* 2:115–138.
- Hammerstein P, Parker GA. 1982. The asymmetric war of attrition. *J Theor Biol.* 96:647–682.
- Hasselquist D, Nilsson JÅ. 2012. Physiological mechanisms mediating costs of immune responses: what can we learn from studies of birds? *Anim Behav.* 83:1303–1312.
- Hess S, Fischer S, Taborsky B. 2016. Territorial aggression reduces vigilance but increases aggression towards predators in a cooperatively breeding fish. *Anim Behav.* 113:229–235.
- Jakobsson S, Brick O, Kullberg C. 1995. Escalated fighting behaviour incurs increased predation risk. *Anim Behav.* 49:235–239.
- Jennings DJ, Boys RJ, Gammell, MP. 2017. Weapon damage is associated with contest


- dynamics but not mating success in fallow deer (*Dama dama*). *Biol Lett.* 13:20170565.
- Johnson HE, Bleich VC, Krausman PR, Koprowski JL. 2007. Effects of antler breakage on mating behavior in male tule elk (*Cervus elaphus nannodes*). *Eur J Wildl Res.* 53:9–15.
- Kappeler PM. 1997. Intrasexual selection in *Mirza coquereli*: evidence for scramble competition polygyny in a solitary primate. *Behav Ecol Sociobiol.* 41:115–127.
- Krause J, Herbert-Read JE, Seebacher F, Domenici P, Wilson AD, Marras S, Svendsen MBS, Strömbom D, Steffensen JF, Krause S, Viblanc PE. 2017. Injury-mediated decrease in locomotor performance increases predation risk in schooling fish. *Philos Trans R Soc B.* 372:20160232.
- Lane SM, Briffa M. 2017. The price of attack: rethinking damage costs in animal contests. *Anim Behav.* 126:23–29.
- Lailvaux SP, Herrel A, VanHooydonck B, Meyers JJ, Irschick DJ. 2004. Performance capacity, fighting tactics and the evolution of life-stage male morphs in the green anole lizard (*Anolis carolinensis*). *Proc R Soc B* 271:2501–2508.
- Lighton JR. 2008. *Measuring metabolic rates: a manual for scientists.* Oxford (UK): Oxford University Press.
- Lincoln G. 1972. The role of antlers in the behaviour of red deer. *J Exp Zoo.* 182:233–249.
- Lochmiller RL, Deerenberg C. 2000. Trade-offs in evolutionary immunology: just what is the cost of immunity? *Oikos.* 88:87–98.
- Maginnis TL. 2006. The costs of autotomy and regeneration in animals: a review and framework for future research. *Behav Ecol.* 17:857–872.
- Mathot KJ, Dingemanse NJ. 2015. Energetics and behavior: unrequited needs and new directions. *Trends Ecol Evol.* 30:199–206.

- Mountcastle AM, Alexander TM, Switzer CM, Combes SA. 2016. Wing wear reduces bumblebee flight performance in a dynamic obstacle course. *Biol Lett.* 12:20160294.
- Olivier-Espejel S, Sabree ZL, Noge K, Becerra JX. 2011. Gut microbiota in nymph and adults of the giant mesquite bug (*Thasus neocalifornicus*) (Heteroptera: Coreidae) is dominated by *Burkholderia* acquired de novo every generation. *Environ Entomol.* 40:1102–1110.
- Palaoro AV, Briffa M. 2017. Weaponry and defenses in fighting animals: how allometry can alter predictions from contest theory. *Behav Ecol.* 28:328–336.
- Parker GA, Rubenstein DI. 1981. Role assessment, reserve strategy, and acquisition of information in asymmetric animal conflicts. *Anim Behav.* 29:221–240.
- Payne RJ. 1998. Gradually escalating fights and displays: The cumulative assessment model. *Anim Behav.* 56:651–662.
- Rojas R, Morales MC, Rivadeneira MM, Thiel M. 2012. Male morphotypes in the Andean river shrimp *Cryphiops caementarius* (Decapoda: Caridea): morphology, coloration and injuries. *J Zool.* 288:21–32.
- Rovero F, Hughes RN, Whiteley NM, Chelazzi G. 2000. Estimating the energetic cost of fighting in shore crabs by noninvasive monitoring of heartbeat rate. *Anim Behav.* 59:705–713.
- Salvador A, Martin J, López P. 1995. Tail loss reduces home range size and access to females in male lizards, *Psammmodromus algirus*. *Behav Ecol.* 6:382–387.
- Smith IP, Taylor AC. 1993. The energetic cost of agonistic behaviour in the velvet swimming crab, *Necora* (= *Liocarcinus*) *puber* (L.). *Anim Behav.* 45:375–391.
- Smith JM, Price GR. 1973. The logic of animal conflict. *Nature.* 246:15–18.
- Somjee U, Woods HA, Duell M, Miller CW. 2018. The hidden cost of sexually selected traits:


the metabolic expense of maintaining a sexually selected weapon. *Proc R Soc B*  
285:20181685.

Somjee U. 2021. Positive allometry of sexually selected traits: Do metabolic maintenance costs  
play an important role? *BioEssays*. 2000183.


West-Eberhard MJ. 1983. Sexual selection, social competition, and speciation. *Q Rev Biol*.  
58:155–183.



**Figure 1** Male *Thasus neocalifornicus* (A) use their weaponized hind legs in fights with other males over access to territories (Emberts and Wiens 2021). During these fights, males will press their tibial spines into their opponents wings, resulting in puncture wounds (B). Note that wing punctures are all located in the top half of the wing (i.e. the corium). The maximum diameter of the puncture wounds shown in B (from left to right) are 0.4 mm, 1.1 mm, and 2.2 mm. Wing photos in this figure and Figures 2 and 3 were modified (i.e. had their backgrounds removed) so that the puncture wounds could more easily be visualized.



**Figure 2** Simulated fighting injuries increase an individual's resting metabolic rate (RMR). The RMR of experimentally damaged individuals increased, whereas the RMR of non-injured individuals (i.e. baseline) slightly decreased. The dashed line indicates where an individual (i.e. a datapoint) would be if they had the exact same resting metabolic rate both before and after they received their treatment. Rectangles indicate a treatment's mean value, while error bars visualize 95% confidence intervals. Note that the 95% confidence intervals for our baseline treatment include zero, while the 95% confidence intervals for our experimental manipulation treatment exclude zero. Wing photographs adjacent to each rectangle visually depict the damage treatments (no damage or 3 mm punctures).



**Figure 3** Fighting-related damage, and the severity of that damage, influence an individual's flying performance. We found that *Thasus neocalifornicus* males with wing damage flew significantly shorter distances than non-damaged individuals. Moreover, if the damage was more severe (i.e. puncture wounds were larger) the distance that individuals flew appeared to decrease. However, among damaged individuals, only individuals with 4 mm puncture wounds flew significantly shorter distances than individuals with 1 mm puncture wounds ( $z=-4.184$ ,  $P<0.001$ ; Table S4). For box and whisker plots, boxes indicate a treatment's interquartile range, while the whiskers indicate a treatment's range when excluding outliers. Wing photographs show examples of the damage treatments.