

Longitudinal parental perception of COVID-19 vaccines for children in a multi-site, cohort study

Patrick Rivers^{a,*}, Cynthia Porter^a, Lindsay B. LeClair^b, Zuha Jeddy^b, Ashley L. Fowlkes^c, Julie Mayo Lamberte^c, Katherine Herder^a, Madeleine Smith^f, Ramona Rai^b, Lauren Grant^c, Kurt T. Hegmann^f, Krystal Jovel^a, Molly Vaughan^b, Clare Mathenge^b, Andrew L. Phillips^f, Sana Khan^a, Amadea Britton^c, Tamara Pilishvili^c, Jefferey L. Burgess^a, Gabriella Newes-Adeyi^b, Manjusha Gaglani^d, Alberto Caban-Martinez^e, Sarang Yoon^f, Karen Lutrick^a, on behalf of the PROTECT network

^a University of Arizona, Tucson, AZ, United States

^b Abt Associates, Rockville, MD, United States

^c Centers for Disease Control and Prevention, Atlanta, GA, United States

^f Utah University, Salt Lake City, UT, United States

^d Baylor College of Medicine, Temple, TX, United States

^e University of Miami, Coral Gables, FL, United States

ARTICLE INFO

Keywords:

Vaccine acceptance
COVID-19
Children

ABSTRACT

Objectives: Pediatric COVID-19 vaccine hesitancy and uptake is not well understood. Among parents of a prospective cohort of children aged 6 months–17 years, we assessed COVID-19 vaccine knowledge, attitudes, and practices (KAP), and uptake over 15 months.

Methods: The PROTECT study collected sociodemographic characteristics of children at enrollment and COVID-19 vaccination data and parental KAPs quarterly. Univariable and multivariable logistic regression models were used to test the effect of KAPs on vaccine uptake; McNemar's test for paired samples was used to evaluate KAP change over time.

Results: A total of 2,837 children were enrolled, with more than half (61 %) vaccinated by October 2022. Positive parental beliefs about vaccine safety and effectiveness strongly predicted vaccine uptake among children aged 5–11 years (aOR 13.1, 95 % CI 8.5–20.4 and aOR 6.4, 95 % CI 4.3–9.6, respectively) and children aged 12+ years (aOR 7.0, 95 % CI 3.8–13.0 and aOR 8.9, 95 % CI 4.4–18.0). Compared to enrollment, at follow-up parents (of vaccinated and unvaccinated children) reported higher self-assessed vaccine knowledge, but more negative beliefs towards vaccine safety, effectiveness, and trust in government. Parents unlikely to vaccinate their children at enrollment reported more positive beliefs on vaccine knowledge, safety, and effectiveness at follow-up.

Conclusion: The PROTECT cohort allows for an examination of factors driving vaccine uptake and how beliefs about COVID-19 and the COVID-19 vaccines change over time. Findings of the current analysis suggest that these beliefs change over time and policies aiming to increase vaccine uptake should focus on vaccine safety and effectiveness.

1. Background

Despite recommendations from the Centers for Disease Control and

Prevention's (CDC) Advisory Committee on Immunization Practices for all youth aged 6 months and older to receive COVID-19 vaccines [1–3], vaccination rates remain relatively low. Less than 10% of children 6

Abbreviations: PROTECT, Pediatric Research Observing Trends and Exposures in COVID-19 Timelines; HEROES, Arizona Healthcare, Emergency Response, and Other Essential Workers Surveillance; RECOVER, Research on the Epidemiology of SARS-CoV-2 in Essential Response Personnel; KAP, knowledge, attitudes, and practices; IRB, Institutional Review Board.

* Corresponding author at: Department of Family and Community Medicine, University of Arizona, 655 N. Alvernon Way, Tucson 85712, AZ, United States.

E-mail address: privers@arizona.edu (P. Rivers).

<https://doi.org/10.1016/j.vaccine.2024.01.016>

Received 24 May 2023; Received in revised form 13 December 2023; Accepted 2 January 2024

Available online 2 February 2024

0264-410X/© 2024 Abt Global LLC. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

months–4 years, 39% of children aged 5–11 years, and 72% of children aged 12–17 years had received at least one dose of a COVID-19 vaccine as of March 2023 [4]. There is a burgeoning literature on the knowledge, attitudes, and practices (KAP) of adults relating to COVID-19 vaccines [5–9] and their relationship to vaccine uptake; however, data for pediatric populations are limited [10]. KAP data have been previously utilized to understand COVID-19 vaccine uptake and acceptance over time among adults [6,11].

Parental hesitation of routine childhood immunizations have been linked to religious exemption, concerns about vaccine safety, and social determinants of health – such as race/ethnicity and education [5,12,13]. Misconceptions about the safety of COVID-19 vaccines as well as concerns about vaccines in general have also been identified as barriers to uptake of COVID-19 vaccines [14]. COVID-19 vaccine hesitancy may also be linked to the general antivaccine movement [12], and individual political views [15]. Unvaccinated parents have broadly reported an unwillingness to vaccinate their children, especially those less than 5 years of age, with nearly half of parents reporting they will definitely not vaccinate their young children, and one-quarter to one-third undecided [16–18]. Additional research is required to better understand COVID-19 vaccine hesitancy among parents, how vaccine perceptions evolve over time, and vaccine uptake among pediatric populations [10].

An examination of the KAPs of parents as they relate to their children can help elucidate factors associated with COVID-19 vaccine acceptability and uptake. Using data collected through prospective cohorts of children and their parents, we examined parental COVID-19 vaccine KAPs as predictors of vaccine uptake in children and assessed individual-level change in KAP responses over 15 months.

2. Methods

2.1. Study design and population

The Pediatric Research Observing Trends and Exposures in COVID-19 Timelines (PROTECT) study is a prospective cohort of children aged 6 months–17 years initiated July 2021 in Arizona, Florida, Texas, and Utah in both rural and urban sites; for this analysis, we used data collected through October 13, 2022 [19]. PROTECT is a companion study to the HEROES-RECOVER study, two cohorts of frontline workers aged ≥ 18 years: the Arizona Healthcare, Emergency Response, and Other Essential Workers Surveillance (HEROES) study and the Research on the Epidemiology of SARS-CoV-2 in Essential Response Personnel (RECOVER) study [20,21]. PROTECT recruitment included the children of HEROES-RECOVER participants and from the community. Community recruitment strategies included radio and social media advertisements, outreach at local events, and phone banking. Parents (or legal guardians) provided informed consent and children aged 7–17 years provided assent for study participation. This study protocol was reviewed and approved by the Abt Associates Institutional Review Board (IRB) (which serves as the single IRB of record for the Florida, Texas, and Utah sites and CDC), by the University of Arizona IRB for the Arizona site, and CDC.

2.2. Data collection

Upon enrollment, parents reported their child's sociodemographic information (e.g., household income, race/ethnicity, and gender), SARS-CoV-2 infection history, and medical history (e.g., medical insurance, vaccinations, medical conditions, and daily medication use).

Parents' intention to vaccinate their children with COVID-19 vaccine and KAPs were assessed through a series of questions about knowledge, attitudes towards the safety and effectiveness, trust in government information, intent to vaccinate, and perceived risk of COVID-19 infection without receiving the COVID-19 vaccine (Supplement). Responses at enrollment were considered baseline and questions were readministered every 90 days to assess change.

COVID-19 vaccination status was collected at enrollment if the child was eligible to receive the vaccine. Parents of children enrolled before they were eligible to receive the vaccine were asked as soon as the child became eligible (COVID-19 vaccines in children were approved by the U.S Food and Drug Administration under Emergency Use Authorization on: December 11, 2020 for children aged ≥ 16 years; [22] May 10, 2021 for children aged 12–15 years; [23] October 29, 2021 for children aged 5–11 years; [24] and June 17, 2022 for children aged 6 months–4 years) [25]. Parents continued to receive surveys about their child's initial series of COVID-19 vaccines until the series was completed. Vaccination was verified by parent-submitted vaccination cards, electronic medical records, or State Immunization Information Systems, as available.

2.3. Analysis

The first outcome was COVID-19 vaccine uptake, defined as the proportion of children who had received at least one COVID-19 vaccine dose at the time of analysis (October 13, 2022). Vaccine intention was derived using child's vaccination status at enrollment and parents' response to, "What are the chances that [child's first name] will get a COVID-19 vaccination?" Using responses on an 8-point Likert scale, children were grouped into three intention categories based upon parent responses and the child's vaccination status at time of enrollment: 1) *Reluctant*, defined as answers of "almost zero chance" or "very small chance," and unvaccinated; 2) *Undecided*, defined as answers of "small chance," "do not know," or "moderate chance" and unvaccinated, or 3) *Likely*, defined as answers of "large chance," "very large chance," or "almost certain," or vaccinated at the time of enrollment. Intention categories were established based on information provided at enrollment to allow comparison over time (outcome 2).

Likert scores for each KAP question were condensed into three categories, with responses greater than the midpoint indicating positive associations, midpoint responses indicating neutral associations without direction, and responses below the midpoint indicating negative associations.

2.4. Statistical analysis

General study participant sociodemographic characteristics (age category, sex, race, and ethnicity), recruitment site, previous SARS-CoV-2 infection, parent insurance status, and household annual income were stratified by child's vaccination status and then further broken down by vaccine intention group, with Pearson's chi-squared tests used to determine differences within each group. Statistical significance was based on p-values < 0.05 .

For the first objective, continuous measurements were expressed as means and standard deviations. Counts and percentages were used for categorical variables. Univariable and multivariable logistic regression models with robust standard errors [26] clustered at the household-level were used to test the effect of each KAP on vaccine uptake. Multivariable models were stratified by age group to account for differences in the timing of vaccine availability and adjusted for sociodemographic characteristics (sex, age, race, ethnicity, and household income of the participant), recruitment site, parent health insurance status, and SARS-CoV-2 infection prior to study enrollment as covariates. Statistical significance was based on 95 % confidence intervals.

To assess changes in KAP responses over time, all parents who completed KAP questions at enrollment and 9-month follow-up were included. McNemar's test for paired samples was used to evaluate the proportion individuals whose KAP responses between enrollment and 9-month assessments changed from negative/neutral to positive (positive change) or from positive to negative/neutral (negative change). For parents with multiple children enrolled in the study, stratified random sampling was used to select one child from each household. Differences in KAP responses were examined within households to ensure the validity of the sampling method. Analyses were stratified by vaccine

intention at baseline and vaccination status at follow-up. Statistical significance was based on p-values < 0.05.

All statistical analyses were completed using SAS (version 9.4; SAS Institute Cary NC).

3. Results

3.1. Participant characteristics

July 2021 to October 2022, 2,834 children from 1,712 households, an average of 1.6 children per household, (Fig. 1) were enrolled in PROTECT; including 511 children (18 %) aged 6 months–4 years, 1,529 (53 %) aged 5–11 years, 610 (22 %) aged 12–15 years, and 184 (7 %) aged 16–17 years (Table 1). Nearly half were male (n = 1,363, 48 %), 58 % (n = 1,503) were non-Hispanic White, 30 % (n = 771) were Hispanic, and 67 % (n = 1,910) resided in Arizona. Only 17 % (n = 464) reported a SARS-CoV-2 infection prior to enrollment.

Overall, 1,717 (61 %) were vaccinated with at least one dose of a COVID-19 vaccine by the time of analysis (Table 1), with 696 (40.5 %) of children with at least one dose entering the study vaccinated, and the remaining 1,021 (59.5 %) receiving their first dose after enrollment in the study. Unvaccinated children were more frequently non-Hispanic Black or Hispanic or Other, uninsured, in a home with lower household annual income, or in Texas or Florida; unvaccinated children were also more likely to have reported a SARS-CoV-2 infection prior to enrollment (all p < 0.001).

Parents of 2,709 children answered questions at enrollment about intention to vaccinate their children. Of this, 73 % (n = 1,970) were likely to vaccinate their child and 82 % (n = 1,625) of their children had received at least one dose of a COVID-19 vaccine by the time of analysis in October 2022 (Table 2). Of the 415 (15 %) parents undecided about vaccinating their child at enrollment, 20 % (n = 84) had received at least one dose at follow-up and only 8 (2 %) of the 324 (12 %) parents unlikely to vaccinate their child were vaccinated.

Among children of parents likely to vaccinate at enrollment, children residing in Texas, Florida, and Utah, or in a household with a lower income, with children aged <5 years, uninsured, or non-Hispanic Black and Hispanic were less frequently vaccinated (p < 0.001). Similarly, children of parents likely to vaccinate at enrollment, but who reported a

SARS-CoV-2 infection prior to enrollment, were more likely to be unvaccinated at time of analysis (p = 0.02).

3.2. KAP as predictor for vaccine uptake

In adjusted models, beliefs regarding vaccine safety, vaccine effectiveness, trust in government information about the vaccine, and chances of getting sick without the vaccine strongly predicted vaccine uptake in children aged 5 years and older. Parents of children aged 12 and older with positive beliefs towards COVID-19 vaccine safety were 7 times more likely to vaccinate their child (aOR 7.0, 95 % CI 3.8–13.0) compared to parents responding neutrally and nearly 9 times more likely if reporting positive beliefs about COVID-19 vaccine effectiveness (aOR 8.9, 95 % CI 4.4–18.0, Table 3a). Parents of children in this group who believed their child had higher chances of getting sick without a COVID-19 vaccine were over 8 times more likely to vaccinate their child (aOR 8.4, 95 % CI 4.0–17.7) compared to parents responding neutrally, and nearly 5 times more likely if reporting more trust in government information about the COVID-19 vaccines (aOR 4.7, 95 % CI 2.6–8.6).

Parents of children aged 5–11 years with positive beliefs in COVID-19 vaccine safety were 13 times more likely to vaccinate their child (aOR 13.1, 95 % CI 8.5–20.4, Table 3b) than parents responding neutrally, and more than 6 times more likely to vaccinate their child if reporting positively about COVID-19 vaccine effectiveness (aOR 6.4, 95 % CI 4.3–9.6). Parents of children in this age group who believed their child had higher chances of getting sick without a COVID-19 vaccine were 3 times more likely to vaccinate their child (aOR 3.0, 95 % CI 1.9–4.6) compared to parents responding neutrally, and over 7 times more likely if reporting more trust in government information about the COVID-19 vaccines (aOR 7.4, 95 % CI 4.7–11.9). Parents of children in this group were twice as likely to vaccinate their child if they reported positively about their vaccine knowledge compared to parents responding neutrally (aOR 2.1, 95 % CI 1.4–3.1).

A smaller number of vaccinated individuals in the under 5 years group led to model convergence issues, though similar trends held in the unadjusted models for both negative and positive KAPs (Table 3c).




Fig. 1. Flow diagram of PROTECT study and KAP analytic populations.

Table 1
 Characteristics of children aged 6 months and older in PROTECT^a, July 2021 to October 2022, stratified by COVID-19 vaccination status (n = 2834).

Characteristic	Vaccinated ^b		Unvaccinated		p-value ^c	Total (n = 2834)	
	(n = 1717)		(n = 1117)			No.	Col %
	No.	Col %	No.	Col %		No.	Col %
Recruitment site					<0.001*		
Tucson, AZ	920	53.6 %	349	31.2 %		1269	44.8 %
Phoenix, AZ	267	15.6 %	114	10.2 %		381	13.4 %
Other AZ sites	155	9.0 %	105	9.4 %		260	9.2 %
BSWH, ^d TX	85	5.0 %	131	11.7 %		216	7.6 %
Salt Lake City, UT	156	9.1 %	131	11.7 %		287	10.1 %
Miami, FL	134	7.8 %	287	25.7 %		421	14.9 %
Age at Enrollment					0.321		
<5 years	145	8.4 %	366	32.8 %		511	18.0 %
5–11 years	1023	59.6 %	506	45.3 %		1529	54.0 %
12+ years	549	32.0 %	245	21.9 %		794	28.0 %
Sex^e					<0.001*		
Male	832	48.5 %	530	47.4 %		1362	48.1 %
Female	861	50.1 %	492	44.0 %		1353	47.7 %
Missing	24	1.4 %	95	8.5 %		119	4.2 %
Race/Ethnicity^f					<0.001*		
NH White ^f	1004	58.5 %	499	44.7 %		1503	53.0 %
NH Black	29	1.7 %	47	4.2 %		76	2.7 %
NH Asian	53	3.1 %	23	2.1 %		76	2.7 %
Hispanic	431	25.1 %	340	30.4 %		771	27.2 %
Other	131	7.6 %	59	5.3 %		190	6.7 %
Missing	69	4.0 %	149	13.3 %		218	7.7 %
SARS-CoV-2 Infection Prior to Enrollment					<0.001*		
Yes	248	14.4 %	216	12.6 %		464	16.4 %
No	1469	85.6 %	865	50.4 %		2327	82.1 %
Parent Insured					<0.001*		
Yes	1644	95.7 %	893	52.0 %		2537	89.5 %
No	45	2.6 %	111	6.5 %		156	5.5 %
Missing	28	1.6 %	113	6.6 %		141	5.0 %
Household Annual Income					<0.001*		
\$0–49,999	121	7.0 %	185	10.8 %		306	10.8 %
\$50,000–\$99,999	383	22.3 %	307	17.9 %		690	24.3 %
\$100,000–\$149,999	437	25.5 %	185	16.6 %		622	21.9 %
\$150,000+	608	35.4 %	222	19.9 %		830	29.3 %
Missing	168	9.8 %	218	19.5 %		386	13.6 %

* Denotes statistically significant p-value.

^a PROTECT = Pediatric Research Observing Trends and Exposures in COVID-19 Timelines.

^b Child received at least one dose of a COVID-19 vaccine.

^c Pearson's chi-squared test used to generate p-values.

^d Baylor Scott & White Hospital.

^e Assigned at birth.

^f NH = Non-Hispanic.

3.3. Individual-level change in KAPs over time

A total of 1,134 PROTECT parents completed both enrollment and 9-month follow-up surveys and were included in the individual-level change over time analysis (Fig. 2). One hundred and eighteen (17 %) parents of children who were vaccinated at enrollment and 57 (15 %) parents of unvaccinated children changed responses regarding COVID-19 vaccine knowledge from a negative response at enrollment to a positive response at follow up (p < 0.0001, and p = 0.02 respectively) (Table 4a). A larger proportion of parents changed from a positive response at enrollment to negative at follow up regarding COVID-19 vaccine effectiveness among vaccinated (n = 312, 44 %, p < 0.001) and unvaccinated (n = 87, 23 %, p = 0.001) children. About a quarter of parents of vaccinated (n = 175, 25 %) and unvaccinated (n = 85, 23 %) children with positive responses at enrollment toward trust in government information about COVID-19 vaccines changed to negative at follow up (p < 0.001). One hundred and thirty-one (19 %) parents of vaccinated children also changed from a positive response at enrollment to a negative response at 9 months reporting their views about COVID-19 vaccine safety (p = 0.003). The KAPs of the majority of parents did not change over time, with 18 %–35 % of parents of unvaccinated parents changing, and 23 %–49 % of parents of vaccinated parents changing individual KAPs.

When analyzing the changes in KAPs from enrollment to the 9-month follow-up by vaccine intention group, a significant percentage of parents unlikely to vaccinate their children changed from a negative response at enrollment to a positive response at follow up, including 25 (20 %) responding positively towards COVID-19 vaccine knowledge (p = 0.03), 19 (16 %) towards COVID-19 vaccine safety (p = 0.007), and 27 (22 %) towards COVID-19 vaccine effectiveness (p = 0.004) (Table 4b). Among parents undecided about vaccination, 28 (17 %) changed from providing negative responses at enrollment to positive responses at follow up to questions on COVID-19 vaccine knowledge (p = 0.005) and 30 (18 %) changed to positive responses on COVID-19 vaccine safety (p = 0.002); however, 32 (20 %) parents changed from providing positive responses at enrollment to negative at follow up on questions regarding trust in government information about COVID-19 vaccines (p = 0.011). Parents likely to vaccinate their children changed from negative at enrollment to positive at follow up on COVID-19 vaccine knowledge (n = 134, 16 %, p < 0.001), but a larger proportion changed from positive at enrollment to negative at follow up on COVID-19 vaccine safety (n = 172, 21 %), COVID-19 vaccine effectiveness (n = 369, 45 %), and trust in government information about COVID-19 vaccines (n = 213, 27 %, all p < 0.001). Overall, between 20 % and 67 % of individual parent KAPs changed over time in the unlikely to vaccinate group, 21 %–32 % changed in the unsure group, and 23 %–49 % in the likely to vaccinate group.

4. Discussion

The prospective nature of the PROTECT cohort enabled investigation of vaccine perception and uptake across a wide geographical area and the shifting nature of beliefs between July 2021 and October 2022, throughout youth vaccine availability. Children who were uninsured or had a lower household annual income were less likely to have received a

Table 2

Characteristics of children in PROTECT, July 2021 to October 2022, aged 6 months and older, stratified by parental intention to vaccinate at enrollment and child's vaccination status at follow-up (n = 2709).

Characteristic	Unlikely to vaccinate child (n = 324)		Undecided (n = 415)		p-value ^c	Likely to vaccinate child (n = 1970)		Total ^a (n = 2709) No. (Col %)
	Vaccinated ^b (n = 8) No. (Col %)	Unvaccinated (n = 316) No. (Col %)	Vaccinated ^b (n = 84) No. (Col %)	Unvaccinated (n = 331) No. (Col %)		Vaccinated ^b (n = 1625) No. (Col %)	Unvaccinated (n = 345) No. (Col %)	
Recruitment Site					*0.001			
Tucson, AZ	1 (12.5)	92 (29.1)	38 (45.2)	83 (25.1)		881 (54.2)	128 (37.1)	1225 (45.2)
Phoenix, AZ	1 (12.5)	34 (10.8)	21 (25.0)	35 (10.6)		245 (15.1)	34 (9.9)	371 (13.7)
Other areas in AZ	2 (25.0)	39 (12.3)	5 (6.0)	36 (10.9)		148 (9.1)	22 (6.4)	252 (9.3)
BSWH, TX	0 (0.0)	43 (13.6)	8 (9.5)	44 (13.3)		77 (4.7)	39 (11.3)	211 (1.8)
Salt Lake City, UT	1 (12.5)	31 (13.0)	4 (4.8)	41 (12.4)		151 (9.3)	56 (16.2)	284 (10.5)
Miami, FL	3 (37.5)	77 (24.4)	8 (9.5)	92 (27.8)		123 (7.6)	63 (18.3)	366 (13.5)
Age					0.711			*0.001
<5 years	0 (0.0)	53 (16.8)	4 (4.8)	95 (28.7)		141 (8.7)	182 (52.8)	475 (17.5)
5–11 years	4 (50.0)	170 (53.8)	55 (65.5)	151 (45.6)		964 (59.3)	131 (38.0)	1475 (54.4)
12+ years	4 (50.0)	93 (29.4)	25 (29.8)	85 (25.7)		520 (32.0)	32 (9.3)	759 (29.9)
Sex^d					0.782			0.599
Male	6 (75.0)	170 (53.8)	40 (47.6)	167 (50.5)		786 (48.4)	177 (51.3)	1346 (49.7)
Female	2 (25.0)	145 (45.9)	44 (52.4)	163 (49.2)		815 (50.2)	166 (48.1)	1335 (49.3)
Refused	0 (0.0)	1 (0.3)	0 (0.0)	1 (0.3)		24 (1.5)	0 (0.0)	28 (1.0)
Race/Ethnicity					0.285			*0.0001
NH White ^e	2 (25.0)	182 (57.6)	39 (46.4)	158 (47.7)		963 (59.3)	154 (44.6)	1500 (57.9)
NH Black	0 (0.0)	18 (5.7)	3 (3.6)	14 (4.2)		26 (1.6)	11 (3.2)	72 (2.8)
NH Asian	0 (0.0)	5 (1.6)	2 (2.4)	11 (3.3)		51 (3.1)	7 (2.0)	76 (2.9)
Hispanic	3 (37.5)	83 (26.3)	28 (33.3)	117 (35.3)		400 (24.6)	122 (35.4)	753 (29.1)
Other	0 (0.0)	10 (3.2)	8 (9.5)	12 (3.6)		123 (7.6)	35 (10.1)	189 (7.3)
Previous COVID-19 Infection					0.055			*0.020
Yes	2 (25.0)	70 (22.2)	10 (11.9)	70 (21.1)		236 (14.5)	67 (19.4)	456 (16.9)
No	6 (75.0)	246 (77.8)	74 (88.1)	261 (78.9)		1382 (85.1)	275 (79.7)	2246 (83.1)
Parent Insured					*0.006			*<0.001
Yes	7 (87.5)	288 (91.1)	82 (97.6)	280 (84.6)		1555 (95.7)	303 (87.8)	2515 (92.8)
No	0 (0.0)	22 (7.0)	2 (2.4)	45 (13.6)		43 (2.6)	34 (9.9)	146 (5.4)
Missing	1 (12.5)	6 (1.9)	0 (0.0)	6 (1.8)		27 (1.7)	8 (2.3)	48 (1.8)
Household Annual Income					0.362			*<0.001
\$0–\$49,999	1 (12.5)	48 (15.2)	12 (14.3)	66 (19.9)		108 (6.6)	59 (17.1)	294 (12.1)
\$50,000–\$99,999	1 (12.5)	126 (39.9)	23 (27.4)	105 (31.7)		359 (22.1)	72 (20.9)	686 (28.3)
\$100,000–\$149,999	2 (25.0)	57 (18.0)	18 (21.4)	52 (15.7)		417 (25.7)	72 (20.9)	618 (25.5)
\$150,000+	3 (37.5)	56 (17.7)	20 (23.8)	68 (20.5)		585 (36.0)	95 (27.5)	829 (34.2)

^a Children with missing vaccine intention group not included.

^b Child received at least one dose of the vaccine by 6-month follow-up.

^c Differences in characteristic distributions between unvaccinated and vaccinated children within the undecided and likely intention group evaluated using Pearson's chi-square tests. P-values not calculated for the unlikely intention group given the sparse cells.

^d Assigned at birth.

^e NH = Non-Hispanic.

vaccination. The cohort was more vaccinated than the general population in all age groups, though children in younger age groups were less likely to be vaccinated. Differences in vaccination rates between this cohort and the general population for their state was particularly pronounced in Arizona (71.6 % vs. 37 %), though similar for other sites. These broad sociodemographic and age trends match those found in the general population [27]. Children who experienced a SARS-CoV-2 infection prior to enrollment were also less likely to receive a vaccination, which is consistent with other findings [28,29], and may be due to a belief in a protective effect or immunity from the previous infection. Even amongst parents likely to vaccinate their children, children under 5 years of age were less likely to be vaccinated, which may be partially explained by the shorter duration of availability.

After adjusting for sex, race/ethnicity, health insurance, household income, prior SARS-CoV-2 positivity, and site, favorable KAPs predicted vaccination. Vaccine effectiveness, vaccine safety, chances of getting sick without vaccination, and trust in government information about the vaccine were significant predictors for parents of children aged 5–11 and 12 and older. In parents of the older group, beliefs around vaccine effectiveness were the strongest predictor of child vaccination, with vaccine safety being the strongest in parents of those 5–11. These trends align with findings of parents of children <5 in this cohort [30]. Other studies have previously found vaccine intention in parents for their

children to be associated with different sociodemographic factors and medical history, including parent vaccination status, though the number of studies has been limited, and their findings have been inconsistent [31–34].

Vaccine perceptions among parents changed over the 15-month observation period. Parents who were likely to vaccinate their children felt more knowledgeable about COVID-19 vaccines 9 months post-enrollment compared to enrollment, but their beliefs about vaccine safety and effectiveness, and their level of trust in government information about the vaccines became more negative. Undecided parents were more positive about vaccine knowledge and vaccine safety, but trusted government information about the vaccines less. On the other hand, parents who were unlikely to vaccinate their child reported increased vaccine knowledge and more positive views on vaccine safety and vaccine effectiveness after 9 months.

This broad shift from more positive to negative over time may be due to most enrollment surveys being completed before the Omicron waves of COVID-19 in the United States, while most 9-month follow-up surveys were completed afterwards. It would seem likely the high rates of infection in child and adult populations would lead to more negative beliefs in several of these KAPs (e.g. beliefs relating to effectiveness and trust in government information about the vaccines), even among parents of vaccinated children. It is notable, though, that the general change

Table 3a

Knowledge, attitudes, and practices (KAPs) at enrollment of parents of children ages 12 + in the PROTECT cohort as predictors of vaccine uptake at 9-month follow-up July 2021 to October 2022 (n = 674).

	Unadjusted		Adjusted ^a	
	OR	95 % CI	OR	95 % CI
Chances of getting sick without vaccine				
Large to almost certain	6.93*	3.87–12.41	8.37*	3.97–17.67
Moderate chance	REF	REF	REF	REF
Small to almost zero	0.22*	0.12–0.39	0.23*	0.10–0.50
Vaccine knowledge				
A lot to a great deal	1.58*	1.03–2.44	1.26	0.74–2.13
Some	REF	REF	REF	REF
A little to nothing at all	0.83	0.49–1.39	1.34	0.70–2.57
Vaccine safety				
Extremely safe or very safe	7.13*	4.32–11.76	7.02*	3.78–13.01
Somewhat safe	REF	REF	REF	REF
Not too safe or not at all safe	0.08*	0.03–0.19	0.04*	0.01–0.15
Vaccine effectiveness				
Extremely effective or very effective	7.75*	4.63–12.97	8.90*	4.39–18.04
Somewhat effective	REF	REF	REF	REF
Not too effective or not at all effective	0.18*	0.09–0.36	0.17*	0.07–0.41
Trust in government information about the vaccine				
Strongly or mildly agree	7.20*	4.28–12.12	4.74*	2.61–8.61
Neutral	REF	REF	REF	REF
Strongly or mildly disagree	0.51*	0.30–0.87	0.34*	0.18–0.67

^a Adjusted for age, sex, race/ethnicity, health insurance, household income, prior COVID-19 positivity, and site.

* Statistically significant.

from positive to negative responses in the likely to vaccinate group did not appear to significantly slow vaccine uptake, and the broad change to more positive associations in the unlikely to vaccinate group suggest some positive change. This was particularly true in individuals who

Table 3b

Knowledge, attitudes, and practices (KAPs) at enrollment of parents of children ages 5–11 in the PROTECT cohort as predictors of vaccine uptake at 9-month follow-up, July 2021 to October 2022 (n = 1447).

	Unadjusted		Adjusted ^b	
	OR	95 % CI	OR	95 % CI
Chances of getting sick without vaccine				
Large to almost certain	3.96*	2.74–5.72	2.98*	1.92–4.63
Moderate chance	REF	REF	REF	REF
Small to almost zero	0.45*	0.31–0.64	0.41*	0.26–0.63
Vaccine knowledge				
A lot to a great deal	2.18*	1.54–3.09	2.06*	1.37–3.09
Some	REF	REF	REF	REF
A little to nothing at all	0.55*	0.39–0.78	0.75	0.49–1.13
Vaccine safety				
Extremely safe or very safe	11.25*	7.81–16.20	13.14*	8.47–20.40
Somewhat safe	REF	REF	REF	REF
Not too safe or not at all safe	0.08*	0.03–0.21	0.07*	0.02–0.24
Vaccine effectiveness				
Extremely effective or very effective	8.05*	5.67–11.45	6.43*	4.30–9.63
Somewhat effective	REF	REF	REF	REF
Not too effective or not at all effective	0.07*	0.03–0.18	0.04*	0.01–0.15
Trust in government information about the vaccine				
Strongly or mildly agree	8.43*	5.75–12.34	7.43*	4.65–11.88
Neutral	REF	REF	REF	REF
Strongly or mildly disagree	1.44	0.92–2.23	1.59	0.95–2.65

^b Adjusted for age, sex, race/ethnicity, health insurance, household income, prior COVID-19 positivity, and site.

* Statistically significant.

Table 3c

Knowledge, attitudes, and practices (KAPs) at enrollment of parents of children ages < 5 years in the PROTECT cohort as predictors of vaccine uptake at 9-month follow-up, July 2021 to October 2022 (n = 441).^b

	Unadjusted	
	OR	95 % CI
Chances of getting sick without vaccine		
Large to almost certain	2.90*	1.84–4.59
Moderate chance	REF	REF
Small to almost zero	0.51*	0.28–0.91
Vaccine knowledge		
A lot to a great deal	1.68*	1.05–2.68
Some	REF	REF
A little to nothing at all	0.41*	0.28–0.72
Vaccine safety		
Extremely safe or very safe	6.26*	3.65–10.75
Somewhat safe	REF	REF
Not too safe or not at all safe	N/A	N/A
Vaccine effectiveness		
Extremely effective or very effective	3.26*	2.07–5.16
Somewhat effective	REF	REF
Not too effective or not at all effective	N/A	N/A
Trust in government information about the vaccine		
Strongly or mildly agree	7.31*	3.68–14.53
Neutral	REF	REF
Strongly or mildly disagree	3.27*	1.37–7.83

^b Adjusted model failed to converge due to cell sparseness, so is omitted.

* Statistically significant.

entered the study having had at least one dose: only 5 % of this group did not have at least their initial series completed at the end of the analysis period. Therefore, increasing the likelihood of vaccination may be possible over time even among parents who had no intention of vaccinating their children when enrolled in PROTECT.

This study is subject to several limitations. First, due to enrollment and follow-up survey timing being tied to when a participant enrolled, and the fact that the follow-up surveys are not administered cross-sectionally to the entire cohort at the same time, it is difficult to draw clear conclusions about cohort-level changes over time. Secondly, some parents may have interpreted the different KAP response options (e.g., a little knowledge, or somewhat safe, or very effective) differently. Third, because approximately two-thirds of the cohort came from Arizona, there may be somewhat limited generalizability, though the other sites span a wide geographical area. Similarly, most participants were recruited from urban areas, and generalizability to non-urban individuals may be somewhat limited. In addition, the study population of families willing to complete weekly nasal swabs and complete 9 months or more of follow-up may not be generalizable to the entire population. Fifth, KAP responses for children less than 5 years of age were relatively sparse due in part to the timing of vaccine availability and time enrolled in the study, which limited the ability to examine predictors of vaccine uptake accounting for potential confounders. Finally, the surveys do not capture parents' precise reasons for changes in KAPs over time, and further investigation would be warranted to elucidate why some KAPs may change over time.

5. Conclusion

The PROTECT cohort allows for useful examinations of the factors driving vaccine uptake and how beliefs about COVID-19 and the COVID-19 vaccines change over time. Our findings suggest that many of these beliefs do change over time and policies aiming to increase vaccine uptake should focus on reinforcing messaging around the safety profiles of the vaccines and their effectiveness.

6. Disclosures

The findings and conclusions in this report are those of the authors




Fig. 2. Distribution of completed enrollment and 9-month follow-up parental KAP surveys in the PROTECT cohort by date, July 2021–September 2022.

Table 4a

Change in parental knowledge, attitude, and practice (KAP) responses of parents in regards to vaccination of their children in the PROTECT cohort during July 2021 to October 2022 by vaccine status from baseline to 9-month follow-up surveys^a (n = 1134).

n, (% of sample) at 9 months	Vaccinated 710 (62.6)				Unvaccinated 424 (37.4)			
	n	n (%) Change: Neg to Pos ^b	n (%) Change: Pos to Neg ^c	p-value ^d	n	n (%) Change: Neg to Pos	n (%) Change: Pos to Neg	p-value
Chances of Getting Sick Without Vaccine	463	90 (19.4)	83 (17.9)	0.648	359	24 (6.7)	40 (11.2)	0.059
Vaccine Knowledge	707	118 (16.7)	42 (5.9)	<0.0001*	381	57 (15.0)	33 (8.7)	0.015*
Vaccine Safety	700	78 (11.4)	131 (18.7)	0.003*	371	40 (10.8)	57 (15.4)	0.104
Vaccine Effectiveness	706	35 (5.0)	312 (44.2)	<0.001*	377	43 (11.4)	87 (23.1)	0.001*
Trust in Government information about the vaccine	695	66 (9.5)	175 (25.2)	<0.0001*	367	45 (12.3)	85 (23.2)	0.0006*

^a Responses included if parents had completed KAP questions for both enrollment and 9-month follow-up surveys.

^b Number of parents (percentage of responses) who changed from negative/neutral KAP response to positive KAP response at follow-up.

^c Number of parents (percentage of responses) who changed from positive KAP response to negative/neutral KAP response at follow-up.

^d P-value from McNemar’s exact test of matched pairs. H₀: Proportion of parents who changed from positive KAP response to negative/neutral KAP response is the same as the proportion of parents who changed from negative/neutral KAP response to positive KAP response at follow-up.

* Indicates statistically significant p-value.

and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Funding/Support

All phases of this study were funded by the Centers for Disease Control and Prevention.

CRedit authorship contribution statement

Patrick Rivers: Conceptualization, Data curation, Formal analysis, Supervision, Validation, Visualization, Writing – original draft, Writing – review & editing. **Cynthia Porter:** Data curation, Validation, Writing – review & editing. **Lindsay B. LeClair:** Conceptualization, Writing – review & editing, Project administration. **Zuha Jeddy:** Conceptualization, Writing – review & editing, Project administration. **Ashley L. Fowlkes:** Conceptualization, Project administration, Supervision,

Writing – review & editing. **Julie Mayo Lamberte:** Project administration, Writing – review & editing. **Katherine Herder:** Formal analysis, Writing – original draft, Writing – review & editing, Visualization. **Madeleine Smith:** Data curation, Writing – review & editing. **Ramona Rai:** Project administration, Writing – review & editing. **Lauren Grant:** Data curation, Formal analysis, Writing – review & editing. **Kurt T. Hegmann:** Data curation, Writing – review & editing, Project administration. **Krystal Jovel:** Data curation, Writing – review & editing. **Molly Vaughan:** Project administration, Writing – review & editing. **Clare Mathenge:** Data curation, Writing – review & editing. **Andrew L. Phillips:** Data curation, Project administration, Writing – review & editing. **Sana Khan:** Data curation, Writing – review & editing. **Amadea Britton:** Conceptualization, Investigation, Methodology, Project administration, Writing – review & editing. **Tamara Pilishvili:** Conceptualization, Investigation, Methodology, Project administration, Writing – review & editing. **Jefferey L. Burgess:** Conceptualization, Funding acquisition, Investigation, Methodology, Project

Table 4b

Change in parental knowledge, attitude, and practice responses of parents regarding vaccination of their children in the PROTECT cohort during July 2021 to October 2022 by vaccine intention from baseline to 9-month follow-up surveys^a (n = 1121).

n, % of sample at 9 months	Unlikely to vaccinate 125 (11.2)				Undecided 168 (15.0)				Likely to vaccinate 828 (73.9)			
	n	n (%) Change Neg to Pos ^b	n (%) Change Pos to Neg ^c	p- value ^d	n	n (%) Change Neg to Pos	n (%) Change Pos to Neg	p- value	n	n (%) Change Neg to Pos	n (%) Change Pos to Neg	p-value
Chances of Getting Sick Without Vaccine	117	5 (4.3)	105 (62.5)	0.99	147	17 (11.6)	14 (9.5)	0.720	576	94 (16.3)	109 (28.9)	0.326
Vaccine Knowledge	123	25 (20.3)	11 (17.7)	0.029*	165	28 (17.0)	7 (4.2)	0.005*	827	134 (16.2)	55 (6.7)	<0.001*
Vaccine Safety	118	19 (16.1)	5 (4.2)	0.007*	164	30 (18.3)	7 (4.3)	0.002*	816	76 (9.3)	172 (21.1)	<0.001*
Vaccine Effectiveness	122	27 (22.1)	9 (7.4)	0.004*	165	22 (13.3)	32 (19.4)	0.220	824	32 (3.9)	369 (44.5)	<0.001*
Trust in Government information about the vaccine	121	25 (20.7)	16 (13.2)	0.211	162	14 (8.6)	32 (19.8)	0.011*	808	74 (9.2)	213 (26.7)	<0.0001*

^a Responses included if parents had completed KAP questions for both enrollment and 9-month follow-up surveys.
^b Number of parents (percentage of responses) who changed from negative/neutral KAP response to positive KAP response at follow-up.
^c Number of parents (percentage of responses) who changed from positive KAP response to negative/neutral KAP response at follow-up.
^d P-value from McNemar’s exact test of matched pairs. H0: Proportion of parents who changed from positive KAP response to negative/neutral KAP response is the same as the proportion of parents who changed from negative/neutral KAP response to positive KAP response at follow-up.
* Indicates statistically significant p-value.

administration, Resources, Supervision, Writing – review & editing. **Gabriella Newes-Adeyi:** Project administration, Writing – review & editing. **Manjusha Gaglani:** Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Writing – review & editing. **Alberto Caban-Martinez:** Funding acquisition, Investigation, Methodology, Project administration, Writing – review & editing. **Sarang Yoon:** Funding acquisition, Investigation, Methodology, Project administration, Writing – review & editing. **Karen Lutrick:** Conceptualization, Data curation, Investigation, Methodology, Project administration, Supervision, Writing – original draft, Writing – review & editing.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Kurt T. Hegmann, MD serves as the editor-in-chief of the American College of Occupational and Environmental Medicine Guidelines; and Manjusha Gaglani, MD serves on the Texas Pediatric Society and Texas Chapter of the American Academy of Pediatrics. The other authors have no conflicts of interest to disclose.

Data availability

The authors do not have permission to share data.

Acknowledgements

CDC: Eduardo Azziz-Baumgartner, Melissa L. Arvay, Allison Ciesla, Aron Hall, Adam MacNeil, Clifford McDonald, Sue Reynolds, Ryan Wiegand, Melissa Briggs-Hagen, Morgan Najdowski, Leora Feldstein.
 Abt Associates: Danielle R. Hunt, Laura J. Edwards, Robin Bloodworth, Claire Douglas, Jazmin Duque, Jini Etolue, Deanna E. Fleary, Jessica Flores, Isaiah Gerber, Kimberly Groover, Louise Hadden, Jenna Harder, Nancy McGarry, Peenaz Mistry, Tyler C. Morrill, Kelly Patlan, Brandon P. Poe, Khaila Prather, Meghan Shea, John Thacker, Pearl Zheng.
 Baylor Scott and White Health: Kayan Dunnigan, Spencer Rose, Nicole Calhoun, Leah Odame-Bamfo, Michael E. Smith, Kempapura Murthy, Tnelda Zunie, Eric Hoffman, Martha Zayed, Ashley Graves, Joel Blais, Jason Ettlinger, Sharla Russell, Natalie Settele, Tiya Searcy, Rupande Patel, Elisa Priest, Jennifer Thomas, Muralidhar Jatla, Madhava Beeram, Javed Butler, Alejandro Arroliga.

University of Miami: Damena Gallimore-Wilson, Roger Noriega, Cynthia Beaver, Alexandra Cruz, Annabel Reyes, Brigitte Madan, Addison Testoff, John Jones.

University of Arizona: Gayatri Arani, Ariyah Armstrong, Nora Baccam, Jordan Baker, Shawn Beitel, Sienna Bigler, Shelby Capell, Andrea Carmona, Alissa Coleman, Hannah Cowling, Carly Deal, Kiara Earley, Katherine D. Ellingson, Julia Fisher, Joe K. Gerald, Lynn Gerald, Sofia Grijalva, Hanna Hanson, Olivia Healy, James Hollister, Rezwana Islam, Meccah Jarrah, Sally Littau, Amelia Lobos, Ashley Lowe, Jeremy Makar, Mayra Martinez, Flavia Nakayima Miuro, Joe Mirabito, Raul Nava, Janko Nikolich-Zugich, Assumpta Nsengiyunva, Cielo Perez, Celia Pikowski, Ferris A. Ramadan, James K. Romine, Priyanka Sharma, Alison Slocum, Lili Steffen, Xiaoxiao Sun, Nicholas Tang, Gavin Tovar, Italia Trejo, Erin Woods, April Yingst.

University of Utah: Rachel T. Brown, Camie Schaefer, Arlyne Arteaga, Matthew Bruner, Daniel Dawson, Emilee Eden, Jenna Pragastis, Joseph Stanford, Jeanmarie Mayer, Marcus Stucki, Riley Campbell, Kathy Tran, Braydon Black, Christina Pick, Madison Tallman, Chapman Cox, Derrick Wong, Michael Langston, Adriele Fugal, Fiona Tsang, Maya Wheeler, Gretchen Maughan, Megan Wilson, Pasha Stinson, Jesse Williams, Seon Reed, Jinyi Mao, Nikki Gallacher, Kendal Chatard, Jenna Vo, Katie Luong, Ryder Jordin, Grace Stewart, Brock Bourdelle, Timina Powaukee, Max Minoughan

Appendix A. Supplementary material

KAP questions asked to parents during quarterly follow-up surveys. Supplementary data to this article can be found online at <https://doi.org/10.1016/j.vaccine.2024.01.016>.

References

- [1] Fleming-Dutra KE, Wallace M, Moulia D, et al. Interim recommendations of the advisory committee on immunization practices for use of Moderna and Pfizer-BioNTech COVID-19 vaccines in children aged 6 months-5 years - United States, June 2022. *MMWR Morb Mortal Wkly Rep* 2022;71:859–68.
- [2] Woodworth K, Moulia D, Collins J, et al. The advisory committee on immunization practices’ interim recommendation for use of Pfizer-BioNTech COVID-19 vaccines in children aged 5–11 years - United States, November 2021. *MMWR Morb Mortal Wkly Rep* 2021;70:1579–83.
- [3] Wallace M, Woodworth K, Gargano J, et al. The advisory committee on immunization practices’ interim recommendation for use of Pfizer-BioNTech COVID-19 vaccine in adolescents aged 12–15 years - United States, May 2021. *MMWR Morb Mortal Wkly Rep* 2021;70:749–52.

- [4] Centers for Disease Control and Prevention. Trends in demographic characteristics of people receiving COVID-19 vaccinations in the United States. March 2, 2023; 2023.
- [5] Willis DE, Andersen JA, Bryant-Moore K, et al. COVID-19 vaccine hesitancy: race/ethnicity, trust, and fear. *Clin Transl Sci* 2021;14(6):2200–7.
- [6] Lutrick K, Groom H, Fowlkes AL, et al. COVID-19 vaccine perceptions and uptake in a national prospective cohort of essential workers. *Vaccine* 2022;40(3):494–502.
- [7] Allen JD, Matsunaga M, Lim E, Zimet GD, Nguyen KH, Fontenot HB. Parental decision making regarding COVID-19 vaccines for children under age 5: does decision self-efficacy play a role? *Vaccines (Basel)* 2023;11(2).
- [8] Szilagyi PG, Shah MD, Delgado JR, et al. Parents' intentions and perceptions about COVID-19 vaccination for their children: results from a national survey. *Pediatrics* 2021;148(4).
- [9] Ruggiero KM, Wong J, Sweeney CF, et al. Parents' intentions to vaccinate their children against COVID-19. *J Pediatr Health Care* 2021;35(5):509–17.
- [10] Willis DE, Presley J, Williams M, Zaller N, McElfish PA. COVID-19 vaccine hesitancy among youth. *Hum Vaccin Immunother* 2021;17(12):5013–5.
- [11] Akarsu B, Canbay Özdemiş D, Ayhan Baser D, Aksoy H, Fidancı İ, Cankurtaran M. While studies on COVID-19 vaccine is ongoing, the public's thoughts and attitudes to the future COVID-19 vaccine. *Int J Clin Pract* 2021;75(4):e13891.
- [12] Olusanya OA, Bednarczyk RA, Davis RL, Shaban-Nejad A. Addressing parental vaccine hesitancy and other barriers to childhood/adolescent vaccination uptake during the coronavirus (COVID-19) pandemic. *Front Immunol* 2021;12:663074.
- [13] Guerin RJ, Naeim A, Baxter-King R, Okun AH, Holliday D, Vavreck L. Parental intentions to vaccinate children against COVID-19: findings from a U.S. National Survey. *Vaccine* 2022.
- [14] Romer D, Winneg KM, Jamieson PE, Brensinger C, Jamieson KH. Misinformation about vaccine safety and uptake of COVID-19 vaccines among adults and 5–11-year-olds in the United States. *Vaccine* 2022;40(45):6463–70.
- [15] Albrecht D. Vaccination, politics and COVID-19 impacts. *BMC Public Health* 2022; 22(1):96.
- [16] Lopes L, Hamel L, Sparks G, Montero A, Presiado M, Brodie M. KFF COVID-19 vaccine monitor: July 2022. Kaiser Family Foundation; July 26, 2022; 2022.
- [17] Santibanez TA, Black CL, Zhou T, et al. Sociodemographic variation in early uptake of COVID-19 vaccine and parental intent and attitudes toward vaccination of children aged 6 months–4 years — United States, July 1–29, 2022. *MMWR Morb Mortal Wkly Rep* 2022;71(46):1479–84.
- [18] Santibanez TA, Lendon JP, Singleton JA, et al. Factors associated with receipt and parental intent for COVID-19 vaccination of children ages 5–11 years. *MedRxiv* 2022.
- [19] Burns J, Rivers P, LeClair LB, et al. Pediatric research observing trends and exposures in COVID-19 timelines (PROTECT): protocol for a multisite longitudinal cohort study. *JMIR Res Protoc* 2022;11(7):e37929.
- [20] Lutrick K, Ellingson KD, Baccam Z, et al. COVID-19 infection, reinfection, and vaccine effectiveness in a prospective cohort of Arizona frontline/essential workers: the AZ HEROES research protocol. *JMIR Res Protoc* 2021;10(6).
- [21] Edwards LJ, Fowlkes AL, Wesley MG, et al. Research on the epidemiology of SARS-CoV-2 in essential response personnel (RECOVER): protocol for a multisite longitudinal cohort study. *JMIR Res Protoc* 2021;10(12):e31574.
- [22] FDA Takes Key Action in Fight against COVID-19 by Issuing Emergency Use Authorization for First COVID-19 Vaccine [press release]. December 11, 2020; 2020.
- [23] Coronavirus (COVID-19) update: FDA authorizes Pfizer-BioNTech COVID-19 vaccine for emergency use in adolescents in another important action in fight against pandemic [press release]. May 10, 2021; 2021.
- [24] FDA authorizes Pfizer-BioNTech COVID-19 vaccine for emergency use in children 5 through 11 years of age [press release]. October 29, 2021; 2021.
- [25] Coronavirus (COVID-19) update: FDA authorizes Moderna and Pfizer-BioNTech COVID-19 vaccines for children down to 6 months of age [press release]. June 17, 2022; 2022.
- [26] White H. Maximum likelihood estimation of misspecified models. *Econometrica* 1982;50(1):1–25.
- [27] American Academy of Pediatrics. Children and COVID-19 vaccination trends. <https://www.aap.org/en/pages/2019-novel-coronavirus-covid-infections/children-and-covid-19-vaccination-trends/>. Published 2022. Accessed.
- [28] Do DP, Frank R. Prior COVID-19 infection: an underappreciated factor in vaccine hesitancy in the USA. *J Public Health (Oxf)* 2022;44(2):471–4.
- [29] Nguyen KH, Huang J, Mansfield K, Corlin L, Allen JD. COVID-19 vaccination coverage, behaviors, and intentions among adults with previous diagnosis, United States. *Emerg Infect Dis* 2022;28(3):631–8.
- [30] Lutrick K, Fowlkes A, Rivers P, et al. Parental intentions and perceptions toward COVID-19 vaccination among children aged 4 months to 4 years - PROTECT cohort, four states, July 2021-May 2022. *MMWR Morb Mortal Wkly Rep* 2022;71(35):1109–14.
- [31] Phan TT, Enlow PT, Wong MK, Lewis AM, Kazak AE, Miller JM. Medical factors associated with caregiver intention to vaccinate their children against COVID-19. *Vaccine X* 2022;10:100144.
- [32] Szilagyi PG, Albertin CS, Gurfinkel D, et al. Prevalence and characteristics of HPV vaccine hesitancy among parents of adolescents across the US. *Vaccine* 2020;38(38):6027–37.
- [33] Opel DJ, Mangione-Smith R, Taylor JA, et al. Development of a survey to identify vaccine-hesitant parents: the parent attitudes about childhood vaccines survey. *Hum Vaccin* 2011;7(4):419–25.
- [34] Kempe A, Saville AW, Albertin C, et al. Parental hesitancy about routine childhood and influenza vaccinations: a national survey. *Pediatrics* 2020;146(1).